

# **Taylor Polynomials: Examples: Part 2**

Created by

Barbara Forrest and Brian Forrest

# Taylor Polynomials

---

## Definition: [Taylor Polynomials]

Assume that  $f(x)$  is  $n$ -times differentiable at  $x = a$ . The  $n$ -th degree Taylor polynomial for  $f(x)$  centered at  $x = a$  is the polynomial

$$\begin{aligned}T_{n,a}(x) &= \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k \\ &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \\ &\quad \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n\end{aligned}$$

## Taylor Polynomials for $\sin(x)$

---

**Example 2:** Find all of the Taylor polynomials up to degree 5 for the function  $f(x) = \sin(x)$  with center  $x = 0$ .

**Solution :** We can see that

$$\begin{aligned}f(0) &= \sin(0) = 0, \\f'(0) &= \cos(0) = 1, \\f''(0) &= -\sin(0) = 0, \\f'''(0) &= -\cos(0) = -1, \\f^{(4)}(0) &= \sin(0) = 0, \text{ and} \\f^{(5)}(0) &= \cos(0) = 1.\end{aligned}$$

Therefore

$$T_{0,0}(x) = 0,$$

$$T_{1,0}(x) = L_0(x) = 0 + 1(x - 0) = x$$

and

$$\begin{aligned}T_{2,0}(x) &= 0 + 1(x - 0) + \frac{0}{2!}(x - 0)^2 \\&= x \\&= T_{1,0}(x).\end{aligned}$$

## Taylor Polynomials for $\sin(x)$

---

**Example 2 (continued):** Find all of the Taylor polynomials up to degree 5 for the function  $f(x) = \sin(x)$  with center  $x = 0$ .

**Solution (continued):** Recall that  $f(0) = \sin(0) = 0$ ,  
 $f'(0) = \cos(0) = 1$ ,  $f''(0) = -\sin(0) = 0$ ,  
 $f'''(0) = -\cos(0) = -1$ ,  $f^{(4)}(0) = \sin(0) = 0$ , and  
 $f^{(5)}(0) = \cos(0) = 1$ .


Next we have

$$\begin{aligned}T_{3,0}(x) &= 0 + 1(x - 0) + \frac{0}{2!}(x - 0)^2 + \frac{-1}{3!}(x - 0)^3 \\ &= x - \frac{x^3}{6} \\ &= T_{4,0}(x)\end{aligned}$$

Finally,

$$T_{5,0}(x) = T_{4,0}(x) + \frac{x^5}{5!} = x - \frac{x^3}{6} + \frac{x^5}{120}$$

# Taylor Polynomials for $\sin(x)$



**Note:** The diagram displays the graph of  $\sin(x)$  with its Taylor Polynomials up to degree 5 (excluding  $T_{0,0}(x)$  since its graph is the  $x$ -axis).

For  $k \geq 0$

$$T_{2k+1,0}(x) = T_{2k+2,0}(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} + \cdots + (-1)^k \frac{x^{2k+1}}{(2k+1)!}$$


# Taylor Polynomials for $\sin(x)$

**Question:** How accurate is the approximation

$$\sin(x) \cong T_{13,0}(x)$$

where

$$\begin{aligned} T_{13,0}(x) = & x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 \\ & + \frac{1}{362880}x^9 - \frac{1}{39916800}x^{11} + \frac{1}{6227020800}x^{13} \end{aligned}$$



**Note:** If  $x \in [-1, 1]$ , then

$$|\sin(x) - T_{13,0}(x)| < 10^{-12}$$

while for  $x \in [-0.01, 0.01]$ ,

$$|\sin(x) - T_{13,0}(x)| < 10^{-42}.$$

# Taylor Polynomials for $e^x$

---

**Example:** Let  $f(x) = e^x$ . Then

$$f^{(k)}(x) = e^x \Rightarrow f^{(k)}(0) = e^0 = 1$$

for any  $k$ . Therefore

$$\begin{aligned} T_{n,0}(x) &= \sum_{k=0}^n \frac{f^{(k)}(0)}{k!} (x-0)^k \\ &= \sum_{k=0}^n \frac{e^0}{k!} x^k \\ &= \sum_{k=0}^n \frac{x^k}{k!}. \end{aligned}$$

In particular,

$$T_{0,0}(x) = 1,$$

$$T_{1,0}(x) = 1 + x,$$


$$T_{2,0}(x) = 1 + x + \frac{x^2}{2},$$

$$T_{3,0}(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6},$$

$$T_{4,0}(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24},$$

$$T_{5,0}(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120}.$$

# Taylor Polynomials for $e^x$



$$T_{n,0}(x) = \sum_{k=0}^n \frac{x^{(k)}}{k!} = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \cdots + \frac{x^n}{n!}$$