

Taylor Polynomials: Examples: Part 2

Created by

Barbara Forrest and Brian Forrest

Taylor Polynomials

Definition: [Taylor Polynomials]

Assume that $f(x)$ is n -times differentiable at $x = a$. The n -th degree Taylor polynomial for $f(x)$ centered at $x = a$ is the polynomial

$$\begin{aligned}T_{n,a}(x) &= \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x - a)^k \\ &= f(a) + f'(a)(x - a) + \frac{f''(a)}{2!} (x - a)^2 + \\ &\quad \dots + \frac{f^{(n)}(a)}{n!} (x - a)^n\end{aligned}$$

Taylor Polynomials for $\sin(x)$

Example 2: Find all of the Taylor polynomials up to degree 5 for the function $f(x) = \sin(x)$ with center $x = 0$.

Solution : We can see that

$$\begin{aligned}f(0) &= \sin(0) = 0, \\f'(0) &= \cos(0) = 1, \\f''(0) &= -\sin(0) = 0, \\f'''(0) &= -\cos(0) = -1, \\f^{(4)}(0) &= \sin(0) = 0, \text{ and} \\f^{(5)}(0) &= \cos(0) = 1.\end{aligned}$$

Therefore

$$\begin{aligned}T_{0,0}(x) &= 0, \\T_{1,0}(x) &= L_0(x) = 0 + 1(x - 0) = x\end{aligned}$$

and

$$\begin{aligned}T_{2,0}(x) &= 0 + 1(x - 0) + \frac{0}{2!}(x - 0)^2 \\&= x \\&= T_{1,0}(x).\end{aligned}$$

Taylor Polynomials for $\sin(x)$

Example 2 (continued): Find all of the Taylor polynomials up to degree 5 for the function $f(x) = \sin(x)$ with center $x = 0$.

Solution (continued): Recall that $f(0) = \sin(0) = 0$,
 $f'(0) = \cos(0) = 1$, $f''(0) = -\sin(0) = 0$,
 $f'''(0) = -\cos(0) = -1$, $f^{(4)}(0) = \sin(0) = 0$, and
 $f^{(5)}(0) = \cos(0) = 1$.


Next we have

$$\begin{aligned}T_{3,0}(x) &= 0 + 1(x - 0) + \frac{0}{2!}(x - 0)^2 + \frac{-1}{3!}(x - 0)^3 \\ &= x - \frac{x^3}{6} \\ &= T_{4,0}(x)\end{aligned}$$

Finally,

$$T_{5,0}(x) = T_{4,0}(x) + \frac{x^5}{5!} = x - \frac{x^3}{6} + \frac{x^5}{120}$$

Taylor Polynomials for $\sin(x)$



Note: The diagram displays the graph of $\sin(x)$ with its Taylor Polynomials up to degree 5 (excluding $T_{0,0}(x)$ since its graph is the x -axis).

For $k \geq 0$

$$T_{2k+1,0}(x) = T_{2k+2,0}(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} + \cdots + (-1)^k \frac{x^{2k+1}}{(2k+1)!}$$


Taylor Polynomials for $\sin(x)$

Question: How accurate is the approximation

$$\sin(x) \cong T_{13,0}(x)$$

where

$$\begin{aligned} T_{13,0}(x) = & x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 \\ & + \frac{1}{362880}x^9 - \frac{1}{39916800}x^{11} + \frac{1}{6227020800}x^{13} \end{aligned}$$



Note: If $x \in [-1, 1]$, then

$$|\sin(x) - T_{13,0}(x)| < 10^{-12}$$

while for $x \in [-0.01, 0.01]$,

$$|\sin(x) - T_{13,0}(x)| < 10^{-42}.$$

Taylor Polynomials for e^x

Example: Let $f(x) = e^x$. Then

$$f^{(k)}(x) = e^x \Rightarrow f^{(k)}(0) = e^0 = 1$$

for any k . Therefore

$$\begin{aligned} T_{n,0}(x) &= \sum_{k=0}^n \frac{f^{(k)}(0)}{k!} (x-0)^k \\ &= \sum_{k=0}^n \frac{e^0}{k!} x^k \\ &= \sum_{k=0}^n \frac{x^k}{k!}. \end{aligned}$$

In particular,

$$T_{0,0}(x) = 1,$$

$$T_{1,0}(x) = 1 + x,$$


$$T_{2,0}(x) = 1 + x + \frac{x^2}{2},$$

$$T_{3,0}(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6},$$

$$T_{4,0}(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24},$$

$$T_{5,0}(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120}.$$

Taylor Polynomials for e^x



$$T_{n,0}(x) = \sum_{k=0}^n \frac{x^{(k)}}{k!} = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \cdots + \frac{x^n}{n!}$$