

# Bounded low and high sets

Bernard A. Anderson  
Department of Mathematics and Physical Sciences  
Gordon State College  
419 College Drive  
Barnesville, GA 30204, United States  
banderson@gordonstate.edu  
faculty.gordonstate.edu/banderson/

Barbara F. Csima  
Department of Pure Mathematics  
University of Waterloo,  
Waterloo, ON N2L 3G1, Canada  
csima@uwaterloo.ca  
www.math.uwaterloo.ca/~csima

Karen M. Lange  
Department of Mathematics  
Wellesley College  
106 Central St.  
Wellesley, MA 02481, United States  
karen.lange@wellesley.edu  
palmer.wellesley.edu/~klange/

August 21, 2015

## Abstract

Anderson and Csima [1] defined a jump operator, the *bounded jump*, with respect to bounded Turing (or weak truth table) reducibility. They showed that the bounded jump is closely related to the Ershov hierarchy and that it satisfies an analogue of Shoenfield jump inversion. We show that there are high bounded low sets and low bounded high sets. Thus, the information coded in the bounded jump is quite different from that of the standard jump. We also consider whether the analogue of the Jump Theorem holds for the bounded jump; do we have  $A \leq_{bT} B$  if and only if  $A^b \leq_1 B^b$ ? We show the forward direction holds but not the reverse.

# 1 Introduction

There are many models of computation based on restrictions of Turing reducibility  $\leq_T$ . Here we consider *bounded Turing reducibility* (also known as *weak truth table reducibility*). For two sets  $A, B \subset \omega$ ,  $A$  is *bounded Turing reducible to  $B$* , written  $A \leq_{bT} B$ , if  $A \leq_T B$  and there is a computable function  $g : \omega \rightarrow \omega$  such that determining  $A(n)$  requires only consulting the first  $g(n)$  digits of  $B$ . In [1], Anderson and Csima defined an analogue of the standard jump tailored to work with  $\leq_{bT}$ . They called this operator the *bounded jump* (see Definition 1). They explored the properties of the bounded jump and compared it to other potential jump operators for weaker forms of computation. In particular, the bounded jump is strictly increasing and order preserving on the bounded Turing degrees, just as the standard jump is on the Turing degrees. They also showed that the bounded jump hierarchy is closely connected to the Ershov hierarchy of  $\alpha$ -c.e. sets (see Theorem 1.2). Here we compare bounded low and bounded high sets to their standard jump counterparts. Recall that a set is *low* (respectively *high*) if the set's standard jump has the minimum (respectively maximum) possible complexity. We can similarly define bounded low and bounded high sets. We show in Theorems 2.1 and 2.5 that there are low bounded high sets and high bounded low sets. Hence, the amount of information coded into the bounded jump does not help us understand the complexity of the standard jump, and vice versa.

It is natural to ask which classical results hold for the bounded Turing degrees, possibly when using the bounded jump. Anderson [2], building on work of Mohrherr [9], proved the analogue of Friedberg's jump inversion in the truth table degrees: for all  $X \geq_{tt} \emptyset'$ , there exists a set  $A$  such that  $A' \equiv_{tt} X \equiv_{tt} A \oplus \emptyset'$ . Moreover, the same proof gives the result in the bounded Turing degrees. However, Csima, Downey, and Ng [5] showed that Schoenfield (and hence Sacks) jump inversion fails when the standard jump is used. Anderson and Csima obtained the analogue with the bounded jump: given a set  $B$  such that  $\emptyset^b \leq_{bT} B \leq_{bT} \emptyset^{2b}$ , there is a set  $A \leq_{bT} \emptyset^b$  such that  $A^b \equiv_{bT} B$ . They leave unanswered whether  $A$  can be taken to be computably enumerable, i.e., whether the analogue of Sacks jump inversion holds. In §3, we consider the classical Jump Theorem, which states that for any sets  $A$  and  $B$ , we have  $A \leq_T B \Leftrightarrow A' \leq_1 B'$ . We wish to determine whether the analogue of the Jump Theorem holds for the bounded jump; do we have  $A \leq_{bT} B \Leftrightarrow A^b \leq_1 B^b$ ? We find that the forward direction holds, but the converse does not.

## 1.1 Notation and Definitions

For notation and background not described below, see Cooper [4] and Soare [12] [11]. All sets considered are subsets of  $\omega$ . We let  $\varphi_0, \varphi_1, \varphi_2, \dots$  be an effective enumeration of the partial computable functions and  $\Phi_0, \Phi_1, \Phi_2, \dots$  be an effective enumeration of the Turing functionals. We assume our enumerations are acceptable.

We let  $\emptyset' = \{x \mid \varphi_x(x) \downarrow\}$ , and for an arbitrary set  $A$ , let  $A' = \{x \mid \Phi_x^A(x) \downarrow\}$ . In the case that the enumeration  $\{\varphi_n\}_{n \in \omega}$  is such that  $\varphi_n = \Phi_n^\emptyset$ , there is no confusion with the two definitions of  $\emptyset'$ . But under any enumeration, the two definitions are 1-equivalent.

For a set  $A$ , we let  $A \restriction x = \{n \in A \mid n \leq x\}$ . We follow an expression with a stage number in brackets (i.e.  $[s]$ ) to indicate the stage number applies to everything in the

expression that is indexed by stage.

For sets  $A$  and  $B$  we write that  $A \leq_{bT} B$ , and say  $A$  is *bounded Turing reducible* to  $B$ , if there exist  $i$  and  $j$  such that  $\varphi_j$  is total and for all  $x$ ,  $A(x) = \Phi_i^{B \parallel \varphi_j(x)}(x) \downarrow$ .

**Definition 1.** Given a set  $A$ , the *bounded jump* of  $A$ , is

$$A^b = \{x \in \omega \mid (\exists i \leq x)[\varphi_i(x) \downarrow \wedge \Phi_x^{A \parallel \varphi_i(x)}(x) \downarrow]\}.$$

We let  $A^{nb}$  denote the  $n$ -th bounded jump. We list a few basic properties of the bounded jump.

**Lemma 1.1** (Anderson & Csimá [1]). *For all sets  $A$ , we have  $A \leq_1 A^b \leq_1 A'$  and  $A^b \equiv_T A \oplus \emptyset'$ . Moreover,  $\emptyset^b \equiv_1 \emptyset'$ .*

As mentioned earlier, a set is low (high) with respect to a given jump operator if its jump encodes the least (most) possible information. A set  $A$  is *low* if  $A' \leq_T \emptyset'$  and is *bounded low* if  $A^b \leq_{bT} \emptyset'$ . A set  $A \leq_T \emptyset'$  is *high* if  $A' \geq_T \emptyset''$ , and a set  $A \leq_{bT} \emptyset^b$  is *bounded high* if  $A' \geq_{bT} \emptyset^{2b}$ .

We briefly review the  $\alpha$ -c.e. sets so that we can state Anderson and Csimá's result connecting the bounded jump hierarchy and the Ershov hierarchy. Fix a canonical, computable coding of the ordinals less than  $\omega^\omega$ . Since we do not use ordinals above  $\omega^\omega$  in this paper, the details of the coding are not significant. We say a function on an ordinal  $\alpha$  is (partial) computable if the corresponding function on codes for the ordinal  $\alpha$  is (partial) computable.

For  $\alpha \geq \omega$ , we say that a set  $A$  is  $\alpha$ -c.e. if there is a partial computable  $\psi : \omega \times \alpha \rightarrow \{0, 1\}$  such that for every  $n \in \omega$ , there exists a  $\beta < \alpha$  where  $\psi(n, \beta) \downarrow$  and  $A(n) = \psi(n, \gamma)$  where  $\gamma$  is least such that  $\psi(n, \gamma) \downarrow$  [8]. We remark that a set  $A$  is  $\omega$ -c.e. if and only if there are computable functions  $f : \omega \times \omega \rightarrow \{0, 1\}$  and  $g : \omega \rightarrow \omega$  such that  $A(n) = \lim_{s \rightarrow \infty} f(n, s)$  and  $|\{s \mid f(n, s+1) \neq f(n, s)\}| \leq g(n)$ . Thus, the above definition of  $\alpha$ -c.e. generalizes the common usage of the term “ $\omega$ -c.e.” Note that this definition differs slightly from the definitions in [3].

The following theorem generalizes the classical result that  $A \leq_{bT} \emptyset'$  iff  $A$  is  $\omega$ -c.e. Though this was not made clear in [1], the theorem holds only for notations of  $\omega^n$  for which we can computably work with the Cantor normal forms of all ordinals less than  $\omega^n$ . (In particular, the notation allows us to computably view  $\omega^n$  as the lexicographical order on  $n$ -tuples from  $\omega$ .)

**Theorem 1.2** (Anderson & Csimá [1]). *For any set  $X$  and  $n \geq 2$ ,*

$$X \leq_{bT} \emptyset^{nb} \iff X \text{ is } \omega^n\text{-c.e.} \iff X \leq_1 \emptyset^{nb}.$$

## 2 Comparing jump classes

### 2.1 Bounded Low

The next example shows that bounded low sets can code substantial information in the Turing degrees. We will need the following definitions and result (see [11]). Given

functions  $f$  and  $g$ , the function  $f$  *dominates*  $g$  if  $f(n) \geq g(n)$  for all but finitely many  $n$ ; the function  $f$  is called *dominant* if  $f$  dominates every total computable function. Martin proved that  $A$  satisfies  $\emptyset'' \leq_T A'$  iff there is a dominant function  $f \leq_T A$ . Given an infinite set  $X = \{a_0 < a_1 < a_2 < \dots\}$ , the *principal function of  $X$*  is defined such that  $p_X(n) = a_n$ .

**Theorem 2.1.** *There exists a c.e. bounded low set that is high.*

*Proof.* To ensure that  $A$  is high, we build  $A$  so that  $p_{\bar{A}}$  is dominant by satisfying for all  $i \in \omega$ :

**R<sub>i</sub>:** If  $\varphi_i$  is total, then  $(\exists m)(\forall l \geq m)[p_{\bar{A}}(l) \geq \varphi_i(l)]$ .

Next note that  $A$  is bounded low iff  $A^b \leq_{bT} \emptyset'$  iff  $A^b$  is  $\omega$ -c.e. The requirement **P<sub>x</sub>** imposes restraint on all requirements **R<sub>i</sub>** for  $i \geq x$  so that we can ensure  $A^b$  is  $\omega$ -c.e.

**P<sub>x</sub>:** If  $\varphi_{n,s}(x) \downarrow$  with  $n \leq x$ , then all requirements **R<sub>i</sub>** for  $i \geq x$  are restrained from enumerating any  $y \leq \varphi_n(x)$  into  $A$  after stage  $s$ .

In fact, the computable function  $g(x) = 2x(x+1)$  will bound the number of changes in the natural approximation of  $A^b(x)$ ,  $\Phi_x^{A \upharpoonright \varphi_n(x)}[s](x)$  where  $n \leq x$  and  $\varphi_n(x) \downarrow$ .

Finally, we ensure that  $A$  is coinfinite by satisfying for all  $e \in \omega$ :

**N<sub>e</sub>:**  $|\bar{A}| \geq e$ .

At each stage  $s$ , we have a c.e. approximation  $A_s$  of  $A$  and a current approximation of  $\bar{A}_s = \{a_{0,s} < a_{1,s} < a_{2,s} < \dots\}$ . To satisfy **N<sub>e</sub>**, no requirement **R<sub>i</sub>** for  $i \geq e$  may enumerate any element  $a_{l,s}$  into  $A$  with  $l \leq e$  at stage  $s$ . We assume, without loss of generality, for all  $i, z, s \in \omega$  that if  $\varphi_i(z+1)[s]$  converges then  $\varphi_i(z)[s]$  converges.

At stage  $s = 0$ , let  $A_0 = \emptyset$  and  $a_{j,0} = j$  for all  $j \in \omega$ .

At stage  $s + 1$ , we first calculate the restraints imposed by requirements **P<sub>x</sub>** and **N<sub>e</sub>** for each requirement **R<sub>i</sub>**.

Let  $r(x, s) = \max_{n \leq x} \varphi_n(x)[s]$ . Let

$$r_{i,s} = \max_{x \leq i} \{r(x, s), a_{i,s}\} = \max_{n \leq x \leq i} \{\varphi_n(x)[s], a_{i,s}\}. \quad (1)$$

Note that  $r(x, s) \leq r(x+1, s)$ ,  $r_{i,s} \leq r_{i,s+1}$ , and  $r_{i,s} \leq r_{i+1,s}$  for all  $x, i, s \in \omega$ . We allow requirement **R<sub>i</sub>** to enumerate any element greater than  $r_{i,s}$  into  $A$ .

We check whether any **R<sub>i</sub>** may act while obeying the restraints. We ask whether

$$(\exists i \leq s)(\exists x \leq s)[x > r_{i,s} \ \& \ \varphi_i(x)[s] \downarrow > a_{x,s}] \quad (2)$$

If so, let  $\langle i, x \rangle$  be the least pair satisfying (2). Let  $l$  be the least number such that  $a_{l,s} \geq \max_{m, y \leq s} \{\varphi_m(y)[s]\} \geq \varphi_i(x)$ . We enumerate  $a_{x,s}, a_{x+1,s}, \dots, a_{l-1,s}$  into  $A_{s+1} \supseteq A_s$  so that  $a_{x,s+1} = a_{l,s} \geq \varphi_i(x)$ . This action ends stage  $s + 1$ . This completes our description of the construction.

**Lemma 2.2.** *For all  $x, e, i \in \omega$ , the limits  $\lim_{s \rightarrow \infty} r(x, s)$ ,  $\lim_{s \rightarrow \infty} a_{e,s} = a_e$ , and  $\lim_{s \rightarrow \infty} r_{i,s} = r_i$  all exist. Hence, **N<sub>e</sub>** is satisfied for all  $e \in \omega$ , so  $A$  is coinfinite.*

*Proof.* Given  $x \in \omega$ , there exists  $s_0$  such that

$$(\forall n \leq x)[\varphi_n(x)[s_0] \downarrow \iff \varphi_n(x) \downarrow], \quad (3)$$

so  $\lim_{s \rightarrow \infty} r(x, s)$  exists.

Given  $e \in \omega$ , suppose for all  $j < e$ ,  $\lim_{s \rightarrow \infty} a_{j,s} = a_j$  is defined. We show that  $\lim_{s \rightarrow \infty} a_{e,s} = a_e$  is defined. Let  $s_0$  be a stage such that for all  $s' \geq s_0$   $a_{j,s'} = a_j$  for  $j < e$ . Hence, no elements below  $a_{e,s_0}$  are enumerated into  $A$  after stage  $s_0$ . Note that if requirement  $\mathbf{R}_i$  acts on behalf of a given pair  $\langle i, x \rangle$  as in (2), requirement  $\mathbf{R}_i$  never satisfies (2) for pair  $\langle i, x \rangle$  again. Thus, there exists a stage  $s_1 \geq s_0$  such that for all  $i \leq e$ , we have  $\varphi_i(e)[s_1] \downarrow$  iff  $\varphi_i(e) \downarrow$  and requirement  $\mathbf{R}_i$  does not act on behalf of a pair  $\langle i, x \rangle$  for  $x \leq e$  at any stage  $s \geq s_1$ . We show  $\lim_{s \rightarrow \infty} a_{e,s} = a_{e,s_1}$ . If not, we have that

$$(\exists s' \geq s_1)(\exists i' \leq s')(\exists x > r_{i',s'})[\varphi_{i'}(x)[s'] \downarrow > a_{x,s'} \text{ \& } x \leq e]. \quad (4)$$

Since  $a_{j,s} = a_{j,s_1} = a_{j,s_0}$  for all  $j < e$  and  $s \geq s_0$ , we must have that  $x = e$  in (4). Then,  $e > r_{i',s'}$  and  $\varphi_{i'}(e)[s'] \downarrow > a_{e,s'}$ . By definition,  $r_{i,s} \geq a_{i,s} \geq i$  for all  $i$  and  $s$ , so  $i' \leq e$  in (4). By construction,  $\mathbf{R}_i$  would act at stage  $s'$  for the pair  $\langle i', e \rangle$ , contradicting our choice of  $s_1$ . Hence,  $a_{e,s} = a_{e,s_1}$  for all  $s \geq s_1$ , so  $\lim_{s \rightarrow \infty} a_{e,s}$  exists.

Since  $r_{i,s} = \max_{x \leq i} \{r(x, s), a_{i,s}\}$  and  $\lim_{s \rightarrow \infty} r(x, s)$  both  $\lim_{s \rightarrow \infty} a_{i,s} = a_i$  exist,  $\lim_{s \rightarrow \infty} r_{i,s} = r_i$  exists. □

**Lemma 2.3.** *The requirement  $\mathbf{R}_i$  is satisfied for all  $i \in \omega$ . Thus,  $p_{\bar{A}}$  is a dominant function, so  $A$  is high.*

*Proof.* Suppose  $\varphi_i$  is total. We show that  $r_i = \lim_{s \rightarrow \infty} r_{i,s}$  satisfies

$$(\forall l > r_i)[a_l = \lim_{s \rightarrow \infty} a_{l,s} = p_{\bar{A}}(l) \geq \varphi_i(l)] \quad (5)$$

Let  $l > r_i$ . Let  $s_0$  be a stage such that  $r_{i,s} = r_i$  and  $a_{l,s} = a_l$  for all  $s \geq s_0$ . Since  $a_{l,s}$  never changes after stage  $s_0$ , we have that  $\mathbf{R}_i$  never acts for  $\langle i, l \rangle$  after stage  $s_0$ . Thus,  $\varphi_i(l) \leq a_l$  since  $l > r_i$ . □

**Lemma 2.4.** *The set  $A^b$  is  $\omega$ -c.e.*

*Proof.* To show that

$$A^b = \{x \in \omega \mid (\exists n \leq x)[\Phi_x^{A \parallel \varphi_n(x) \downarrow}(x) \downarrow]\} \quad (6)$$

is  $\omega$ -c.e., we construct a computable  $\{0, 1\}$ -function  $f(x, s)$  such that

1.  $\lim_{s \rightarrow \infty} f(x, s) = A^b(x)$  and
2.  $f(x, s) \neq f(x, s+1)$  for at most  $g(x) = 2x(x+1)$  many stages  $s$ .

We let

$$f(x, s) = \begin{cases} 1 & \text{if } (\exists n \leq x) [\Phi_x^{A \parallel \varphi_n(x) \downarrow}[s](x) \downarrow] \\ 0 & \text{else} \end{cases}$$

Suppose  $f(x, s) = 1$ , i.e., there is some  $n \leq x$  such that  $\Phi_x^{A \parallel \varphi_n(x) \downarrow}[s](x) \downarrow$ . By construction, the only requirements that may enumerate elements below  $\varphi_n(x)$  into  $A$  after stage  $s$  are  $\mathbf{R}_i$  for  $i < x$ . Hence, only these  $x$ -many requirements may injure the computation  $\Phi_x^{A \parallel \varphi_n(x) \downarrow}[s](x) \downarrow$ , potentially causing  $f(x, t)$  to equal 0 for some  $t > s$ . We show that each of these  $x$ -many requirements may injure a computation  $\Phi_x^{A \parallel \varphi_n(x) \downarrow}(x)$  on a given use  $\varphi_n(x)$  with  $n \leq x$  at most once.

Suppose requirement  $\mathbf{R}_k$  for  $k < x$  enumerates elements below  $\varphi_n(x)$  into  $A$  at stage  $s' > s$  while acting for the pair  $\langle k, y \rangle$ . By construction,  $\mathbf{R}_k$  enumerates  $a_{y, s'}, \dots, a_{l-1, s'}$  into  $A$  where  $a_{y, s'} < \varphi_n(x)$  and  $l$  is the least value such that

$$a_{l, s'} \geq \max_{m, z \leq s'} \{\varphi_m(z)[s']\}. \quad (7)$$

So, we have that  $a_{y, s'+1} = a_{l, s'} \geq \varphi_n(x)[s']$ .

We assumed for all  $i, z, s \in \omega$  that if  $\varphi_i(z+1)[s]$  converges then  $\varphi_i(z)[s]$  converges. It is also easy to see that the restraint  $r_{i, s}$  on  $\mathbf{R}_i$  is non-decreasing in  $s$ . Moreover, at any given stage of the construction, the least pair  $\langle i, z \rangle$  satisfying (2) receives attention. These observations imply that if requirement  $\mathbf{R}_i$  acts on behalf of the pairs  $\langle i, z \rangle$  and  $\langle i, z' \rangle$  with  $z < z'$  at stages  $s$  and  $s'$  respectively, then  $s < s'$ . Since requirement  $\mathbf{R}_k$  acts at stage  $s' > s$  for the pair  $\langle k, y \rangle$ , requirement  $\mathbf{R}_k$  will never act for a pair  $\langle k, y' \rangle$  where  $y' < y$  after stage  $s'$ . If  $\mathbf{R}_k$  acts for a pair  $\langle k, y'' \rangle$  where  $y < y''$  after stage  $s'$ , the requirement will only enumerate elements  $a$  into  $A$  at stage  $t > s'$  where

$$a \geq a_{y'', t} \geq a_{y'', s'+1} > a_{y, s'+1} \geq \varphi_n(x)[s']. \quad (8)$$

Therefore, a single requirement  $\mathbf{R}_k$  with  $k < x$  can injure a given use  $\varphi_n(x)$  for the computation  $\Phi_x^{A \parallel \varphi_n(x) \downarrow}(x)$  at most once.

There are  $(x+1)$ -many  $\varphi_n$  for  $n \leq x$  that may compute a use  $\varphi_n(x)$  for the computation  $\Phi_x^{A \parallel \varphi_n(x) \downarrow}(x)$ . Thus,  $f(x, s) \neq f(x, s+1)$  for at most  $g(x) = 2x(x+1)$  stages  $s$ , since for each of the  $(x+1)$ -many potential uses, there are  $x$ -many requirements that may injure that use at most once, and each injury may lead to two changes in  $f(x, s)$  (divergence followed by reconvergence).  $\square$

$\square$

## 2.2 Bounded high

We show a set's bounded jump may be maximally complex while its standard jump codes the minimum amount.

**Theorem 2.5.** *There exists a low set  $A \leq_{bT} \emptyset'$  that is bounded high.*

Note that this result gives an example of a low set that is not bounded low. It is also an example of an incomplete bounded high set.

*Proof.* We construct an  $\omega$ -c.e. set  $A$  (and hence  $A \leq_{bT} \emptyset'$ ) that satisfies  $A' \leq_T \emptyset'$  and  $A^b \geq_{bT} \emptyset^{2b} = (\emptyset^b)^b$ . We build a partial computable approximation function  $\theta : \omega \rightarrow \{0, 1\}$  to witness that  $A$  is  $\omega$ -c.e. Specifically, we start by having  $\theta(n, n) \downarrow$ , and the action of the construction ensures that  $A(n) = \theta(n, \tilde{n})$  where  $\tilde{n}$  is the least index such that  $\theta(n, \tilde{n}) \downarrow$ . The first condition intuitively says that  $A$  can change its guess at whether  $n \in A$  at most  $n$  times. We satisfy the standard requirements to guarantee that  $A$  is low.

$$\mathbf{N}_e: \quad (\exists^\infty s) [\Phi_e^A(e)[s] \downarrow] \implies \Phi_e^A(e) \downarrow.$$

To meet  $\mathbf{N}_e$ , we define the *restraint function*  $r(e, s) = \max_{t \leq s} u(e, t)$ , where  $u(e, s)$  is the use of the computation  $\Phi_e^A(e)[s]$  if it converges and 0 otherwise. At any stage in the construction, we do not allow any  $\mathbf{P}_i$  for  $i \geq e$  to change  $A(x)$  for any  $x < r(e, s)$ .

To show that  $A^b \geq_{bT} \emptyset^{2b}$ , we use that  $\emptyset^{2b}$  is  $\omega^2$ -c.e. by Theorem 1.2. Hence, there is a partial computable approximation function  $\psi : \omega \times \omega^2 \rightarrow \{0, 1\}$  for  $\emptyset^{2b}$ . We use  $\psi$  to produce an effective procedure  $\Psi$  and a computable function  $f$  so that  $\Psi$  computes whether  $n \in \emptyset^{2b}$  using only the oracle  $A^b \parallel f(n)$ , i.e., we satisfy the following positive requirements for all  $n \in \omega$ :

$$\mathbf{P}_n: \quad \Psi^{A^b \parallel f(n)}(n) \downarrow = \emptyset^{2b}(n).$$

We will encode  $\emptyset^{2b}$  into  $A^b \parallel f(n)$  by approximating  $\emptyset^{2b}$  with  $\psi$ , and we will define the computable bound  $f$  on the reduction  $\Psi$  in terms of how many times the approximation  $\psi$  can change. For each  $n \in \omega$ , let  $\beta_n$  be the first ordinal less than  $\omega^2$  (when dovetailing computations) for which we see  $\psi(n, \beta_n) \downarrow$ . Since  $\psi$  is a computable approximation function for  $\emptyset^{2b}$ , the function  $n \rightarrow \beta_n$  is computable. Since  $\beta_n < \omega^2$ ,  $\beta_n = \omega \cdot l_n + k_n$  for some  $l_n, k_n \in \omega$  where  $l_n$  and  $k_n$  are computable from the notation for  $\beta_n$ . Similarly, let  $\beta_{n,s}$  be the least  $\beta < \omega^2$  such that  $\psi(n, \beta)[s] \downarrow$  if such a  $\beta$  exists. Otherwise, let  $\beta_{n,s} = \beta_n$ . We then define  $l_{n,s}, k_{n,s} \in \omega$  so that  $\beta_{n,s} = \omega \cdot l_{n,s} + k_{n,s}$ . Without loss of generality, we may assume that if  $\beta_{n,s+1} \neq \beta_{n,s}$ , then  $\psi(n, \beta_{n,s+1}) \neq \psi(n, \beta_{n,s})$ .

In order to construct the reduction  $\Psi$  from  $A^b$  to  $\emptyset^{2b}$ , we will need to control whether or not infinitely many elements reside in  $A^b$ . To that end, we fix computable injective functions  $g$  and  $h$  such that for all  $x \in \omega$  we control  $\Phi_{g(x)}$  and  $\phi_{h(x)}$  where  $h(x) < g(x)$ . The procedure  $\Psi$  will only use the membership of (finitely many) indices  $g(\langle n, i, y \rangle)$  in  $A^b$  to compute  $\emptyset^{2b}(n)$ . We call this collection of indices the  $A^b$ -block of indices for  $\mathbf{P}_n$ , and we define it as follows.

The  $A^b$ -block of indices for  $\mathbf{P}_n$  consists of disjoint subblocks of indices, one for each  $i \leq l_n$ , called the *location  $i$   $A^b$ -block* for  $\mathbf{P}_n$ . The location  $i$   $A^b$ -block for  $\mathbf{P}_n$  consists of

1. A single index, called the *location index*, used to indicate whether  $l_{n,s} < i$  at some stage  $s$ ;
2. A collection of indices, called the *coding indices*, used to code  $\emptyset^{2b}(n)$  if  $i = \lim_s l_{n,s}$ ; and
3. A collection of indices, one for each coding index, called the *injury accounting indices*, used to change the functional  $\Psi$  when  $\mathbf{P}_n$  is injured. We give a computable function  $q(n)$  in Lemma 2.7 and prove there that reserving  $q(n)$  many coding indices and  $q(n)$  many injury indices suffices.

We can (uniformly in  $n$  and  $i \leq l_n$ ) computably assign  $1 + 2q(n)$  indices of the form  $g(\langle n, i, y \rangle)$  to the location  $i$   $A^b$ -block for  $\mathbf{P}_n$ . Given this assignment of indices, one can find a computable function  $f : \omega \rightarrow \omega$  such that  $f(n)$  is larger than all indices in the  $A^b$ -block for  $\mathbf{P}_n$ . Note that the  $A^b$ -block of indices for  $\mathbf{P}_n$  remains constant throughout the construction and that the  $A^b$ -blocks for  $\mathbf{P}_n$  and  $\mathbf{P}_m$  with  $n \neq m$  are disjoint.

At each stage  $s$ , for  $n \leq s$ , we define a partial function  $\alpha_{n,s}^b : \omega \rightarrow \{0, 1\}$  whose domain is the  $A^b$ -block for  $\mathbf{P}_n$  such that for  $i$  in this block,  $\alpha_{n,s}^b(i) = 0$  if  $i \notin A^b[s]$  and  $\alpha_{n,s}^b(i) = 1$  if  $i \in A^b[s]$ . We call  $\alpha_{n,s}^b$  the *stage  $s$  approximation to the  $A^b$ -block for  $\mathbf{P}_n$* . During the construction, we let  $v_{n,s}$  denote the stage  $s$  coding index for  $\emptyset^{2b}(n)$  in the location  $l_{n,s}$   $A^b$ -block for  $\mathbf{P}_n$ . To encode the stage  $s$  approximation of  $\emptyset^{2b}(n)$  into  $v_{n,s}$ , we will ask whether a (dynamic) location  $c_{n,s}$  is a member of  $A$ . We similarly define the partial function  $\alpha_{n,s} : \omega \rightarrow \{0, 1\}$  with domain  $\{c_{n,s}\}$  so that  $\alpha_{n,s}(c_{n,s}) = 0$  if  $c_{n,s} \notin A[s]$  and  $\alpha_{n,s}(c_{n,s}) = 1$  if  $c_{n,s} \in A[s]$ . During the construction, when we say “define  $\Psi^{\alpha_{n,s}^b}(n)$  to halt” (respectively “define  $\Phi_j^{\alpha_{n,s}}(x) = y$ ”), we mean that the computation holds for all oracles that agree with the stage  $s$  approximation of the  $A^b$ -block for  $\mathbf{P}_n$  (respectively the stage  $s$  approximation of  $A$  at  $c_{n,s}$ ).

In addition to coding  $\emptyset^{2b}(n)[s]$  into  $v_{n,s}$ , we will also enumerate other indices into  $A^b$  during the construction. To enumerate some index  $g(x)$  other than  $v_{n,s}$  into  $A^b$ , we will simply let  $\phi_{h(x)} \downarrow = 0$  and let  $\Phi_{g(x)}^X(g(x)) \downarrow$  for all oracles  $X$ .

We now describe a module that we will use repeatedly in the construction.

### 2.2.1 Initializing $\mathbf{P}_n$

We say we *initialize for  $\mathbf{P}_n$  at stage  $s$*  when we complete the following construction module, which encodes  $\emptyset^{2b}(n)[s]$  into  $A^b$  at location  $v_{n,s}$ . First, we let  $c_{n,s}$  be a fresh element larger than  $k_{n,s}$ ,  $\max_{e \leq s} r(e, s)$ , and all previously used elements in  $A$ .

The module depends on whether  $\mathbf{P}_n$  has been initialized previously. If  $\mathbf{P}_n$  has been initialized previously, we consider whether  $l_{n,s} = l_{n,s-1}$ . If  $l_{n,s} < l_{n,s-1}$ , for each  $l$  satisfying  $l_{n,s} < l \leq l_{n,s-1}$ , we enumerate the location index in the location  $l$   $A^b$ -block for  $\mathbf{P}_n$  into  $A^b$ , and we let  $v_{n,s}$  be the first coding index in the location  $l_{n,s}$   $A^b$ -block for  $\mathbf{P}_n$ . Now suppose  $l_{n,s} = l_{n,s-1}$ . We will see that we only initialize  $\mathbf{P}_n$  in this case when  $\mathbf{P}_n$  is injured. Hence, we enumerate into  $A^b$  the next unused injury accounting index in the location  $l_{n,s}$   $A^b$ -block for  $\mathbf{P}_n$ , and we let  $v_{n,s}$  be the next unused coding index in the location  $l_{n,s}$   $A^b$ -block for  $\mathbf{P}_n$ . Regardless of whether  $l_{n,s} = l_{n,s-1}$ , these actions destroy any previously defined computation  $\Psi^{\alpha_{n,s-1}^b}(n) \downarrow$ . (If stage  $s$  is the first stage at which we initialize  $\mathbf{P}_n$ , we have no such computations to destroy.)

In all cases, we encode  $\emptyset^{2b}(n)[s]$  into  $A^b$  at location  $v_{n,s}$ , the coding index defined above in the location  $l_{n,s}$   $A^b$ -block for  $\mathbf{P}_n$ . We enumerate  $v_{n,s}$  into  $A^b$  if and only if  $\psi(n, \beta_{n,s}) = 1$ . We accomplish this enumeration by picking  $z$  so that  $g(z) = v_{n,s}$ , setting  $\phi_{h(z)}$  equal to the constant function  $c_{n,s}$ , and defining  $\Phi_{v_{n,s}}^{\alpha_{n,s}}$  to halt on all inputs. Finally, we define

$$\Psi^{\alpha_{n,s}^b}(n) = \alpha_{n,s}^b(v_{n,s}).$$

$$\text{So, } \Psi^{\alpha_{n,s}^b}(n) = \psi(n, \beta_{n,s}).$$



### 2.2.2 The construction

At stage 0, we initialize  $\mathbf{P}_0$  for the first time. At stage  $s + 1$ , we suppose the following statements hold for  $n \leq s$ .

1.  $\Psi^{\alpha_{n,s}^b}(n) := \alpha_{n,s}^b(v_{n,s}) = \psi(n, \beta_{n,s})$ .
2. If  $v_{n,s} \in A^b[s]$ , then  $\phi_{j_{n,s}} = c_{n,s}$  provides the required oracle bound and  $c_{n,s} \notin A[s]$ . Moreover, the computation witnessing  $v_{n,s} \in A^b[s]$  only asks about the membership of  $c_{n,s}$  in  $A$ .
3.  $c_{n,s} > \max_{e \leq n} r(e, s)$ .

At stage  $s + 1$ , we consider whether we need to take action for  $\mathbf{P}_n$  for each  $n \leq s$  (in order of priority). First, we consider whether  $c_{n,s} \leq \max_{e \leq n} r(e, s + 1)$ . If so, we say that  $\mathbf{P}_n$  is *injured*, and we initialize  $\mathbf{P}_n$  at stage  $s + 1$ . If not, we consider whether  $\beta_{n,s+1} = \beta_{n,s}$ . If  $\beta_{n,s+1} = \beta_{n,s}$ , we do nothing beyond updating all values, e.g., setting  $v_{n,s+1} = v_{n,s}$ ,  $\alpha_{n,s+1}^b = \alpha_{n,s}^b$ , etc. We now suppose that  $\beta_{n,s+1} \neq \beta_{n,s}$ , so  $\psi(n, \beta_{n,s+1}) \neq \psi(n, \beta_{n,s})$  by our assumptions about  $\psi$ . Our action depends on whether  $l_{n,s+1} = l_{n,s}$ .

If  $l_{n,s+1} < l_{n,s}$ , we initialize  $\mathbf{P}_n$ . Now assume  $l_{n,s+1} = l_{n,s}$  so  $k_{n,s+1} < k_{n,s}$ . We let  $v_{n,s+1} = v_{n,s}$  and  $c_{n,s+1} = c_{n,s}$ . By our inductive hypotheses, if  $v_{n,s+1} \in A^b[s]$ , then  $\psi(n, \beta_{n,s}) = 1$  and  $\psi(n, \beta_{n,s+1}) = 0$ . Furthermore,  $c_{n,s+1} \notin A[s]$ . Suppose that  $t \leq c_{n,s+1}$  is the least element for which  $\theta(c_{n,s+1}, t)[s]$  is defined. Since  $c_{n,s+1} \notin A[s]$ , we have that  $\theta(c_{n,s+1}, t)[s] = 0$ . We define  $\theta(c_{n,s+1}, t - 1)[s + 1] = 1$ , changing the approximation of  $A$  so that  $c_{n,s+1} \in A[s + 1]$ . This action destroys the computation  $\Phi_{v_{n,s+1}}^{\alpha_{n,s}}(v_{n,s+1}) \downarrow$ , so  $v_{n,s+1} \notin A^b[s + 1]$ . Finally, we define (if it has not already been defined)  $\Psi^{\alpha_{n,s+1}^b}(n) = \alpha_{n,s+1}^b(v_{n,s+1})$ . Thus,  $\Psi^{\alpha_{n,s+1}^b}(n) = 0 = \psi(n, \beta_{n,s+1})$ .

By our inductive hypotheses, if  $v_{n,s+1} \notin A^b[s]$ , then  $\psi(n, \beta_{n,s}) = 0$  and  $\psi(n, \beta_{n,s+1}) = 1$ . If  $c_{n,s+1} \notin A[s]$ , we have not changed the approximation of  $A$  at  $c_{n,s+1}$  since  $\mathbf{P}_n$  was last initialized. In this case, we let  $A[s + 1] = A[s]$ , and we enumerate  $v_{n,s+1}$  into  $A^b$  at stage  $s + 1$  by letting  $z$  be such that  $g(z) = v_{n,s+1}$ , setting  $\phi_{h(z)}$  equal to the constant function  $c_{n,s+1}$ , and defining  $\Phi_{v_{n,s+1}}^{\alpha_{n,s+1}^b}$  to halt on all inputs. Finally, we define  $\Psi^{\alpha_{n,s+1}^b}(n) = \alpha_{n,s+1}^b(v_{n,s+1})$ . Otherwise,  $c_{n,s} \in A[s]$ , and we have changed the approximation of  $A$  at  $c_{n,s+1}$  since  $\mathbf{P}_n$  was last initialized. Suppose that  $t \leq c_{n,s+1}$  is the least element for which  $\theta(c_{n,s+1}, t)[s]$  is defined. We have that  $\theta(c_{n,s+1}, t)[s] = 1$ . We define  $\theta(c_{n,s+1}, t - 1)[s + 1] = 0$ , changing the approximation of  $A$  so that  $c_{n,s+1} \notin A[s + 1]$ . This action reinstates the previously defined computations  $\Phi_{v_{n,s+1}}^{\alpha_{n,s+1}^b}(v_{n,s+1}) \downarrow$  and  $\Psi^{\alpha_{n,s+1}^b}(n) = \alpha_{n,s+1}^b(v_{n,s+1})$ , so  $v_{n,s+1} \in A^b[s + 1]$ . Regardless of whether  $c_{n,s} \in A[s]$ , we have  $\Psi^{\alpha_{n,s+1}^b}(n) = 1 = \psi(n, \beta_{n,s+1})$  in the case  $v_{n,s+1} \notin A^b[s]$ .

The final step of stage  $s + 1$  is to initialize  $\mathbf{P}_{s+1}$ . This completes the construction.

### 2.2.3 Verification

As usual, we say requirement  $\mathbf{N}_e$  is *injured* at stage  $s + 1$  whenever some  $\mathbf{P}_n$  with  $n < e$  enacts a change  $A[s](x) \neq A[s + 1](x)$  for some  $x < r(e, s)$ . We say  $\mathbf{P}_n$  acts at stage  $s + 1$  if  $\mathbf{P}_n$  is injured at that stage or  $\beta_{n,s+1} \neq \beta_{n,s}$ .

**Lemma 2.6.** *For all  $e \in \omega$ , requirement  $\mathbf{N}_e$  is met (and is injured at most finitely often) and  $r(e) = \lim_s r(e, s)$  exists.*

*Proof.* We prove the lemma for  $\mathbf{N}_e$  by induction. By the inductive hypothesis, there is a stage  $s$  such that for all  $t \geq s$  and  $n < e$ , requirement  $\mathbf{N}_n$  is not injured at stage  $t$  and  $r(n) = r(n, t) = r(n, s)$ . Since  $\psi$  is an approximation function for the  $\omega^2$ -c.e. set  $\emptyset^{2b}$ , there is a stage  $s' > s$  such that  $\beta_{n, s'} = \beta_{n, t}$  for all  $t \geq s$  and  $n < e$ . By construction and choice of  $s'$ , no  $\mathbf{P}_n$  with  $n < e$  acts after stage  $s'$ . (Note that  $c_{n, s'} > \max_{j \leq n} r(j)$  since  $s' > s$ .) Hence,  $\mathbf{N}_e$  is never injured after stage  $s'$ . Suppose  $\Phi_e^A(e)[t'] \downarrow$  at some stage  $t' > s'$ . Then  $r(e, t) = r(e, t')$  for all  $t \geq t'$  so  $A[t]$  never changes below  $r(e, t')$  and  $\Phi_e^A(e) \downarrow$ .  $\square$

**Lemma 2.7.** *There is a computable function  $q(e)$  such that, for all  $i \in \omega$ , requirement  $\mathbf{P}_e$  is injured at most  $q(e)$  many times. Each requirement  $\mathbf{P}_e$  acts finitely often and is eventually satisfied. Moreover,  $A$  is  $\omega$ -c.e.*

*Proof.* We define  $q(e)$  in terms of two computable helper functions  $w(e)$  and  $d(e)$ , which we define inductively below. We ensure that  $w(e)$  is an upper bound on the number of times the restraint  $r(e, s)$  increases and that  $d(e)$  is an upper bound on the number of times  $\mathbf{P}_e$  redefines the dynamic index  $c_{e, s}$ . By construction,  $\mathbf{P}_e$  is injured at most  $\sum_{n=0}^e w(n)$  many times, so we define  $q(e)$  to be this sum for all  $e \in \omega$ .

We now define  $w(e)$  and  $d(e)$ . Since  $\mathbf{N}_0$  is never injured once it sets a restraint, we set  $w(0) = 1$ . For any  $e \in \omega$ , requirement  $\mathbf{P}_e$  changes the index  $c_{e, s}$  only when  $l_{e, s}$  changes or when injured by some higher priority  $\mathbf{N}_n$ . Thus, we set  $d(e) = l_e + \sum_{n=0}^e w(n)$ . Finally, we consider when  $r(e, s)$  can increase. Any  $\mathbf{P}_n$  with  $n < e$  can injure  $\mathbf{N}_e$  by changing the approximation of whether  $c_{n, s} \in A$ . The restraint  $r(e, s)$  can increase before and after each such injury. Consider a fixed collection of indices  $\{c_{n, s}\}_{n < e}$ . By construction, the approximation of  $A$  below  $\max\{c_{n, s}\}_{n < e}$  only changes on these indices. There are  $2^e$  many configurations of the values of  $A$  at these indices. Since  $r(e, s)$  is increasing,  $\mathbf{N}_e$  is injured at most  $2^e$  many times, for each fixed collection of indices  $\{c_{n, s}\}_{n < e}$ . Hence, the definition  $w(e) = 2^{e+1} \prod_{n=0}^e d(n)$  serves as the desired upper bound. This completes the definitions of  $q(e)$ ,  $d(e)$ , and  $w(e)$  and the proof of the first claim.

Now suppose all  $\mathbf{P}_n$  with  $n < e$  act finitely often and are eventually satisfied. Choose some  $s$  such that no  $\mathbf{P}_n$  with  $n < e$  nor  $\mathbf{N}_n$  with  $n \leq e$  ever acts after stage  $s$ ,  $r(n, s) = \lim_{t \rightarrow \infty} r(n, t)$  for all  $n \leq e$ , and  $\mathbf{P}_e$  is not injured after stage  $s$ . We may further choose  $s$  large enough so that  $\beta_{e, s} = \beta_{e, t}$  for all  $t \geq s$ , i.e.,  $\psi(e, \beta_{e, s}) = \emptyset^{2b}(e)$ . By the choice of  $s$ , we also have that  $v_{e, s}$ ,  $c_{e, s}$ , and  $\alpha_{e, s}^b$  never change after stage  $s$ . Moreover, the construction ensures that

$\Psi^{\alpha_{e, s}^b}(e) := \alpha_{e, s}^b(v_{e, s}) = \psi(e, \beta_{e, s})$ , which equals  $\emptyset^{2b}(e)$ . Thus,  $\Psi^{A^b \upharpoonright f(e)}(e) \downarrow = \emptyset^{2b}(e)$ .

(Recall that  $f(e)$  is larger than any element in the  $A^b$  block for  $\mathbf{P}_e$ .)

Finally, observe that we always redefine  $c_{n, s}$  whenever  $l_{n, s}$  changes and choose  $c_{n, s}$  larger than  $k_{n, s}$ . For a fixed index  $c_{n, s}$ , we redefine the approximation of whether  $c_{n, s} \in A[s]$  only when  $\beta_{n, s}$ , and hence  $k_{n, s}$ , decrease. Thus,  $A(c_{n, s})[s]$  changes at most  $k_{n, s} < c_{n, s}$  times. So,  $A$  is  $\omega$ -c.e.  $\square$

$\square$

### 3 The Jump Theorem

The Jump Theorem states in part that for any sets  $A$  and  $B$  we have  $A \leq_T B \Leftrightarrow A' \leq_1 B'$ . In Theorem 3.1, we prove the forward direction of the bounded jump analogue of the Jump Theorem. In particular, for all sets  $A$  and  $B$ , we have  $A \leq_{bT} B$  implies  $A^b \leq_1 B^b$ . We give a counterexample to the converse in Theorem 3.2. Anderson and Csima [1] showed the forward direction holds for the  $b_0$  jump, which is defined, for any set  $A$ , as

$$A^{b_0} = \{ \langle e, i, j \rangle \in \omega \mid \varphi_i(j) \downarrow \wedge \Phi_e^A \parallel \varphi_i(j) \downarrow \}.$$

The converse to that result is also false. It follows from work by Downey and Greenberg [6] that there are sets  $A$  and  $B$  such that  $A^{b_0} \leq_1 B^{b_0}$  but  $A \not\leq_{bT} B$ . (Note that Downey and Greenberg use the notation  $A^\dagger$  for a jump operator they define that satisfies  $A^\dagger \equiv_1 A^{b_0}$ .) The results for the bounded jump and the  $b_0$  jump do not follow from one another; although  $A^{b_0} \leq_1 A^b$  and  $A^b \leq_{bT} A^{b_0}$ , we do not in general have  $A^b \equiv_1 A^{b_0}$  (see [1]).

**Theorem 3.1.** *Let  $A$  and  $B$  be sets such that  $A \leq_{bT} B$ . Then  $A^b \leq_1 B^b$ .*

*Proof.* Let  $\Gamma$  and  $g$  witness that  $A \leq_{bT} B$ . Let  $j$  be an injective computable function defined by  $\varphi_{j(x,i)}(z) = g(\varphi_i(x))$  ( $z$  is a dummy variable). Let  $f$  be a computable injective function defined by

$$\Phi_{f(x)}^Y(z) \downarrow \Leftrightarrow (\exists i \leq x) [\varphi_i(x) \downarrow \wedge \Phi_x^{\Gamma^Y \parallel g(\varphi_i(x)) \parallel \varphi_i(x)}(x) \downarrow].$$

By the padding lemma, we may assume  $f(x) \geq j(x, i)$  for all  $x$  and all  $i \leq x$ . We will show that  $x \in A^b \Leftrightarrow f(x) \in B^b$  to prove the theorem.

For the forward direction we apply definitions and substitute  $n = j(x, i)$ .

$$\begin{aligned} x \in A^b &\Leftrightarrow (\exists i \leq x) [\varphi_i(x) \downarrow \wedge \Phi_x^A \parallel \varphi_i(x)(x) \downarrow] \Leftrightarrow \\ &(\exists i \leq x) [\varphi_i(x) \downarrow \wedge \Phi_x^{\Gamma^B \parallel g(\varphi_i(x)) \parallel \varphi_i(x)}(x) \downarrow] \Leftrightarrow \Phi_{f(x)}^B \parallel g(\varphi_i(x))(f(x)) \downarrow \Leftrightarrow \\ &\varphi_{j(x,i)}(f(x)) \downarrow \wedge \Phi_{f(x)}^B \parallel \varphi_{j(x,i)}(f(x))(f(x)) \downarrow \Rightarrow \\ &(\exists n \leq f(x)) [\varphi_n(f(x)) \downarrow \wedge \Phi_{f(x)}^B \parallel \varphi_n(f(x))(f(x)) \downarrow] \Leftrightarrow f(x) \in B^b \end{aligned}$$

For the backwards direction, we ignore the original witness that  $f(x) \in B^b$  and apply the use principle.

$$\begin{aligned} f(x) \in B^b &\Leftrightarrow (\exists m \leq f(x)) [\varphi_m(f(x)) \downarrow \wedge \Phi_{f(x)}^B \parallel \varphi_m(f(x))(f(x)) \downarrow] \Leftrightarrow \\ &(\exists m \leq f(x)) [\varphi_m(f(x)) \downarrow \wedge (\exists i \leq x) [\varphi_i(x) \downarrow \wedge \Phi_x^{\Gamma^B \parallel \varphi_m(f(x)) \parallel g(\varphi_i(x)) \parallel \varphi_i(x)}(x) \downarrow]] \Rightarrow \\ &(\exists i \leq x) [\varphi_i(x) \downarrow \wedge \Phi_x^{\Gamma^B \parallel g(\varphi_i(x)) \parallel \varphi_i(x)}(x) \downarrow] \Leftrightarrow \\ &(\exists i \leq x) [\varphi_i(x) \downarrow \wedge \Phi_x^A \parallel \varphi_i(x)(x) \downarrow] \Leftrightarrow x \in A^b \end{aligned}$$

Thus,  $f$  witnesses  $A^b \leq_1 B^b$ . □

**Theorem 3.2.** *There are c.e. sets  $A$  and  $B$  such that  $A^b \leq_1 B^b$  and  $A \not\leq_{bT} B$ .*

*Proof.* Let  $g$  and  $h$  be injective computable functions such that for all  $x$  and all  $k \leq 6^x$  we will control  $\Phi_{g(x)}$  and  $\varphi_{h(x,k)}$  and we have  $h(x,k) < g(x)$ .

We construct c.e. sets  $A$  and  $B$  that satisfy the following requirements:

$$\mathbf{R}_{\langle e,i \rangle} : (\exists x) [A(x) \neq \Phi_e^{B \parallel \varphi_i(x)}(x) \vee \varphi_i(x) \uparrow \vee \Phi_e^{B \parallel \varphi_i(x)}(x) \uparrow]$$

$$\mathbf{Q}_n : n \in A^b \leftrightarrow g(n) \in B^b$$

It is clear that these requirements suffice to prove the theorem. We build the sets  $A$  and  $B$  via a finite injury construction where the requirements are given priority  $\mathbf{R}_0, \mathbf{Q}_0, \mathbf{R}_1, \mathbf{Q}_1, \mathbf{R}_2, \mathbf{Q}_2, \dots$ . We will use a movable marker  $x_{e,i}$  that eventually settles on the witness for  $\mathbf{R}_{\langle e,i \rangle}$ . We use the indices  $k_n \leq 6^n$  for bookkeeping, and  $z$  represents a dummy variable throughout the construction. We will assume the convention that for any stage  $s$  and any  $c, d$  if  $\varphi_c(d)[s] \downarrow = t$  then  $t, c, d \leq s$ .

To satisfy  $\mathbf{R}_{\langle e,i \rangle}$ , we will wait until we see  $\varphi_i(x_{e,i}) \downarrow$  and  $\Phi_e^{B \parallel \varphi_i(x_{e,i})}(x_{e,i}) \downarrow = 0$  (if this never happens, we are done). We then enumerate  $x_{e,i}$  into  $A$  and attempt to preserve the now incorrect computation by fixing  $x_{e,i}$  and  $B \parallel \varphi_i(x_{e,i})$ . If this restraint is injured by a stronger requirement, we choose a new  $x_{e,i}$  and start over.

To satisfy  $\mathbf{Q}_n$ , we will wait until we see  $n$  enter our current estimate for  $A^b$ . We then define  $\varphi_{h(n,k_n)}(z)$  to be a fresh large number (the same number for all  $z$ ) and set  $\Phi_{g(n)}^{B \parallel \varphi_{h(n,k_n)}(z)}(z)$  to converge for all  $z$  so that  $g(n)$  is now in  $B^b$ . We try to fix enough of  $A$  and  $B$  to preserve the statements  $n \in A^b$  and  $g(n) \in B^b$ . If either statement is injured, we enumerate  $\varphi_{h(n,k_n)}(z)$  into  $B$  so that our earlier declaration that  $g(n)$  is in  $B^b$  no longer applies. We then increment  $k_n$  by one and start over.

We note that the computable function  $g$  is not allowed to have any errors. Hence, we may be required to void declarations placing  $g(n)$  into  $B^b$  by enumerating new elements into  $B$ , even if doing so injures stronger requirements. To prevent such injuries, we act preemptively. As soon as we see  $\varphi_i(x_{e,i})$  converge (but before we place any restraint on  $B$  for  $\mathbf{R}_{\langle e,i \rangle}$ ), we immediately void any active declarations from weaker requirements. To simplify the construction, whenever we act for a requirement, we will injure all weaker requirements, regardless of whether such an injury appears necessary. For brevity, we call this process the *injury procedure*, and we describe it at the end of the construction.

We start with  $x_{e,i}$  as the least number in the  $\langle e, i \rangle$  column and all  $k_n = 0$ . We set  $A(0) = 0$  and  $B(0) = 1$  and never change those values (this ensures our declarations for  $B^b$  do not change  $A^b$ ). No other elements begin in  $A$  and  $B$ .

At each stage  $s$  we perform the following procedures:

**Preemptive injury of  $\mathbf{R}_{\langle e,i \rangle}$**  : If some convergence  $\varphi_i(x_{e,i})[s] \downarrow$  has just occurred, then we carry out the injury procedure.

**Diagonalization for  $\mathbf{R}_{\langle e,i \rangle}$**  : If we observe that  $\Phi_e^{B \parallel \varphi_i(x_{e,i})}(x_{e,i})[s] \downarrow = 0$  holds, then we enumerate  $x_{e,i}$  into  $A$  and execute the injury procedure.

**$Q_n$  Strategy :** If  $n \in A^b[s]$  and  $Q_n$  has not acted (since its last injury), we first run the injury procedure. We then set  $\phi_{h(n,k_n)}(z)$  (for all  $z$ ) to be the least unused number greater than  $s$ , and we declare that  $\Phi_{g(n)}^{B \parallel \phi_{h(n,k_n)}(z)}(z) \downarrow$  for all  $z$ .

We describe the injury procedure on all requirements weaker than the one under consideration:

- For all weaker  $R_{\langle e,i \rangle}$  requirements with  $x_{e,i} \leq s$ , we assign a new unused value greater than  $s$  to  $x_{e,i}$ .
- For all weaker  $Q_n$  requirements, if  $\phi_{h(n,k_n)}(z)$  has been defined, we enumerate  $\phi_{h(n,k_n)}(z)$  into  $B$  and increment  $k_n$  by one.

This completes the construction. It is clear that  $A$  and  $B$  are c.e. The usual argument that the  $n^{\text{th}}$  requirement in a standard finite injury argument acts at most  $2^n$  times can be routinely altered to show that every  $Q_n$  acts at most  $6^n$  many times (we multiply by an additional  $3^n$  since the requirement  $R_{\langle e,i \rangle}$  between  $Q_m$  and  $Q_{m+1}$  may act twice). Hence, we have sufficiently many possible values for  $k_n$  to run the construction.

We show  $R_{\langle e,i \rangle}$  is satisfied, as is  $Q_n$ . Suppose that  $s_0$  is the last stage at which any requirement of higher priority than  $R_{\langle e,i \rangle}$  acts or is injured. Let  $x_{e,i}$  now denote the final value of the marker  $x_{e,i}$ , that is, its value at stage  $s_0$ . If  $\phi_i(x_{e,i})$  diverges or  $\Phi_e^{B \parallel \phi_i(x_{e,i})}(x_{e,i})$  either diverges or equals one, then we are done as  $x_{e,i} \notin A$ . Otherwise, we preemptively injure  $R_{\langle e,i \rangle}$  at some stage  $s > s_0$  and diagonalize on behalf of  $R_{\langle e,i \rangle}$  at some stage  $t > s$ . At stage  $s$ , the construction ensures that  $\phi_{h(n,k_n)}(z) > \phi_i(x_{e,i})$  for all  $Q_n$  of lower priority than  $R_{\langle e,i \rangle}$ . As a result, for all  $s_1 > s$  we have  $B_{s_1} \parallel \phi_i(x_{e,i}) = B \parallel \phi_i(x_{e,i})$ . Thus,

$$\Phi_e^{B \parallel \phi_i(x_{e,i})}(x_{e,i}) = \Phi_e^{B_t \parallel \phi_i(x_{e,i})}(x_{e,i}) = 0 \neq 1 = A(x_{e,i})$$

so  $R_{\langle e,i \rangle}$  is satisfied.

Suppose that  $s_0$  is the last stage at which any requirement of higher priority than  $Q_n$  acts or is injured, so  $k_n$  and  $h(n,k_n)$  are fixed for all  $s > s_0$ . By construction,  $\Phi_{g(n)}^{B \parallel s_0} \uparrow$  so  $g(n) \notin B^b[s_0]$ . We show  $n \in A^b \Leftrightarrow g(n) \in B^b$ . We consider two cases. First, suppose that  $Q_n$  never acts at any stage  $s > s_0$ . Then,  $n \notin A^b[s]$  for any  $s > s_0$ , so  $n \notin A^b$ . Also, no new convergence declarations are made about  $\Phi_{g(n)}^\sigma$  for any string  $\sigma \in 2^{<\omega}$  in  $B^b$  after stage  $s_0$  (and all strings  $\sigma$  for which there are existing convergence declarations are incomparable with  $B_{s_0}$ ). All lower priority  $Q_m$  requirements satisfy  $\phi_{h(m,k_m)}(z) > s_0$ , so  $B \parallel s_0[s_0] = B \parallel s_0$ . Thus,  $g(n) \notin B^b$ .

Second, suppose that  $Q_n$  acts at some stage  $s > s_0$ . Then,  $n \in A^b[s]$  and  $g(n) \in B^b[s]$ . At stage  $s$ , the injury procedure ensures that  $x_{e,i} > s$  for all lower priority  $R_{\langle e,i \rangle}$  requirements and  $\phi_{h(m,k_m)}(z) > s$  for all lower priority  $Q_m$  requirements. Hence,  $A \parallel s[s] = A \parallel s$  and  $B \parallel s[s] = B \parallel s$ . Thus,  $n \in A^b$  and  $g(n) \in B^b$ .

We conclude  $n \in A^b \Leftrightarrow g(n) \in B^b$  so  $Q_n$  is satisfied, completing our proof.  $\square$

## 4 Questions

Here we provided examples of extreme kinds of bounded low and bounded high sets. It is natural to ask what other kinds of examples exist. We constructed a c.e. high bounded low set in Theorem 2.1, but we only constructed an  $\omega$ -c.e. low bounded high set in Theorem 2.5. We conjecture that there is no c.e. example of a low set that is bounded high.

**Question 4.1.** *Does there exist a c.e. set that is both low and bounded high?*

A set  $A \leq_T \emptyset'$  is called *superlow* if  $A' \leq_{\text{tt}} \emptyset'$  and *superhigh* if  $A' \geq_{\text{tt}} \emptyset''$ . Mohrherr [10] constructed an example of an incomplete superhigh set. We have a low set that is bounded high, which is clearly bounded high and not high. Similarly, we have an example of a bounded low set that is not low.

**Question 4.2.** *Does there exist a bounded low set that is low but not superlow? Does there exist a high and bounded high set that is not superhigh?*

Observe that any superlow set is bounded low since  $A^b \leq_1 A'$ . However, we can ask:

**Question 4.3.** *Does there exist a superhigh set that is not bounded high?*

Recall that  $A$  is bounded low iff  $A^b$  is  $\omega$ -c.e. Could we provide a better characterization?

**Question 4.4.** *Provide characterizations of bounded low and bounded high sets.*

It is also natural to consider these questions for other restricted computation jump operators, such as Gerla's operator [7], which is discussed in [1].

## Acknowledgements

The second author was partially supported by Canadian NSERC Discovery Grant 312501. The third author was partially supported by National Science Foundation grants DMS-0802961 and DMS-1100604.

## References

- [1] B. Anderson and B. Csima. A bounded jump for the bounded Turing degrees. *Notre Dame J. Formal Logic*, 55(2):245–264, 2014.
- [2] B. A. Anderson. Automorphisms of the truth-table degrees are fixed on a cone. *J. Symbolic Logic*, 74(2):679–688, 2009.
- [3] C. J. Ash and J. Knight. *Computable structures and the hyperarithmetical hierarchy*, volume 144 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 2000.

- [4] B. Cooper. *Computability Theory*. Chapman and Hall/CRC, 2004.
- [5] B. Csima, R. Downey, and K. M. Ng. Limits on jump inversion for strong reducibilities. *J. Symbolic Logic*, 76(4):1287–1296, 2011.
- [6] R. Downey and N. Greenberg. A transfinite hierarchy of notions of lowness in the computably enumerable degrees, unifying classes, and natural definability. Preprint.
- [7] G. Gerla. Una generalizzazione della gerarchia di Ershov. *Bolletino U.M.I.*, 16-B(5):765–778, 1979.
- [8] C. G. Jockusch, Jr. and R. A. Shore. Pseudo-jump operators II: Transfinite iterations, hierarchies, and minimal covers. *J. Symbolic Logic*, 49:1205–1236, 1984.
- [9] J. Mohrherr. Density of a final segment of the truth-table degrees. *Pacific J. Math.*, 115(2):409–419, 1984.
- [10] J. Mohrherr. A refinement of low  $n$  and high  $n$  for the r.e. degrees. *Z. Math. Logik Grundlag. Math.*, 32(1):5–12, 1986.
- [11] R. I. Soare. *Recursively Enumerable Sets and Degrees*. Springer-Verlag, 1987.
- [12] R. I. Soare. *Computability Theory and Applications*. Springer-Verlag, preprint.