

Low-Rank Matrix Completion with Facial Reduction

Shimeng Huang¹ Henry Wolkowicz²

¹Department of Statistics and Actuarial Science
University of Waterloo

²Department of Combinatorics and Optimization
University of Waterloo

AT ECM2017, Thurs. June 1, 10:50AM

Outline

- Finding **low rank matrix completions** is a numerically hard **nonconvex** problem.
- A popular convex relaxation is the **nuclear norm** which is SDP-representable; and, both the SDP and its dual satisfy strict feasibility (Slater's constraint qualification).
- For inequality constrained optimization problems, perhaps the most important **key** is to identify the **active constraints**. We aim to do **facial reduction** for the optimal face of the SDP, i.e., identify the “**active**” face.
- Thus we (try to) **avoid a need for a SDP solver**.

Low-Rank Matrix Completion

Example (Partial Matrix with Noise — BUT Low Rank)

1.01	2	?
1	?	2.99

Low-Rank Matrix Completion

Example (Partial Matrix with Noise — BUT Low Rank)

1.01	2	?
1	?	2.99

1	2	3
1	2	3

Low-Rank Matrix Completion

Example (Partial Matrix with Noise — BUT Low Rank)

1.01	2	?
1	?	2.99

1	2	3
1	2	3

Problem Statement (non-convex & intractable)

Given a real **partial matrix** $z \in \mathbb{R}^{\hat{E}}$ with some level of noise,

$$\text{(LRMC)} \quad \min \quad \text{rank}(M) \\ \text{s.t.} \quad \| \mathcal{P}_{\hat{E}}(M) - z \| \leq \delta, \quad M \in \mathbb{R}^{m \times n}$$

- \hat{E} indices for **known entries** (sampled data) in $Z \in \mathbb{R}^{m \times n}$;
with **coordinate projection/partial matrix** $z = \mathcal{P}_{\hat{E}}(Z) \in \mathbb{R}^{\hat{E}}$
- $\delta > 0$ is a tuning parameter

Applications Include:

- data science
- model reduction
- collaborative filtering (Netflix problem)
- sensor network localization
- pattern recognition
- various machine learning scenarios

Low-Rank Matrix Completion

Minimizing rank is a hard nonconvex problem

Rank is a lower semi-continuous function.

Nuclear Norm Minimization (convex relaxation)

The problem (LRMC) can be approximated by

$$\begin{array}{ll} \text{(NN-LRMC)} & \min \quad \|M\|_* \\ & \text{s.t.} \quad \|\mathcal{P}_{\hat{E}}(M) - z\| \leq \delta \end{array}$$

- $\|M\|_* = \sum_i \sigma_i(M)$, sum of singular values, **nuclear norm** (Schatten 1-norm, Ky-Fan r -norm, trace norm)
- $\|UXV^T\|_* = \|X\|_*$ unitarily invariant

Nuclear Norm Minimization, Fazel-'02 thesis

Theorem (Fazel,Hindi,Boyd '01)

$\|X\|_*$ is the convex envelope of rank X on $\{X \in \mathbb{R}^{m \times n} : \|X\| \leq 1\}$.

Properties of nuclear norm:

- “best” convex lower approximation of rank function
- The nuclear ball is the convex hull of the intersection of rank-1 matrices with the unit ball:
 $\text{conv}\{uv^T : u \in \mathbb{R}^n, v \in \mathbb{R}^m, \|u\| = 1, \|v\| = 1\}$
- SDP-representable
- Related references by: Candes,Fazel,Parrilo,Recht

SDP Representable

SDP Embedding Lemma

Let $M \in \mathbb{R}^{m \times n}$ and $t \in \mathbb{R}$. Then:

$$\|M\|_* \leq t$$

if, and only if,

there exist (symmetric) W_1 and W_2 such that

$$\begin{bmatrix} W_1 & M \\ M^T & W_2 \end{bmatrix} \succeq 0, \quad \text{trace}(W_1) + \text{trace}(W_2) \leq 2t.$$

- compact SVD: $M = U\Sigma V^T$, $\|M\|_* = \text{trace} \Sigma \leq t$
- $\begin{bmatrix} U\Sigma^{1/2} \\ V\Sigma^{1/2} \end{bmatrix} \begin{bmatrix} U\Sigma^{1/2} \\ V\Sigma^{1/2} \end{bmatrix}^T = \begin{bmatrix} U\Sigma U^T & U\Sigma V^T \\ V\Sigma U^T & V\Sigma V^T \end{bmatrix} \succeq 0$
- For necessity, set $W_1 = U\Sigma U^T$, $W_2 = V\Sigma V^T$; for sufficiency, exploit $\text{range } M \subseteq \text{range } W_1, \text{range } M^T \subseteq \text{range } W_2$

Nuclear Norm Low Rank Problem, (NN-LRMC)

Semidefinite Embedding: Trace Minimization

Problem (NN-LRMC) can be formulated as:

$$\begin{aligned} \text{(SDP-LRMC)} \quad & \min && \frac{1}{2} \text{trace}(Y) \\ & \text{s.t.} && \|\mathcal{P}_{\bar{E}}(Y) - z\| \leq \delta \\ & && Y \succeq 0 \end{aligned}$$

where $Q = \begin{bmatrix} 0 & Z \\ Z^T & 0 \end{bmatrix}$, $z = \mathcal{P}_{\bar{E}}(Z) = \mathcal{P}_{\bar{E}}(Q)$;

\bar{E} is set of indices in Q corresponding to known entries of Z .

$$Y = \begin{array}{|c|c|} \hline \text{W1} & \text{M} \\ \hline \text{M}' & \text{W2} \\ \hline \end{array}$$

First, an Example of Facial Reduction, FR

Example (Facial Reduction in Linear Programming)

$$\begin{aligned} \min \quad & (2 \ 5 \ -1 \ 4 \ 7)x \\ \text{s.t.} \quad & \begin{bmatrix} 1 & 1 & -1 & 3 & 1 \\ -1 & 1 & 2 & 2 & -1 \end{bmatrix} x = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ & x \geq 0, x \in \mathbb{R}^5 \end{aligned}$$

First, an Example of Facial Reduction, FR

Example (Facial Reduction in Linear Programming)

$$\begin{aligned} \min \quad & (2 \ 5 \ -1 \ 4 \ 7)x \\ \text{s.t.} \quad & \begin{bmatrix} 1 & 1 & -1 & 3 & 1 \\ -1 & 1 & 2 & 2 & -1 \end{bmatrix} x = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ & x \geq 0, x \in \mathbb{R}^5 \end{aligned}$$

If we sum the two constraints we get a *facial* constraint

$$2x_2 + x_3 + 5x_4 = 0 \implies x \in \mathcal{F} = \{x \in \mathbb{R}_+^5 : x_2 = x_3 = x_4 = 0\}$$

First, an Example of Facial Reduction, FR

Example (Facial Reduction in Linear Programming)

$$\begin{aligned} \min \quad & (2 \ 5 \ -1 \ 4 \ 7) x \\ \text{s.t.} \quad & \begin{bmatrix} 1 & 1 & -1 & 3 & 1 \\ -1 & 1 & 2 & 2 & -1 \end{bmatrix} x = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ & x \geq 0, x \in \mathbb{R}^5 \end{aligned}$$

If we sum the two constraints we get a *facial* constraint

$$2x_2 + x_3 + 5x_4 = 0 \implies x \in \mathcal{F} = \{x \in \mathbb{R}_+^5 : x_2 = x_3 = x_4 = 0\}$$

strict feasibility fails; problem can be reduced

$$\begin{aligned} \min \quad & (2 \ 7) v \\ \text{s.t.} \quad & (1 \ 1) v = 1 \\ & v \geq 0 \end{aligned}$$

First Example of Facial Reduction, cont...

Example (Facial Reduction in Linear Programming)

$$\begin{array}{ll} \min & (2 \ 5 \ -1 \ 4 \ 7)x \\ \text{s.t.} & \begin{bmatrix} 1 & 1 & -1 & 3 & 1 \\ -1 & 1 & 2 & 2 & -1 \end{bmatrix} x = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ & x \geq 0 \end{array}$$

First Example of Facial Reduction, cont...

Example (Facial Reduction in Linear Programming)

$$\begin{array}{ll} \min & (2 \ 5 \ -1 \ 4 \ 7)x \\ \text{s.t.} & \begin{bmatrix} 1 & 1 & -1 & 3 & 1 \\ -1 & 1 & 2 & 2 & -1 \end{bmatrix} x = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ & x \geq 0 \end{array}$$

Find y with $y^T b = 0, 0 \neq w = A^T y \geq 0$ to get:

$$y = (1 \ 1)^T, 0 \neq w^T = (A^T y)^T = (0 \ 2 \ 1 \ 5 \ 0) \geq 0.$$

Then w is an exposing vector of the feasible set:

First Example of Facial Reduction, cont...

Example (Facial Reduction in Linear Programming)

$$\begin{array}{ll} \min & (2 \ 5 \ -1 \ 4 \ 7)x \\ \text{s.t.} & \begin{bmatrix} 1 & 1 & -1 & 3 & 1 \\ -1 & 1 & 2 & 2 & -1 \end{bmatrix} x = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ & x \geq 0 \end{array}$$

Find y with $y^T b = 0, 0 \neq w = A^T y \geq 0$ to get:

$$y = (1 \ 1)^T, 0 \neq w^T = (A^T y)^T = (0 \ 2 \ 1 \ 5 \ 0) \geq 0.$$

Then w is an exposing vector of the feasible set:

$$w^T x = 0, \forall \text{ feasible } x \implies x = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_5 \end{bmatrix}; x_2 = x_3 = x_4 = 0;$$

(simplified) FR problem is

$$\min \{(2 \ 7)v : (1 \ 1)v = 1, v \geq 0\}$$

Faces of a Closed Convex Cone, ccc

Face of a ccc \mathcal{K} , $\mathcal{K} + \mathcal{K} \subseteq \mathcal{K}$, $\mathbb{R}\mathcal{K} \subseteq \mathcal{K}$

Let \mathcal{K} be a ccc. A cone $F \subseteq \mathcal{K}$ is a **face** of \mathcal{K} , $F \trianglelefteq \mathcal{K}$, if

$$x, y \in \mathcal{K}, \quad x + y \in F \quad \Rightarrow \quad x, y \in F,$$

If $\emptyset \neq F \subsetneq \mathcal{K}$, then it is called a **proper face**.

Faces of a Closed Convex Cone, ccc

Face of a ccc \mathcal{K} , $\mathcal{K} + \mathcal{K} \subseteq \mathcal{K}$, $\mathbb{R}\mathcal{K} \subseteq \mathcal{K}$

Let \mathcal{K} be a ccc. A cone $F \subseteq \mathcal{K}$ is a **face** of \mathcal{K} , $F \trianglelefteq \mathcal{K}$, if

$$x, y \in \mathcal{K}, \quad x + y \in F \quad \Rightarrow \quad x, y \in F,$$

If $\emptyset \neq F \subsetneq \mathcal{K}$, then it is called a **proper face**.

Characterization of Faces of PSD Cone \mathbb{S}_+^n

Let $X \in \text{relint}(F)$, $F \trianglelefteq \mathbb{S}_+^n$;

let $X = [U \quad V] \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} [U \quad V]^T$, $D \in \mathbb{S}_{++}^k$

be the spectral decomposition.

two views are: $F = U\mathbb{S}_+^k U^T = \mathbb{S}_+^n \cap (VV^T)^\perp$

Properties of Faces

Some Useful Facts about Faces

- a face of a face is a face;
- an intersection of two faces is a face
- $F_i \trianglelefteq K, F_i = K \cap \phi_i^\perp, i = 1, \dots, k$, implies

$$\cap_i F_i = K \cap \left(\sum_i \phi_i \right)^\perp$$

i.e., intersection exposed faces - exposed by sum of exposing vectors

Properties of Faces

Some Useful Facts about Faces

- a face of a face is a face;
- an intersection of two faces is a face
- $F_i \trianglelefteq K, F_i = K \cap \phi_i^\perp, i = 1, \dots, k$, implies

$$\cap_i F_i = K \cap \left(\sum_i \phi_i \right)^\perp$$

i.e., intersection exposed faces - exposed by sum of exposing vectors

For PSD cone

- Self-replicating: a face of a PSD cone is *still* a PSD cone;
- Facially exposed: every face of PSD cone has exposing vector
- Self-dual: $\mathcal{K} = \mathcal{K}^* = \{x : \langle x, y \rangle \geq 0, \forall y \in \mathcal{K}\}$

Back to the Low-Rank Matrix Completion Problem

Recall (SDP-LRMC) Problem: Given $z \in \mathbb{R}^{\hat{E}}$ a partial matrix, find the matrix Z of **minimum rank** to complete z ,
i.e., $\mathcal{P}_{\hat{E}}(Z) = \mathcal{P}_{\hat{E}}(Q) = z$,

Minimize nuclear norm using SDP

$$\begin{array}{ll} \text{(SDP-LRMC)} & \min \|Y\|_* = \frac{1}{2} \text{trace}(Y) \\ & \text{s.t. } \mathcal{P}_{\bar{E}}(Y) = z \\ & Y \succeq 0, \end{array}$$

where \bar{E} is the set of indices in Y that correspond to \hat{E} , the known entries of the upper right block of $\begin{bmatrix} 0 & Z \\ Z^T & 0 \end{bmatrix} \in \mathbb{S}_+^{m+n}$.

Back to the Low-Rank Matrix Completion Problem

Recall (SDP-LRMC) Problem: Given $z \in \mathbb{R}^{\hat{E}}$ a partial matrix, find the matrix Z of **minimum rank** to complete z ,
i.e., $\mathcal{P}_{\hat{E}}(Z) = \mathcal{P}_{\hat{E}}(Q) = z$,

Minimize nuclear norm using SDP

$$\begin{array}{ll} \text{(SDP-LRMC)} & \min \|Y\|_* = \frac{1}{2} \text{trace}(Y) \\ & \text{s.t. } \mathcal{P}_{\bar{E}}(Y) = z \\ & Y \succeq 0, \end{array}$$

where \bar{E} is the set of indices in Y that correspond to \hat{E} , the known entries of the upper right block of $\begin{bmatrix} 0 & Z \\ Z^T & 0 \end{bmatrix} \in \mathbb{S}_+^{m+n}$.

- Since the **diagonal is free**, note that the **Slater condition (strict feasibility)** does hold for (SDP-LRMC). (And it holds for its dual.)

Facial Reduction of (SDP-LRMC) for Optimal Face

Bipartite Graph, $G_Z = (U_m, V_n, \hat{E})$

With Z and the sampled elements we get a bipartite graph G_Z .

Find Fully Known Submatrix X – a biclique α , $X \cong z[\alpha] \in \mathbb{R}^{p \times q}$

After permutation of rows and columns, WLOG

$$Z = \begin{bmatrix} Z_1 & Z_2 \\ X & Z_3 \end{bmatrix}, \quad z = Z[\hat{E}], \quad \alpha \subseteq \hat{E}, \quad X \cong z[\alpha].$$

Facial Reduction of (SDP-LRMC) for Optimal Face

Bipartite Graph, $G_Z = (U_m, V_n, \hat{E})$

With Z and the sampled elements we get a bipartite graph G_Z .

Find Fully Known Submatrix X – a biclique α , $X \cong z[\alpha] \in \mathbb{R}^{p \times q}$

After permutation of rows and columns, WLOG

$$Z = \begin{bmatrix} Z_1 & Z_2 \\ X & Z_3 \end{bmatrix}, \quad z = Z[\hat{E}], \quad \alpha \subseteq \hat{E}, \quad X \cong z[\alpha].$$

Our algorithm is based on finding **bicliques** in G_Z ; we do this by finding (nontrivial/nondiagonal-block) cliques within symmetric matrix Y .

$$Y = \begin{bmatrix} W_1 & Z \\ Z^T & W_2 \end{bmatrix}$$

Bipartite Graph and Biclique

Partial matrix

$$z \cong \begin{bmatrix} -5 & \text{NA} & 10 & -20 & \text{NA} & -6 \\ 4 & 0 & 4 & 4 & 6 & 6 \\ -3 & \text{NA} & \text{NA} & 32 & 27 & \text{NA} \\ 5 & \text{NA} & 0 & 10 & 12 & \text{NA} \\ \text{NA} & -30 & \text{NA} & \text{NA} & 27 & \text{NA} \\ 3 & -5 & -2 & 8 & \text{NA} & 4 \\ 5 & 5 & \text{NA} & 0 & 3 & \text{NA} \end{bmatrix}, \quad \hat{E} = \{11, 13, 14, 16, 21, \dots, 74, 75\}$$

biclique indices: $\bar{U}_m = \{6, 1, 2\}$, $\bar{V}_n = \{1, 4, 3, 6\}$, $\alpha = \{61, 64, 63, 66, 11, \dots, 26\}$

$$z[\alpha] \equiv X = \begin{bmatrix} 3 & 8 & -2 & 4 \\ -5 & -20 & 10 & -6 \\ 4 & 4 & 4 & 6 \end{bmatrix}.$$

$$Y[\alpha] = \begin{bmatrix} & & & & 3 & 8 & -2 & 4 \\ & & & & -5 & -20 & 10 & -6 \\ & & & & 4 & 4 & 4 & 6 \\ & \text{FREE} & & & & & & \\ 3 & -5 & 4 & & & & & \\ 8 & -20 & 4 & & & & & \\ -2 & 10 & 4 & & & \text{FREE} & & \\ 4 & -6 & 6 & & & & & \end{bmatrix}$$

Our View of Facial Reduction and Exposed Faces

Theorem (Drusvyatskiy, Pataki, W. '15)

Linear transformation $\mathcal{M}: \mathbb{S}^n \rightarrow \mathbb{R}^m$, adjoint \mathcal{M}^* ; feasible set $\mathcal{F} := \{X \in \mathbb{S}_+^n : \mathcal{M}(X) = b\} \neq \emptyset$, $b \in \mathbb{R}^m$. Then a vector v exposes a proper face of $\mathcal{M}(\mathbb{S}_+^n)$ containing $b \iff v$ satisfies the auxiliary system

$$0 \neq \mathcal{M}^*v \in \mathbb{S}_+^n \quad \text{and} \quad \langle v, b \rangle = 0.$$

Let N denote smallest face of $\mathcal{M}(\mathbb{S}_+^n)$ containing b . Then:

- 1 $\mathbb{S}_+^n \cap \mathcal{M}^{-1}N = \text{face}(\mathcal{F})$, the smallest face containing \mathcal{F} .
- 2 For any vector $v \in \mathbb{R}^m$ the following equivalence holds:

$$v \text{ exposes } N \iff \mathcal{M}^*v \text{ exposes } \text{face}(\mathcal{F})$$

Noisy sensor network localization: robust facial reduction and the Pareto frontier

D. Drusvyatskiy, N. Krislock, Y-L. Cheung Voronin, and H. W. '16

Facial Reduction for (SDP-LRMC), r is target rank for Z

Biclique $\alpha \cong$ of G_Z , $z[\alpha] \equiv X \in \mathbb{R}^{p \times q}$

target rank $r \leq \min\{p, q\} < \max\{p, q\}$;

WLOG

$$Z = \begin{bmatrix} Z_1 & Z_2 \\ X & Z_3 \end{bmatrix},$$

$$\text{SVD: } X = [U_1 \quad U_X] \begin{bmatrix} \Sigma \in \mathbb{S}_{++}^r & 0 \\ 0 & 0 \end{bmatrix} [V_1 \quad V_X]^T$$

We get full rank factorization

$$X = \bar{P}\bar{Q}^T = U_1\Sigma V_1^T, \quad \bar{P} = U_1\Sigma^{1/2}, \quad \bar{Q} = V_1\Sigma^{1/2}.$$

Since *rank* is lower semi-continuous: $\text{rank } X = \text{rank } Z$ generically.

In fact our tests form: $Z = PQ^T$

with P, Q random, i.i.d. and full column rank r .

FR using Optimal Y

Rewrite Optimal Y

Assuming we have obtained the desired target rank $Y = r$

$$0 \preceq Y = \begin{bmatrix} U \\ P \\ Q \\ V \end{bmatrix} D \begin{bmatrix} U \\ P \\ Q \\ V \end{bmatrix}^T = \begin{bmatrix} UDU^T & UDP^T & UDQ^T & UDV^T \\ PDU^T & PDP^T & PDQ^T & PDV^T \\ QDU^T & QDP^T & QDQ^T & QDV^T \\ VDU^T & VDP^T & VDQ^T & VDV^T \end{bmatrix}$$

And assume rank $X = r$

$$X = PDQ^T = \bar{P}\bar{Q}^T.$$

implies the ranges satisfy

$$U_1^T U_X = P^T U_X = 0, V_1^T V_X = Q^T V_X = 0$$

$$\begin{aligned} \text{range}(X) &= \text{range}(P) = \text{range}(\bar{P}) = \text{range}(U_1), \\ \text{range}(X^T) &= \text{range}(Q) = \text{range}(\bar{Q}) = \text{range}(V_1). \end{aligned}$$

Constructing Exposing Vectors

Key for facial reduction

We can use an **exposing vector** formed as $U_X U_X^T$ for block PDP^T as well as $V_X V_X^T$ for block QDQ^T and **add appropriate blocks of zeros**:

$$W_X = \left[\begin{array}{c|cc|c} 0 & 0 & 0 & 0 \\ \hline 0 & U_X U_X^T & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right] + \left[\begin{array}{c|cc|c} 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & V_X V_X^T & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right]$$
$$= \left[\begin{array}{c|cc|c} 0 & 0 & 0 & 0 \\ \hline 0 & U_X U_X^T & 0 & 0 \\ \hline 0 & 0 & V_X V_X^T & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right].$$

All three matrices provide exposing vectors.

Facial reduction from exposing vector

$$F^* \triangleq TS_+^{((n+m)-(p+q-2r))} T^T, \quad \text{range } T = \text{null } W_X.$$

Measuring Noise of Biclique $\alpha \in \Theta$

Biclique: $\alpha \subseteq \hat{E}$, $z[\alpha] \cong X \in \mathbb{R}^{p \times q}$, target rank r

singular values of X : $\sigma_1 \geq \dots \geq \sigma_{\min\{p,q\}}$

$$\text{biclique noise: } u_X^P := \frac{\sum_{i=r+1}^{\min\{p,q\}} \sigma_i^2}{0.5p(p-1)} \quad u_X^Q := \frac{\sum_{i=r+1}^{\min\{p,q\}} \sigma_i^2}{0.5q(q-1)}$$

Assign biclique weight

Total noise of all bicliques: $S = \sum_{X \in \Theta} (u_X^P + u_X^Q)$

$$\text{for each } \alpha \in \Theta : \quad w_X^P = 1 - \frac{u_X^P}{S}, \quad w_X^Q = 1 - \frac{u_X^Q}{S}$$

Facial Reduction Process

Facial Reduction Process

- Find set of bicliques Θ , of appropriate sizes

Facial Reduction Process

- Find **set of bicliques** Θ , of appropriate sizes
- Find **corresponding exposing vectors** $\{Y_{\alpha}^{expo}\}_{\alpha \in \Theta}$
calculate their weights $\{\omega_{\alpha}\}_{\alpha \in \Theta}$

Facial Reduction Process

- Find **set of bicliques** Θ , of appropriate sizes
- Find **corresponding exposing vectors** $\{Y_\alpha^{expo}\}_{\alpha \in \Theta}$
calculate their weights $\{\omega_\alpha\}_{\alpha \in \Theta}$
- Calculate the weighted sum of all the exposing vectors

$$Y_{Final}^{expo} = \sum_{\alpha \in \Theta} \omega_\alpha Y_\alpha^{expo}$$

Facial Reduction Process

- Find **set of bicliques** Θ , of appropriate sizes
- Find **corresponding exposing vectors** $\{Y_\alpha^{expo}\}_{\alpha \in \Theta}$
calculate their weights $\{\omega_\alpha\}_{\alpha \in \Theta}$
- Calculate the weighted sum of all the exposing vectors

$$Y_{Final}^{expo} = \sum_{\alpha \in \Theta} \omega_\alpha Y_\alpha^{expo}$$

- Find full column rank V such that $\text{range } V = \text{null } Y_{Final}^{expo}$.

Facial Reduction Process

- Find **set of bicliques** Θ , of appropriate sizes
- Find **corresponding exposing vectors** $\{Y_\alpha^{expo}\}_{\alpha \in \Theta}$
calculate their weights $\{\omega_\alpha\}_{\alpha \in \Theta}$
- Calculate the weighted sum of all the exposing vectors

$$Y_{Final}^{expo} = \sum_{\alpha \in \Theta} \omega_\alpha Y_\alpha^{expo}$$

- Find full column rank V such that $\text{range } V = \text{null } Y_{Final}^{expo}$.
- Solve equivalent smaller problem based on **smaller dimensional matrix** R , where

$$Y = VRV^T$$

Facial Reduction Process

- Find **set of bicliques** Θ , of appropriate sizes
- Find **corresponding exposing vectors** $\{Y_\alpha^{expo}\}_{\alpha \in \Theta}$
calculate their weights $\{\omega_\alpha\}_{\alpha \in \Theta}$
- Calculate the weighted sum of all the exposing vectors

$$Y_{Final}^{expo} = \sum_{\alpha \in \Theta} \omega_\alpha Y_\alpha^{expo}$$

- Find full column rank V such that $\text{range } V = \text{null } Y_{Final}^{expo}$.
- Solve equivalent smaller problem based on **smaller dimensional matrix** R , where

$$Y = VRV^T$$

- (Follows the framework in Drusvyatskiy/Krislock/Cheung-Voronin/W.)

Exploit block structure

Y_{Final}^{expo} has **block structure** so V has a block structure too:

$$Y_{Final}^{expo} = \begin{bmatrix} \sum_{X \in \mathcal{C}} w_X^P W_X^P & 0 \\ 0 & \sum_{X \in \mathcal{C}} w_X^Q W_X^Q \end{bmatrix}, \quad V = \begin{bmatrix} V_P & 0 \\ 0 & V_Q \end{bmatrix}$$

Exploit block structure

Y_{Final}^{expo} has **block structure** so V has a block structure too:

$$Y_{Final}^{expo} = \begin{bmatrix} \sum_{X \in \mathcal{C}} w_X^P W_X^P & 0 \\ 0 & \sum_{X \in \mathcal{C}} w_X^Q W_X^Q \end{bmatrix}, \quad V = \begin{bmatrix} V_P & 0 \\ 0 & V_Q \end{bmatrix}$$

allows a computational speed up for eigenvalue subproblems.

Noiseless Case

FR dramatically **reduces dimension** of now **overdetermined** problem:

$$\begin{aligned} \min \quad & \text{trace}(R) && (= \text{trace}(VRV^T)) \\ \text{s.t.} \quad & \mathcal{P}_{\bar{E}}(V_P R_{pq} V_Q^T) = z \\ & R = \begin{bmatrix} R_p & R_{pq} \\ R_{pq}^T & R_q \end{bmatrix} \succeq 0. \end{aligned}$$

Noiseless Case

FR dramatically **reduces dimension** of now **overdetermined** problem:

$$\begin{aligned} \min \quad & \text{trace}(R) && (= \text{trace}(VRV^T)) \\ \text{s.t.} \quad & \mathcal{P}_{\tilde{E}}(V_P R_{pq} V_Q^T) = z \\ & R = \begin{bmatrix} R_p & R_{pq} \\ R_{pq}^T & R_q \end{bmatrix} \succeq 0. \end{aligned}$$

remove the redundant constraints

Use a **compact QR** to find well-conditioned full rank matrix representation. A simple **semidefinite constrained least squares** solution may be enough!

$$\min_{R \in \mathbb{S}_+^{rv}} \|\mathcal{P}_{\tilde{E}}(V_P R_{pq} V_Q^T) - \tilde{z}\|.$$

(Here \tilde{E}, \tilde{z} denote the corresponding entries after removing redundant constraints. Often R found explicitly.)

Noisy Case

Cannot simply remove redundant constraints;
use random **sketch matrix** A to reduce the number of constraints;
first solve:

$$\delta_0 = \min_{R \in \mathbb{S}_+^{r_0}} \left\| A \left(\mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - z \right) \right\|.$$

and hopefully obtain the target rank!

Noisy Case

Cannot simply remove redundant constraints;
use random **sketch matrix** A to reduce the number of constraints;
first solve:

$$\delta_0 = \min_{R \in \mathbb{S}_+^{r_0}} \left\| A \left(\mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - z \right) \right\|.$$

and hopefully obtain the target rank!
Otherwise, we use a **refinement step**.

Refinement Step in the Noisy Case

We would like to reduce the rank after the previous step using a parametric approach:

$$\begin{array}{ll} \min & \text{trace}(R) \\ \text{s.t.} & \|A(\mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - b)\| \leq \delta_0 \\ & R \succeq 0. \end{array}$$

Refinement Step in the Noisy Case

We would like to reduce the rank after the previous step using a parametric approach:

$$\begin{aligned} \min \quad & \text{trace}(R) \\ \text{s.t.} \quad & \|A(\mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - b)\| \leq \delta_0 \\ & R \succeq 0. \end{aligned}$$

To ensure the rank can be reduced, we flip the problem:

$$\begin{aligned} \varphi(\tau) := \min \quad & \|A(\mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - b)\| + \gamma \|R\|_F \\ \text{s.t.} \quad & \text{trace}(R) \leq \tau \\ & R \succeq 0. \end{aligned}$$

where γ is a regularization parameter, since the least squares problem can be underdetermined.

Table: noiseless: $r = 8$; $m \times n$ size; density p ; mean 20 instances.

Specifications			r_v	Rcvrd (%Z)	Time (s)	Rank	Residual (%Z)
m	n	mean(p)					
1000	3000	0.53	16.10	96.39	37.29	8.0	1.1072e-10
1000	3000	0.50	17.65	88.99	36.50	8.0	4.6569e-10
1000	3000	0.48	32.15	71.66	72.14	8.5	2.0413e-07

Table: noisy: $r = 2$; $m \times n$ size; density p ; mean 20 instances.

Specifications				Rcvd (%Z)	Time (s)		Rank		Residual (%Z)	
m	n	% noise	p		initial	refine	initial	refine	initial	refine
1100	3000	0.50	0.33	100.00	33.72	48.53	2.00	2.00	8.53e-03	8.53e-03
1100	3000	1.00	0.33	100.00	33.67	49.09	2.00	2.00	2.70e-02	2.70e-02
1100	3000	2.00	0.33	100.00	34.13	48.84	2.00	2.00	9.75e-02	9.75e-02
1100	3000	3.00	0.33	100.00	36.34	92.73	5.00	5.00	5.48e-01	1.40e-01
1100	3000	4.00	0.33	100.00	51.45	186.28	11.00	8.00	1.25e+00	1.28e-01

Conclusion

Preprocessing

- Though strict feasibility holds **generically**, failure appears in many applications. Preprocessing based on structure can both *regularize* and simplify the problem.
(New Survey FR: Drusvyatskiy and W. '17)

Exploit structure at optimum

For low-rank matrix completion the structure at the optimum can be exploited to apply FR on the optimal face even though strict feasibility holds. In many cases one gets an optimal solution without the need of any SDP solver.

To do: reduce density/more refinement; real life applications

Thanks for your attention! Questions?

Low-Rank Matrix Completion with Facial Reduction

Shimeng Huang¹ Henry Wolkowicz²

¹Department of Statistics and Actuarial Science
University of Waterloo

²Department of Combinatorics and Optimization
University of Waterloo

AT ECM2017, Thurs. June 1, 10:50AM

References



E.J. Candès and B. Recht, *Exact matrix completion via convex optimization*, *Found. Comput. Math.* 9 (2009), no. 6, 717–772. MR 2565240



D. Drusvyatskiy, N. Krislock, Y-L. Cheung Voronin, and H. Wolkowicz, *Noisy sensor network localization: robust facial reduction and the Pareto frontier*, *SIAM Journal on Optimization* (2017), 1–30, arXiv:1410.6852, 20 pages, to appear.



D. Drusvyatskiy, G. Pataki, and H. Wolkowicz, *Coordinate shadows of semidefinite and Euclidean distance matrices*, *SIAM J. Optim.* 25 (2015), no. 2, 1160–1178. MR 3357643



D. Drusvyatskiy, S. Sremac, and H. Wolkowicz, *Three views of facial reduction in cone optimization*, Tech. report, University of Waterloo, Waterloo, Ontario, 2017, survey in progress.



M. Fazel, *Matrix rank minimization with applications*, Ph.D. thesis, Stanford University, Stanford, CA, 2001.



M. Fazel, H. Hindi, and S.P. Boyd, *A rank minimization heuristic with application to minimum order system approximation*, *Proceedings American Control Conference*, 2001, pp. 4734–4739.



B. Recht, M. Fazel, and P. Parrilo, *Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization*, *SIAM Rev.* 52 (2010), no. 3, 471–501. MR 2680543