

THE MATHEMATICAL LEGACY OF WILLIAM ARVESON

KENNETH R. DAVIDSON

Dedicated to the memory of Bill Arveson

Communicated by the editors

ABSTRACT. This is a retrospective of some of William Arveson's many contributions to operator theory and operator algebras.

KEYWORDS: *dilation theory, completely positive maps, invariant subspaces, maximal subdiagonal operator algebras, dynamical systems, commutative subspace lattices, nest algebras, automorphism groups, multivariable operator theory.*

MSC (2000): 47A, 47D, 47L, 46L

1. INTRODUCTION

William Barnes Arveson, born in November 1934, completed his doctorate in 1964 at UCLA under the supervision of Henry Dye. After an instructorship at Harvard, Bill began a long career at the University of California, Berkeley. He died in November 2011, shortly before his 77th birthday, of complications resulting from surgery.

Arveson's work was deep and insightful, and occasionally completely revolutionary. When he attacked a problem, he always set it in a general framework, and built all of the infrastructure needed to understand the workings. This is one of the reasons that his influence has been so pervasive in many areas of operator theory and operator algebras.

He worked in both operator theory and operator algebras. I think it is fair to say that he did not really distinguish between the two areas, and proved over and over again that these areas are inextricably linked. In [9], he writes in the introduction, and in fact demonstrated many times, that:

"In the study of families of operators on Hilbert space, the self-adjoint algebras have occupied a preeminent position. Nevertheless, many problems in operator theory lead obstinately toward questions about algebras that are not necessarily self-adjoint."

Bill's work moves across the landscape of operator theory and operator algebras, and he developed many foundational ideas which wove the subject together as an integrated whole.

It isn't possible in a short article to review all of Bill's contributions. I will cover many of the highlights. The choices made reflect my personal taste, and I extend my apologies to those who feel that their favourite Arveson paper is omitted. There will be one glaring omission— E_0 -semigroups and CP semigroups. Beginning with [21], Arveson wrote 24 papers (more than 650 pages) and a book (450 pages) on this topic. It is too large a topic to cover in this article, and Masaki Izumi has written a survey [69] of this area to appear in this same volume.

Bill Arveson was well-known to be an outstanding expositor. His papers were always meticulously written. In addition, he wrote three graduate level texts. His *An invitation to C^* -algebras* [17] is distinguished from other C^* -algebra texts in that it develops in detail the structure of type I C^* -algebras and their representation theory. *A short course on spectral theory* [24] is a brief introduction to Banach algebras and operator theory. *Noncommutative dynamics and E -semigroups* [26] is a full development of the theory of E_0 -semigroups.

Bill was also a dedicated supervisor. According to the Mathematics Genealogy Project, he had 27 doctoral students. At the memorial ceremony in Berkeley in February, many of his former students paid their respects and told stories of Bill as a mentor, inspiration and friend. He will be greatly missed. But his mathematics will live on.

2. DILATION THEORY

I will begin with Arveson's contributions to dilation theory. While this is not his first work, it is perhaps his most influential. It has had ramifications in many aspects of operator algebras and operator theory.

In 1953, Bela Sz.Nagy [102] showed that for every operator T on Hilbert space of norm at most one, there is a *dilation* to a unitary operator U on a larger Hilbert space $\mathcal{K} = \mathcal{K}_- \oplus \mathcal{H} \oplus \mathcal{K}_+$ of the form

$$U = \begin{bmatrix} * & 0 & 0 \\ * & T & 0 \\ * & * & * \end{bmatrix},$$

where $*$ represents an unspecified operator. In particular, the map from polynomials in U to polynomials in T obtained by compression to \mathcal{H} is easily seen to be a contractive homomorphism. This recovers the von Neumann inequality for contractions:

$$\|p(T)\| \leq \|p\|_\infty := \sup_{|z| \leq 1} |p(z)| \quad \text{for all } p \in \mathbb{C}[z].$$

When T does not have a unitary summand, this map extends to yield a map from the algebra H^∞ of bounded analytic functions on the unit disk into the weakly closed algebra that T generates. This allows the transference of powerful techniques from function theory to operator theory. See [103, 104] for more in this direction.

These methods extended, in part, to pairs of commuting contractions. In 1964, Ando [5] showed that any commuting pair of contractions simultaneously dilate to a pair of commuting unitaries. In particular, they also satisfy von Neumann's inequality for polynomials in two variables, where the norm on polynomials is the supremum over the bidisk \mathbb{D}^2 . In the one variable case, the minimal dilation is unique in a natural sense. However, difficulties arise in the two variable case because this dilation is no longer unique.

For commuting triples, there is a serious road block. Briefly skipping ahead in the history, Parrott [80] used Arveson's ideas to construct three commuting contractions which do not dilate to commuting unitaries. Furthermore, examples developed later from the work of Varopoulos [108] on tensor product norms showed that even von Neumann's inequality can fail.

In two papers in Acta Math. [11, 13], Arveson revolutionized dilation theory by reformulating it in the framework of an arbitrary operator algebra. The algebra in Sz.Nagy's case is the disk algebra $A(\mathbb{D})$, and for Ando is the bidisk algebra $A(\mathbb{D}^2)$.

There were a number of key ideas that are central to Arveson's approach. First, an operator algebra \mathcal{A} was considered as a subalgebra of a C^* -algebra \mathfrak{B} . Second, the correct representations were *completely contractive*, and yielded *completely positive* maps on $\overline{\mathcal{A} + \mathcal{A}^*}$. Third, he claimed that every operator algebra \mathcal{A} lived inside a unique canonical smallest C^* -algebra called the *C^* -envelope*. These ideas completely changed the way we look at dilations.

To elaborate, the $k \times k$ matrices over the C^* -algebra \mathfrak{B} , $\mathcal{M}_k(\mathfrak{B})$, is itself a C^* -algebra and thus carries a unique norm. This induces a family of norms on the matrix algebras over \mathcal{A} . Today we say that this gives \mathcal{A} an operator space structure. A map φ from \mathcal{A} into $\mathcal{B}(\mathcal{H})$ (or into any C^* -algebra) induces maps φ_k from $\mathcal{M}_k(\mathcal{A})$ into $\mathcal{M}_k(\mathcal{B}(\mathcal{H})) \simeq \mathcal{B}(\mathcal{H}^{(k)})$ by acting by φ on each entry:

$$\varphi_k([a_{ij}]) = [\varphi(a_{ij})].$$

We say that φ is *completely bounded* when

$$\|\varphi\|_{cb} := \sup_{k \geq 1} \|\varphi_k\| < \infty.$$

It is *completely contractive* when $\|\varphi\|_{cb} \leq 1$. When \mathcal{A} is an *operator system* (a unital self-adjoint subspace), we say φ is *completely positive* when φ_k is positive for all $k \geq 1$.

Completely positive maps on C^* -algebras were introduced by Stinespring [100], where he proved a fundamental structure theorem for completely positive

maps. The notion was used by Størmer [101] in a study of positive maps on C^* -algebras. However, Arveson really recognized the power of this idea, and turned completely positive maps into a major tool in operator theory.

Every completely contractive unital map ρ of a unital operator algebra \mathcal{A} extends uniquely to a completely positive map $\tilde{\rho}$ on the operator system $\overline{\mathcal{A} + \mathcal{A}^*}$. Arveson's Extension Theorem established that this map extends to a completely positive map on the C^* -algebra \mathfrak{B} .

Theorem 2.1 (Arveson's extension theorem 1969). *Let \mathcal{S} be an operator system in a unital C^* -algebra \mathfrak{B} . If φ is a completely positive map from \mathcal{S} into $\mathcal{B}(\mathcal{H})$, then there is a completely positive extension Φ of \mathfrak{B} into $\mathcal{B}(\mathcal{H})$.*

In modern terminology, this says that $\mathcal{B}(\mathcal{H})$ is *injective* in the category of operator systems with completely positive maps. This is a generalization of Krein's theorem that positive linear functionals have a positive extension. The norm is not increased because $\|\Phi\| = \|\Phi(1)\| = \|\varphi(1)\| = \|\varphi\|$.

Stinespring's Theorem shows that Φ is a corner of a $*$ -representation π of \mathfrak{B} . When restricted to \mathcal{A} , we obtain a completely contractive representation of the form

$$\pi(a) = \begin{bmatrix} * & 0 & 0 \\ * & \rho(a) & 0 \\ * & * & * \end{bmatrix} \quad \text{for all } a \in \mathcal{A}.$$

This is called a *$*$ -dilation* of ρ . So we have

Theorem 2.2 (Arveson's dilation theorem 1969). *Let \mathcal{A} be a unital operator algebra, and let $\rho : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ be a representation. The following are equivalent:*

- (i) ρ is completely contractive.
- (ii) $\tilde{\rho}$ is completely positive.
- (iii) ρ has a $*$ -dilation.

This result was highlighted in Paulsen's book [83] and the expanded revision [84]. In Arveson's paper, this fundamental result is a remark following Theorem 1.2.9.

Arveson was focussed on the notion of a *boundary representation*, that is, an irreducible $*$ -representation π of \mathfrak{B} with the property that $\pi|_{\mathcal{A}}$ has a unique completely positive extension to \mathfrak{B} . These representations are the non-commutative analogue of points in the spectrum of a function algebra which have a unique representing measure—points in the Choquet boundary. When there are sufficiently many boundary representations to represent \mathcal{A} completely isometrically, then there is a quotient of \mathfrak{B} which is the unique minimal C^* -algebra containing \mathcal{A} completely isometrically. This is the *C^* -envelope* of \mathcal{A} .

Arveson was only able to establish the existence of this C^* -envelope in a variety of special cases. For example, if \mathfrak{B} contains the compact operators and the quotient by the compacts is not completely isometric on $\mathcal{A} + \mathcal{A}^*$, then \mathfrak{B} itself is the C^* -envelope. He analyzed a variety of singly generated examples, always

from the boundary representation viewpoint. In particular, Arveson introduced the notion of *matrix range* of an operator T :

$$W_n(T) = \{ \varphi(T) : \varphi : C^*(T) \rightarrow \mathcal{M}_n \text{ unital completely positive} \} \quad \text{for } n \geq 1.$$

In [12], he proved that the matrix range of an irreducible compact operator formed a complete unitary invariant. Finding unitary invariants for operators was an important open problem. This provides a new invariant even for matrices. This was strengthened in [13] to show that for any irreducible operator T such that $C^*(T)$ contains the compact operators, the matrix range is again a complete unitary invariant.

Theorem 2.3 (Arveson 1972). *If T_1 and T_2 are irreducible operators such $C^*(T_i)$ contains the compact operators, then T_1 and T_2 are unitarily equivalent if and only if*

$$W_n(T_1) = W_n(T_2) \quad \text{for all } n \geq 1.$$

A decade later, Hamana [66] established the existence of a unique minimal injective operator system containing \mathcal{A} , the *injective envelope*, and from this deduced the existence of the C^* -envelope in complete generality. However it shed little light on the existence of boundary representations. In 2005, Dritschel and McCullough [55] produced a very different proof of the existence of the C^* -envelope based on ideas of Agler [1]. It was more direct, and it showed that every representation had a *maximal dilation*, which is a representation such that any further dilation (to a completely contractive representation) can be obtained only by adjoining another representation by a direct sum. This property in another guise was recognized by Muhly and Solel [78], but their result relied on Hamana's theorem. This approach did not deal with the question of irreducible representations either, so it still did not establish the existence of boundary representations. Arveson revisited this question in the light of this new proof, and established the existence of sufficiently many boundary representations when \mathcal{A} is separable [28].

In the past four decades, many tools have been developed for computing C^* -envelopes, and they can now be computed for many classes of examples. They have become a basic tool for the study of nonself-adjoint operator algebras.

Completely bounded maps took on a life of their own in the 1980s. Wittstock [110] proved the extension theorem for completely bounded maps. Paulsen [82] showed that this follows directly from Arveson's extension theorem using a 2×2 matrix trick.

The notion of completely bounded and completely positive maps has had a profound influence on self-adjoint operator algebras. While the goal of Bill's paper was dilation theory, the theory of completely positive maps has had a life of its own in the self-adjoint theory. The injectivity of $\mathcal{B}(\mathcal{H})$ led to deep work on injective von Neumann algebras by Connes [46]. This was followed by work on

nuclear C^* -algebras by Lance [73], and work by Choi, Effros and Lance determining the tight connection between nuclearity of a C^* -algebra and the injectivity of the von Neumann algebra it generates [57, 40, 42].

When Brown, Douglas and Fillmore [35, 36] did their groundbreaking work on essentially normal operators and the Ext functor (a K-homology theory for C^* -algebras), Arveson pointed out how one obtains inverses in the Ext group using completely positive maps. Later he wrote an important paper [18] introducing quasicontral approximate units for C^* -algebras, and used this to provide a unified and transparent approach to Voiculescu's celebrated generalized Weyl-von Neumann theorem [109], the Choi-Effros lifting theorem [41], and the structure of the Ext groups for nuclear C^* -algebras.

Completely positive and completely bounded maps also play a central role in Kadison's similarity problem, which asks whether every bounded representation of a C^* -algebra is similar to a $*$ -representation. This was confirmed for nuclear C^* -algebras by Bunce [37] and Christensen [44]. Then Haagerup [64] established the result for cyclic representations of arbitrary C^* -algebras. Along the way, he showed that a representation was similar to a $*$ -representation if and only if it was completely bounded. See Pisier's monograph [88].

The theory of CP and CB maps led to a considerable revolution on the nonself-adjoint side as well. One important application concerns the Sz.Nagy-Halmos problem. An operator T is called *polynomially bounded operator* if there is a constant C so that

$$\|p(T)\| \leq C\|p\|_\infty \quad \text{for all } p \in \mathbb{C}[x].$$

It is *completely polynomially bounded* if this inequality holds for matrices of polynomials. Halmos's refinement of Sz.Nagy's question asks whether every polynomially bounded operator is similar to a contraction. Paulsen [81] showed that a representation of a unital operator algebra is similar to a completely contractive one if and only if it is completely bounded. He used this to show that every completely polynomially bounded operator is similar to a contraction [82]. Then Pisier [87] showed that there are polynomially bounded operators which are not similar to a contraction, solving the Sz.Nagy-Halmos problem. See [84, 88] for good coverage of this material.

The notions of matrix norms and completely bounded maps led to an abstract categorical approach to abstract operator systems, operator algebras and operator spaces. This is significant because it frees the theory from reliance on spatial representations. Choi and Effros [43] characterized abstract operator systems. Ruan [96] established a GNS style representation theorem for operator spaces; and Blecher, Ruan and Sinclair [34] did the same for unital operator algebras. See [58, 89, 33] for a treatment of this material.

3. MAXIMAL SUBDIAGONAL ALGEBRAS

In his early work, Arveson used nonself-adjoint operator algebras to find non-commutative analogues of results in analytic functions theory, to study dynamics on topological spaces, to study invariant subspaces, and even to study the corona problem of Carleson. We will survey these results in the next few sections.

K. Hoffman [67] defined a *logmodular* function algebra to be a function algebra \mathcal{A} on a compact Hausdorff space X such that the set $\{\log |h| : h \in \mathcal{A}^{-1}\}$ is norm dense in $C_{\mathbb{R}}(X)$. He showed that this class shared many properties that the algebra H^{∞} of bounded analytic functions on the disk enjoys. In fact H^{∞} satisfies the stronger property that every invertible real function f in L^{∞} has the form $\log |h|$ for an invertible H^{∞} function. Indeed one can take $h = \exp(f + i\tilde{f})$ where \tilde{f} is the harmonic conjugate of f . We say that H^{∞} is strongly logmodular, or that it satisfies factorization.

Srinivasan and Wang [99] defined a weak-* closed subalgebra \mathcal{A} of $L^{\infty}(m)$ to be a weak-* Dirichlet algebra if $\mathcal{A} + \mathcal{A}^*$ is weak-* dense in $L^{\infty}(m)$ and the functional $\varphi(f) = \int f dm$ is multiplicative. Hoffman and Rossi [68] show that these algebras are logmodular. A variety of other properties such as factorization, variants of Szegő's theorem and Beurling's invariant subspace theorem are eventually shown to be equivalent to this property. See [32] for a survey of this and the non-commutative generalizations.

In the paper [9] that came out of Arveson's thesis [7], he developed a non-commutative setting for logmodularity. This setting is a finite von Neumann algebra M with a faithful normal state τ . A weak-* closed subalgebra \mathcal{A} is tracial if there is a normal expectation Φ of M onto $\mathcal{A} \cap \mathcal{A}^*$ which is a homomorphism on \mathcal{A} : $\Phi(ab) = \Phi(a)\Phi(b)$ for all $a, b \in \mathcal{A}$, and preserves the trace: $\tau\Phi = \tau$. A (finite) *maximal subdiagonal algebra* is a tracial subalgebra of M such that $\mathcal{A} + \mathcal{A}^*$ is weak-* dense in M .

It isn't hard to see that logmodularity of a function algebra is equivalent to saying that $\{|h|^2 : h \in \mathcal{A}^{-1}\}$ is dense in $C_{\mathbb{R}}(X)$, and that factorization is the stronger property that every invertible real continuous function on X is exactly $|h|^2$ for some invertible $h \in \mathcal{A}$. A subalgebra \mathcal{A} of a C^* -algebra B is called logmodular if $\{a^*a : a \in \mathcal{A}^{-1}\}$ is dense in B_{sa} , and has factorization if this set coincides with the set of all invertible self-adjoint elements. By the polar decomposition, this latter property is equivalent to saying that every invertible element of B factors as $b = ua$ for some $a \in \mathcal{A}$ and unitary $u \in B$.

Arveson [9] established the analogue of the Hoffman-Rossi result:

Theorem 3.1 (Arveson 1967). *Every maximal subdiagonal algebra satisfies factorization.*

An interesting class of examples is obtained as follows. Start with a chain \mathcal{L} of projections (in the usual order) in a finite von Neumann algebra M such that

there is a normal expectation onto the commutant \mathcal{L}' . Then

$$\text{Alg } \mathcal{L} = \{a \in M : ae = eae \text{ for all } e \in \mathcal{L}\}$$

is a maximal subdiagonal algebra. Arveson also provides interesting examples associated to ergodic group actions on measure spaces.

There are a variety of other important properties that H^∞ satisfies, and surprisingly many of them have natural non-commutative analogues. To describe them, we need to introduce the Fuglede-Kadison determinant [60] which they define on II_1 factors, and Arveson extends to all finite von Neumann algebras. Define $\Delta(a) = e^{\tau(\log |a|)}$ when a is invertible, and in general

$$\Delta(a) = \inf_{\varepsilon > 0} e^{\tau(\log(|a| + \varepsilon 1))}$$

This map is multiplicative, positive homogeneous, monotone, self-adjoint; and it is norm continuous on the invertible elements. Arveson makes this much more tractable by showing that

$$\Delta(a) = \inf \tau(|a|b) : b \in M^{-1}, b > 0, \tau(b) \geq 1\}.$$

From this it follows that

$$\Delta(a) \leq \Delta(\Phi(a)) \text{ for all } a \geq 0.$$

Now if $h \in H^\infty$, then Jensen's inequality says that

$$\log |h(0)| \leq \frac{1}{2\pi} \int_0^{2\pi} \log |h(e^{it})| dt.$$

Jensen's formula states that this is an equality when h is invertible. We also mention Szegő's theorem, which states that if f is a positive function in L^1 , then

$$\inf_{h \in H^\infty, h(0)=1} \frac{1}{2\pi} \int_0^{2\pi} |h(e^{it})|^2 f(t) dt = \exp \left(\frac{1}{2\pi} \int_0^{2\pi} \log f(t) dt \right).$$

Arveson defines the analogues in the non-commutative setting: a tracial algebra \mathcal{A} satisfies Jensen's inequality if

$$\Delta(\Phi(a)) \leq \Delta(a) \text{ for all } a \in \mathcal{A}$$

and satisfies Jensen's formula if

$$\Delta(\Phi(a)) = \Delta(a) \text{ for all } a \in \mathcal{A}^{-1}.$$

Szegő's formula becomes:

$$\Delta(b) = \inf \{\tau(ba^*a) : a \in \mathcal{A}, \Delta\Phi(a) \geq 1\}.$$

Arveson [9] established that these three properties are equivalent, and that they are satisfied in many classes of examples. Labuschagne [72] proved that these properties hold for all maximal subdiagonal algebras. Then Blecher and Labuschagne [31] showed that for the class of tracial algebras, these properties are equivalent to logmodularity, and to factorization, and to the weak-* Dirichlet property. See [32] for a survey of this material.

4. DYNAMICAL SYSTEMS

Arveson was interested in using operator algebras to obtain complete invariants for a discrete dynamical system in both the topological and the measure theoretic settings.

In the measure setting, consider an ergodic transformation τ of a measure space (X, m) where m is a probability measure. Two transformations σ and τ are conjugate if there is a measure preserving automorphism Φ of X so that $\sigma = \Phi\tau\Phi^{-1}$. An important problem is to determine when two ergodic transformations are conjugate.

This is easily transferred to a statement about operator algebras. Take $M = L^\infty(m)$ considered as acting on $L^2(m)$ by multiplication. The measure preserving transformation τ induces an automorphism of M by

$$\alpha(M_f) = M_{f \circ \tau} \quad \text{so that} \quad \alpha(\chi_E) = \chi_{\tau^{-1}(E)}.$$

Ergodicity means that the only fixed points of α are the scalars. If σ induces the automorphism β , then conjugacy of σ and τ converts to conjugacy of α and β in the automorphism group of M .

Every automorphism α of M determines a unique unitary map U as follows. Since $L^\infty(m)$ is a dense subset of $L^2(m)$, we set

$$U(g) = \alpha(M_g)1 \quad \text{for all } g \in L^\infty(m).$$

Then one readily checks that this extends to a unitary operator such that

$$\alpha(M_g) = UM_gU^* \quad \text{for all } g \in L^\infty(m).$$

The conjugacy of two transformations reduces to unitary equivalence of the two automorphisms.

Arveson's idea was to construct an operator algebra $\mathcal{A}(\alpha)$ as the norm closure of

$$\mathcal{A}_0(\alpha) = \left\{ \sum_{i=1}^k M_{f_i} U^i : f_i \in L^\infty(m), k \geq 0 \right\}.$$

His main result in [8] is that conjugacy of two automorphisms α and β is equivalent to the unitary equivalence of $\mathcal{A}(\alpha)$ and $\mathcal{A}(\beta)$.

His results actually got a lot more mileage in the topological setting, which he and Josephson developed in [29]. Begin with a locally compact Hausdorff space X and a proper map σ of X into itself. As we shall see, construction of a nonself-adjoint operator algebra does not require the map σ to be invertible. Two dynamical systems (X, σ) and (Y, τ) are (topologically) conjugate if there is a homeomorphism γ of Y onto X so that $\sigma = \gamma\tau\gamma^{-1}$. Again the issue is to decide when two systems are conjugate.

The map σ induces an endomorphism α of $C(X)$ by composition. When σ is a homeomorphism, this map is an automorphism. So we consider a dynamical system as the pair $(C(X), \alpha)$. Arveson and Josephson construct a nonself-adjoint

operator algebra in an analogous way to the measure theoretic setting. Their construction depended on the existence of a quasi-invariant measure and small fixed point set. Indeed they put rather stringent conditions on the maps:

Suppose that (X, σ) and (Y, τ) are two topological dynamical systems satisfying the following conditions

- (i) σ and τ are homeomorphisms.
- (ii) there is a regular Borel probability measure μ on X which is quasi-invariant for σ , i.e. $\mu \circ \sigma$ and μ are mutually absolutely continuous;
- (iii) $\mu(O) > 0$ for every non-empty open set $O \subset X$;
- (iv) the set $P = \{x \in X : \sigma^n(x) = x \text{ for some } n \geq 1\}$ of periodic points has measure zero;
- (v) there is a regular Borel probability measure ν on Y with these same properties, and in addition τ is ergodic with respect to ν .

Arveson and Josephson use these properties to build algebras $\mathcal{A}(\alpha)$ and $\mathcal{A}(\beta)$ as in the measure theoretic setting, but using continuous functions rather than $L^\infty(\mu)$, by constructing an explicit representation on $L^2(\mu)$.

What I will describe instead is a more general construction due to Peters [85] called the *semicrossed product*. This has the major advantages that it does not depend on any measure, and we may consider non-invertible systems (X, τ) where τ is any proper continuous map from X into itself.

The key new notion is that of a *covariant representation* (π, V) where π is a $*$ -representation of $C(X)$ on a Hilbert space \mathcal{H} and $V \in \mathcal{B}(\mathcal{H})$ is a contraction such that

$$V\pi(f) = \pi(\alpha(f))V \quad \text{for all } f \in C(X).$$

Peters defines the operator algebra $C(X) \times_\alpha \mathbb{Z}_+$ to be the universal operator algebra for covariant representations. To form this algebra, let $\mathcal{A}_0(\alpha)$ be the set of all formal polynomials of the form $\sum_{i=0}^k f_i \tau^i$ for $f_i \in C(X)$ and $k \geq 0$. Define a norm by

$$\left\| \sum_{i=0}^k f_i \tau^i \right\| := \sup \left\{ \left\| \sum_{i=0}^k \pi(f_i) V^i \right\| : (\pi, V) \text{ is covariant} \right\}.$$

The *semicrossed product* is the completion of $\mathcal{A}_0(\alpha)$ in this norm.

Peters shows that a sufficient family of covariant representations is given by the orbit representations: fix $x \in X$ and define a representation π_x

$$\pi_x(f) = \text{diag} (f(x), f(\tau(x)), f(\tau^2(x)), \dots)$$

and set V to be the unilateral shift. The direct sum over a dense subset of X yields a faithful completely isometric representation of $\mathcal{A}(\alpha)$. In particular, τ is represented by an isometry.

Peters shows that subject to the Arveson-Josephson hypotheses, their algebra $\mathcal{A}(\alpha)$ coincides with the *semicrossed product* $C(X) \times_\alpha \mathbb{Z}_+$.

I can now state the Arveson-Josephson result in this modern terminology.

Theorem 4.1 (Arveson-Josephson 1969). *Suppose that (X, σ) and (Y, τ) are two topological dynamical systems, and let α and β be the automorphisms of $C(X)$ and $C(Y)$ induced by σ and τ respectively. Furthermore assume conditions (i)–(v) listed above. Then the following are equivalent:*

- (1) (X, σ) and (Y, τ) are conjugate.
- (2) $C(X) \times_{\alpha} \mathbb{Z}_+$ and $C(Y) \times_{\beta} \mathbb{Z}_+$ are completely isometrically isomorphic.
- (3) $C(X) \times_{\alpha} \mathbb{Z}_+$ and $C(Y) \times_{\beta} \mathbb{Z}_+$ are algebraically isomorphic.

Peters improved on this result considerably by removing the conditions (i), (ii), (iii) and (v) at the expense of a slight strengthening of (iv) to

- (a) X is compact.
- (b) σ has no periodic points.

He obtained the same equivalence. Hadwin and Hoover [65] further improved on this. Finally Davidson and Katsoulis [50] removed all of these conditions, and showed that (1), (2) and (3) are equivalent for arbitrary proper maps of locally compact Hausdorff spaces with no conditions on periodic points at all.

5. INVARIANT SUBSPACES

The invariant subspace problem, which remains open today, was a popular topic four decades ago. Arveson made some important contributions, which as usual, stressed operator algebras rather than single operator theory.

If \mathcal{A} is an algebra of operators on a Hilbert space H , let

$$\text{Lat } \mathcal{A} = \{M \text{ is a closed subspace of } \mathcal{H} : \mathcal{A}M \subset M\},$$

and if \mathcal{L} is a collection of subspaces, then

$$\text{Alg}(\mathcal{L}) = \{T \in \mathcal{B}(\mathcal{H}) : TM \subset M \text{ for all } M \in \mathcal{L}\}.$$

Halmos called an operator algebra *reflexive* if $\mathcal{A} = \text{Alg Lat } \mathcal{A}$. This algebra is always unital and closed in the weak operator topology (WOT-closed). An operator algebra is *transitive* if it has no proper closed invariant subspace. The transitive algebra problem asks whether there is a proper unital WOT-closed transitive operator algebra.

Arveson's first contribution to invariant subspace theory recognized the advantage of having a masa (maximal abelian self-adjoint algebra) in \mathcal{A} . Note that he shows that the algebra is dense in the weak-* topology, rather than weak operator topology. This is a subtle, and occasionally non-trivial, strengthening.

Theorem 5.1 (Arveson's density theorem 1967). *A transitive subalgebra of $\mathcal{B}(\mathcal{H})$ which contains a masa is weak-* dense in $\mathcal{B}(\mathcal{H})$.*

This result received a fair bit of interest, and a generalization was found by Radjavi and Rosenthal [93]: if \mathcal{A} is an operator algebra containing a masa, and $\text{Lat } \mathcal{A}$ is a chain (nest), then the WOT-closure of \mathcal{A} is reflexive.

Perhaps this result prompted Bill to revisit the issue. He decided to study weak-* closed operator algebras containing a masa, to determine whether all such operator algebras are reflexive. We assume that the Hilbert space is separable. The masa \mathcal{D} is spatially isomorphic to $L^\infty(\mu)$ acting on $L^2(\mu)$ for some sigma-finite Borel measure μ . If $\mathcal{A} \supset \mathcal{D}$, then $\text{Lat } \mathcal{A} \subset \text{Lat } \mathcal{D}$. If $M \in \text{Lat } \mathcal{D}$, then the orthogonal projection P_M commutes with \mathcal{D} ; and by maximality, P_M belongs to \mathcal{D} . So $\text{Lat } \mathcal{A}$ can be naturally identified with a sublattice of the lattice of projections in \mathcal{D} . In particular, these projections all commute. In addition, the lattice is complete (as a lattice, and as a SOT-closed set of projections). Such a lattice is called a *commutative subspace lattice* (CSL).

In a 100 page tour de force in the Annals of Mathematics, Arveson [14] developed an extensive theory for these operator algebras. The first major result is a spectral theorem for CSLs.

Theorem 5.2 (Arveson 1974). *If \mathcal{L} is a CSL on a separable Hilbert space, there is a compact metric space X , a regular Borel measure μ on X , and a Borel partial order \preceq on X so that \mathcal{L} is unitarily equivalent to the lattice of projections onto the (essentially) increasing Borel sets.*

Then he develops a class of pseudo-integral operators and identifies when one is supported on the graph of the partial order; and hence belongs to $\text{Alg } \mathcal{L}$. Curiously, while the pseudo-integral operators seem to depend on the choice of coordinates, the weak-* closure of the pseudo-integral operators supported on the graph of the partial order is independent of all choices. He calls this algebra $\mathcal{A}_{\min}(\mathcal{L})$. He shows that $\text{Lat } \mathcal{A}_{\min}(\mathcal{L}) = \mathcal{L}$ and hence $\text{Lat } \text{Alg } \mathcal{L} = \mathcal{L}$ for all CSLs. One of the main results is:

Theorem 5.3 (Arveson 1974). *If \mathcal{L} is a CSL on a separable Hilbert space, then there is a weak-* closed operator algebra $\mathcal{A}_{\min}(\mathcal{L})$ containing a masa with $\text{Lat}(\mathcal{A}_{\min}(\mathcal{L})) = \mathcal{L}$ so that: a weak-* closed operator algebra \mathcal{A} containing a masa has $\text{Lat}(\mathcal{A}) = \mathcal{L}$ if and only if $\mathcal{A}_{\min} \subseteq \mathcal{A} \subseteq \text{Alg}(\mathcal{L})$.*

Surprisingly, it is possible for $\mathcal{A}_{\min}(\mathcal{L})$ to be properly contained in $\text{Alg}(\mathcal{L})$, even if it is WOT-closed. An example is constructed based on the failure of spectral synthesis on S^3 in harmonic analysis. He calls a lattice *synthetic* if $\mathcal{A}_{\min}(\mathcal{L}) = \text{Alg}(\mathcal{L})$. Many examples of synthetic lattices are produced, including lattices which are generated by finitely many commuting chains, and lattices associated to certain ordered groups. This is a definitive answer to the question.

The area of CSL algebras has received a lot of attention. See [20] for a survey of some of this material. Also see [49] for further material.

6. NEST ALGEBRAS AND THE CORONA THEOREM

Kadison and Singer [71] initiated a theory of triangular operator algebras. While their algebras were not necessarily WOT-closed or even norm-closed, Ringrose [94, 95] developed a related class that are WOT-closed and reflexive. He calls a complete chain of subspaces \mathcal{N} a *nest*. A *nest algebra* is $\mathcal{T}(\mathcal{N}) := \text{Alg } \mathcal{N}$, the algebra of all operators which are upper triangular with respect to the nest \mathcal{N} . In the first paper, he characterized the Jacobson radical. In the second, he showed that any algebra isomorphism between two nest algebras is given by a similarity.

Arveson's paper [16] was very influential for the subject of nest algebras. In particular, he developed a distance formula from an arbitrary operator to $\mathcal{T}(\mathcal{N})$. Note that if $N \in \mathcal{N}$ and $T \in \mathcal{T}(\mathcal{N})$, the invariance of N may be expressed algebraically by $TP_N = P_N TP_N$, or equivalently $P_N^\perp TP_N = 0$. Therefore if $A \in \mathcal{B}(\mathcal{H})$ and $T \in \mathcal{T}(\mathcal{N})$, we see that

$$\|P_N^\perp AP_N\| = \|P_N^\perp(A - T)P_N\| \leq \inf_{T \in \mathcal{T}(\mathcal{N})} \|T - A\| = \text{dist}(A, \mathcal{T}(\mathcal{N})).$$

This inequality remains valid if we take the supremum over all $N \in \mathcal{N}$. Surprisingly, this leads to an exact formula:

Theorem 6.1 (Arveson's distance formula 1975). *Let \mathcal{N} a nest on a Hilbert space \mathcal{H} ; and let $A \in \mathcal{B}(\mathcal{H})$. Then*

$$\text{dist}(A, \mathcal{T}(\mathcal{N})) = \sup_{N \in \mathcal{N}} \|P_N^\perp AP_N\|.$$

This and other results about nest algebras proved important in the development of the theory.

A separable continuous nest is a nest which is order isomorphic to $I = [0, 1]$. The Volterra nest \mathcal{N} consisting of subspaces

$$N_t = L^2(0, t) \subset \mathcal{H} = L^2(0, 1) \quad \text{for } 0 \leq t \leq 1$$

is an example which is multiplicity free, in that the projections onto N_t generate a masa. The infinite ampliation \mathcal{M} given by $M_t = L^2((0, t) \times I) \subset L^2(I^2)$ generates an abelian von Neumann algebra with non-abelian commutant. So $\mathcal{T}(\mathcal{N})$ and $\mathcal{T}(\mathcal{M})$ are not unitarily equivalent. Ringrose [95] asked whether they could be similar.

Three students of Arveson's eventually solved this problem completely. Andersen [4] showed that any two continuous nests are approximately unitarily equivalent. So in particular, there is a sequence of unitaries U_n so that

$$[0, 1] \ni t \rightarrow P_{M_t} - U_n P_{N_t} U_n^*$$

are continuous, compact operator valued functions converging uniformly to 0. (Arveson [19] generalized this to certain CSL algebras.) Larson [74] used this to solve Ringrose's problem by showing that any two continuous nests are similar. Shortly afterwards, Davidson [48] showed that if two nests are order isomorphic

via a map which preserves dimensions of differences of nested subspaces, then there is a similarity of the two nests implementing the isomorphism; and the similarity may be taken to be a small compact perturbation of a unitary operator. See [49] for more material on nest algebras.

Arveson was also interested in finding an operator theoretic proof of the famous corona theorem of Carleson [39]. This result is formulated as follows: suppose that f_1, \dots, f_n are functions in H^∞ for which there is an $\varepsilon > 0$ so that

$$(\dagger) \quad \inf_{z \in \mathbb{D}} \sum_{i=1}^n |f_i(z)|^2 \geq \varepsilon.$$

Then there are functions $g_i \in H^\infty$ so that $\sum_{i=1}^n f_i g_i = 1$. A reformulation of this result is that the unit disk \mathbb{D} is dense in the maximal ideal space of H^∞ . Carleson's proof was very difficult, and people continue to look for more accessible proofs. An easier (but not easy) proof, based on unpublished ideas of Thomas Wolff, is now available. See Garnett's book [61].

Arveson had an operator theoretic approach to the corona theorem. H^2 is the Hardy space of square integrable analytic functions on the disk \mathbb{D} considered as a subspace of $L^2(\mathbb{T})$. Represent H^∞ on H^2 as multiplication operators T_h (these are analytic Toeplitz operators). In [16], he established this variation of Carleson's theorem:

Theorem 6.2 (Arveson's Toeplitz corona theorem 1975). *If $f_1, \dots, f_n \in H^\infty$ and*

$$(\ddagger) \quad \sum_{i=1}^n T_{f_i} T_{f_i}^* \geq \varepsilon I,$$

then there are $g_i \in H^\infty$ so that $\sum_{i=1}^n f_i g_i = 1$.

This result is weaker than the corona theorem because (\ddagger) is a stronger hypothesis than (\dagger) . To see this, we use the fact that H^2 is a reproducing kernel Hilbert space: the functions

$$k_w(z) = \frac{(1 - |w|^2)^{1/2}}{1 - z\bar{w}}$$

are unit vectors for $w \in \mathbb{D}$ with the property that $T_h^* k_w = \overline{h(w)} k_w$. Applying the vector state $\rho_w(A) = \langle A k_w, k_w \rangle$ to (\ddagger) for $w \in \mathbb{D}$, one recovers Carleson's condition (\dagger) .

It turns out that Toeplitz corona theorems are useful in establishing full corona theorems. Ball, Trent and Vinnikov [30] established a Toeplitz corona theorem for all complete Nevanlinna-Pick kernels as a consequence of their commutant lifting theorem. This was a useful step towards a bona fide corona theorem by Costea, Sawyer and Wick [47] for Drury-Arveson space, among other spaces. For definitions of these concepts, see section 8.

7. AUTOMORPHISM GROUPS

In [15], Arveson developed a spectral analysis of automorphism groups acting on C^* -algebras and von Neumann algebras.

Here G is a locally compact abelian group with Haar measure dt . A representation is a bounded representation $t \rightarrow U_t$ in $\mathcal{B}(X)$, where X is a Banach space, and the representation is assumed to be continuous with respect to a certain weak topology. In particular, SOT-continuous unitary representations on Hilbert space form an important special case. A standard technique extends this to a bounded representation π_U of $L^1(G)$ such that

$$\pi_U(f)\xi = \int_G f(t)U_t\xi dt.$$

We may regard the Fourier transform as the Gelfand transform of the commutative Banach algebra $L^1(G)$ into $C(\hat{G})$, where \hat{G} is the dual group consisting of continuous characters of G . When $f \in L^1(G)$, its zero set is $Z(f) = \{\gamma \in \hat{G} : \hat{f}(\gamma) = 0\}$.

We define the spectrum of the representation U to be

$$\text{sp } U = \bigcap \{Z(f) : f \in L^1(G), \pi_U(f) = 0\}.$$

Moreover, if $\xi \in \mathcal{H}$, then the spectrum of ξ is

$$\text{sp}_U(\xi) = \bigcap \{Z(f) : f \in L^1(G), \pi_U(f)\xi = 0\}.$$

Then if E is a closed subset of \hat{G} , we define the *spectral subspace* to be

$$M^U(E) = \{\xi : \text{sp}_U(\xi) \subset E\}.$$

When U is a strongly continuous unitary representation on Hilbert space, Stone's theorem yields a projection valued measure dP_γ on \hat{G} so that $U_t = \int_{\hat{G}} \gamma(t) dP_\gamma$; and $M^U(E) = P(E)\mathcal{H}$. But in more general contexts, there is often no version of Stone's theorem. Nevertheless, the spectral subspaces form a substitute, and they determine the representation uniquely:

Theorem 7.1 (Arveson 1974). *Let X be a Banach space, and let U and V be two strongly continuous representations of G on X . If $M^U(E) = M^V(E)$ for every compact subset E of \hat{G} , then $U = V$.*

An important application of this theory is to a one parameter group of automorphisms acting on a von Neumann algebra $\mathcal{R} \subset \mathcal{B}(\mathcal{H})$. That is, $t \rightarrow \alpha_t$ is a weak- $*$ continuous representation of \mathbb{R} into $\text{Aut}(\mathcal{R})$. Arveson was interested in finding conditions that determine when this group is unitarily implemented, i.e. when $\alpha_t = \text{Ad } U_t$, where U is a strongly continuous unitary representation with $U_t \in \mathcal{R}$ and $\text{sp } U \subset [0, \infty)$. Let $\mathcal{R}^\alpha(E) = \{A \in \mathcal{R} : \text{sp}_\alpha(A) \subset E\}$. The answer is given by:

Theorem 7.2 (Arveson 1974). *Let (α_t) be a weak-* continuous representation of \mathbb{R} into $\text{Aut}(\mathcal{R})$. Then there is a strongly continuous unitary representation U with $U_t \in \mathcal{R}$ and $\text{sp } U \subset [0, \infty)$ such that $\alpha_t = \text{Ad } U_t$ for $t \in \mathbb{R}$ if and only if*

$$\bigcap_{t \in \mathbb{R}} \mathcal{R}^\alpha[t, \infty) = \{0\}.$$

Moreover $U_t = \int_{-\infty}^{+\infty} e^{itx} dP(x)$ where P is the spectral measure such that

$$P[t, \infty)\mathcal{H} = \bigcap_{s < t} \mathcal{R}^\alpha[s, \infty)\mathcal{H}.$$

Arveson used this to provide a new proof of the famous theorem of Kadison [70] and Sakai [97] that every derivation of a von Neumann algebra is inner. Alain Connes [45] used the Arveson spectrum to define the more refined Connes spectrum of an automorphism group that allows one to define the type III_λ factors.

One can show that

$$\mathcal{R}^\alpha[s, \infty)\mathcal{R}^\alpha[t, \infty) \subset \mathcal{R}^\alpha[s+t, \infty).$$

It follows that $H_\alpha^\infty(\mathcal{R}) = \mathcal{R}^\alpha[0, \infty)$ is a weak-* closed subalgebra of \mathcal{R} . This is sometimes a maximal subdiagonal algebra, but not always. A very curious result of Loeb and Muhly [75] exhibits an example where $H_\alpha^\infty(\mathcal{R})$ is a proper nonself-adjoint subalgebra of \mathcal{R} which is WOT-dense in \mathcal{R} . A variant on the transitive algebra problem deals with *reductive* algebras, meaning that every invariant subspace is reducing (or that the orthogonal complement of every invariant subspace is invariant). It asks whether every WOT-closed reductive operator is a von Neumann algebra. Loeb and Muhly's construct is a reductive *weak-* closed* algebra which is not self-adjoint (but the WOT-closure is a von Neumann algebra).

I will now skip ahead to some more recent results. As I mentioned in the introduction, in the intervening period, Bill was very interested in E_0 -semigroups and CP-semigroups. This work will be covered in the article [69] by Izumi in this volume.

8. MULTIVARIABLE OPERATOR THEORY

As discussed in section 2, the Sz.Nagy dilation theorem led to a development of function-theoretic techniques in single operator theory. In particular, if T is a contraction with no unitary summand, then it has an H^∞ functional calculus. This has many ramifications. See [104] for an updated look at this material.

The examples of Parrott and Varopoulos, also mentioned in section 2, explain why there are problems with a straightforward generalization of this dilation theory to several variables by studying d -tuples of commuting contractions.

It turns out that a different choice of norm condition leads to a better theory. The more tractable norm condition on an d -tuple T_1, \dots, T_d is a row contractive condition: consider $T = [T_1, \dots, T_d]$ as an operator mapping the direct sum $\mathcal{H}^{(d)}$ of d copies of \mathcal{H} into \mathcal{H} and require that

$$\|T\| = \left\| \sum_{i=1}^d T_i T_i^* \right\|^{1/2} \leq 1.$$

In the non-commutative case, the appropriate model for the dilation theorem is best described as the left regular representation of the free semigroup \mathbb{F}_d^+ , consisting of all words in an alphabet $\{1, \dots, d\}$, acting on $\ell^2(\mathbb{F}_d^+)$ by

$$L_i \xi_w = \xi_{iw} \quad \text{for } 1 \leq i \leq d, w \in \mathbb{F}_d^+.$$

The space $\ell^2(\mathbb{F}_d^+)$ can be realized as the full Fock space over $E = \mathbb{C}^d$, namely

$$\ell^2(\mathbb{F}_d^+) \simeq \sum_{k \geq 0}^{\oplus} E^{\otimes k} = \mathbb{C} \oplus \mathbb{C}^d \oplus \mathbb{C}^{d^2} \oplus \dots$$

where $E^{\otimes k} = \mathbb{C}^{d^k}$ is identified with the span of words of length k . In this view, the operators L_i become the *creation operators* that tensor on the left by e_i , where e_1, \dots, e_d is the standard basis for \mathbb{C}^d .

The dilation theorem is that every row contraction T dilates to a row isometry $V = [V_1, \dots, V_d]$ where $V_i^* V_j = \delta_{ij} I$. Furthermore, there is a decomposition $V_i \simeq L_i^{(\alpha)} \oplus W_i$ where α is a cardinal and W_1, \dots, W_d are generators of a representation of the Cuntz algebra \mathcal{O}_d . This was established for $d = 2$ by Frazho [59], for finite d by Bunce [38] and for arbitrary d including infinite values, and uniqueness, by Popescu [90]. Popescu has written a long series of papers developing the Sz.Nagy-Foiaş theory in this context. The WOT-closed algebras generated by row isometries have also been extensively studied by Popescu beginning with [91] and Davidson and Pitts beginning with [51].

The dilation theory for commuting row contractions came earlier, but was exploited later. Drury [56] showed that every commuting row contraction with $\|T\| < 1$ dilates to a direct sum of copies of a certain d -tuple of weighted shifts, $S = [S_1, \dots, S_d]$, now called the *d-shift*. This was refined by Müller and Vasilescu [79] to show that if $\|T\| \leq 1$, then T dilates to a row contraction V with commuting entries so that $V_i \simeq S_i^{(\alpha)} \oplus N_i$ where the N_i are commuting normal operators such that $\sum_{i=1}^d N_i N_i^* = I$, a *spherical isometry*. Perhaps the reason this did not go further at the time is that the exact nature of the d -tuple S was not realized.

Arveson began a program of multivariable operator theory in [22]. He improved the Drury-Müller-Vasilescu dilation theorem. Moreover, he showed that the d -shift has a natural representation on symmetric Fock space over $E = \mathbb{C}^d$. Let E^k denote the symmetric tensor product of k copies of E , which can be considered as the subspace of $E^{\otimes k}$ which is fixed by the action of the permutation group S_k

which permutes the terms of a tensor product of k vectors. Then

$$H_d^2 = \sum_{k \geq 0}^{\oplus} E^k \subset \ell^2(\mathbb{F}_d^+).$$

This space is coinvariant for the creation operators L_i , and $S_i = P_{H_d^2} L_i|_{H_d^2}$.

There is a natural basis

$$z^k = \frac{k!}{|k|!} \sum_{|w|=k, w(z)=z^k} \xi_w,$$

where $k = (k_1, \dots, k_d) \in \mathbb{N}_0^d$, $|k| = \sum_i k_i$, $k! := \prod_i k_i!$ and $z^k := z_1^{k_1} z_2^{k_2} \dots z_d^{k_d}$. This is an orthogonal but not orthonormal basis since $\|z^k\|^2 = \frac{k!}{|k|!} =: c_k$. In this basis, S_i is just multiplication by z_i . The space H_d^2 may now be considered as the space of analytic functions

$$f(z) = \sum_{k \in \mathbb{N}_0^d} a_k z^k \quad \text{such that} \quad \sum_{k \in \mathbb{N}_0^d} |a_k|^2 c_k < \infty.$$

These series converge on the unit ball \mathbb{B}^d of \mathbb{C}^d , so are bona fide functions. This is a reproducing kernel Hilbert space (RKHS), meaning that $w \in \mathbb{B}^d$ separate points and the value of $f(w)$ may be recovered by

$$f(w) = \langle f, k_w \rangle \quad \text{where} \quad k_w(z) = \frac{1}{1 - \langle z, w \rangle}, \quad w \in \mathbb{B}^d.$$

The space H_d^2 is now often called Drury-Arveson space.

Every RKHS has an algebra of multipliers, those functions which multiply the space into itself. The multiplier algebra $\mathcal{M}_d = \text{Mult}(H_d^2)$ of H_d^2 consists of bounded analytic functions on the ball \mathbb{B}_d . It is the WOT-closed unital algebra generated by S_1, \dots, S_d . However the norm is greater than the sup norm, and is not comparable—so \mathcal{M}_d is a proper subalgebra of $H^\infty(\mathbb{B}_d)$.

The C*-algebra $C^*(S) := C^*(S_1, \dots, S_n)$ contains the compact operators \mathfrak{K} , and $C^*(S)/\mathfrak{K} \simeq C(\partial\mathbb{B}^d)$, the boundary sphere of \mathbb{B}^d —which is the spectrum of the generic spherical isometry. So one deduces that this is the C*-envelope of the unital nonself-adjoint algebra \mathcal{A}_d generated by S_1, \dots, S_d .

The classical Nevanlinna-Pick theorem [86] for the unit disk states that given $z_1, \dots, z_n \in \mathbb{D}$ and $w_1, \dots, w_n \in \mathbb{C}$, there is a function $h \in H^\infty$ with $\|h\|_\infty \leq 1$ so that $h(z_i) = w_i$ if and only if

$$\begin{bmatrix} 1 - w_i \bar{w}_j \\ 1 - z_i \bar{z}_j \end{bmatrix}_{n \times n}$$

is positive definite. A matrix version states that given z_1, \dots, z_n in the disk, and $r \times r$ matrices W_1, \dots, W_n , there is a function F in the the unit ball of $\mathcal{M}_r(H^\infty)$

such that $F(z_i) = W_i$ if and only if the Pick matrix

$$\left[\begin{array}{c} I_r - W_i W_j^* \\ 1 - z_i \bar{z}_j \end{array} \right]_{n \times n}$$

is positive semidefinite. Sarason [98] put this into an operator theoretic context.

An RKHS \mathcal{H} of functions on a set X is a natural place to study interpolation. One asks the same question: given $x_1, \dots, x_n \in X$ and $r \times r$ matrices W_1, \dots, W_n , is there a function F in the unit ball of $\mathcal{M}_r(\text{Mult}(\mathcal{H}))$ so that $F(z_i) = W_i$? It is easy to show that a necessary condition is the positive semidefiniteness of the matrix

$$\left[(I_r - W_i W_j^*) \langle k_{x_j}, k_{x_i} \rangle \right]_{n \times n}.$$

A space for which this is also a sufficient condition is called a complete Nevanlinna-Pick kernel.

Complete Nevanlinna-Pick kernels were characterized by Quiggin [92] and McCullough [76]. It follows from their characterization that H_d^2 is a complete Nevanlinna-Pick kernel. This was shown from a completely different angle, by showing that the non-commutative WOT-closed algebra \mathfrak{L}_d on Fock space generated by L_1, \dots, L_d has a good distance formula for ideals [52, 6], and hence has a Nevanlinna-Pick theory based on Sarason's approach. It follows that \mathcal{M}_d is a complete quotient of \mathfrak{L}_d , and hence is a complete Nevanlinna-Pick kernel. Finally, Agler and McCarthy [2] give a new proof of the Quiggin-McCullough theorem, and show that \mathcal{M}_d for $d = \infty$ is the universal complete Nevanlinna-Pick kernel in the sense that every irreducible complete Nevanlinna-Pick RKHS imbeds in a natural way into H_d^2 . See [3] for an overview of this theory.

The confluence of these very different directions leading to the same space, H_d^2 , is remarkable: the good dilation theory for commuting row contractions, the close relationship with the non-commutative model \mathfrak{L}_d , and the fact that this is a complete Nevanlinna-Pick kernel. All play a role in making H_d^2 a very fertile environment in which to study multivariable operator theory.

Associated to any row contraction T is the completely positive contractive map

$$\varphi(A) = \sum_{i=1}^d T_i A T_i^*.$$

Notice that T is spherical if and only if $\varphi(I) = I$. We call a row contraction T *pure* if there is no spherical part. The decreasing sequence

$$I \geq \varphi(I) \geq \varphi^2(I) \geq \dots$$

always has a limit in the strong operator topology. This limit $\varphi_\infty(I)$ is 0 if and only if T is pure.

In [23], Arveson begins a much more algebraic approach to multivariable operator theory. Douglas particularly has espoused the approach of considering

a Hilbert space \mathcal{H} with an operator algebra \mathcal{A} acting on it via a representation $\rho : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ as a Hilbert module over \mathcal{A} with the action $a \cdot \xi = \rho(a)\xi$. This approach was developed for function algebras in the monograph [53] by Douglas and Paulsen.

In this context, one begins with a commuting row contractive d -tuple T acting on \mathcal{H} , and consider \mathcal{H} as a module over $\mathbb{C}[z] := \mathbb{C}[z_1, \dots, z_d]$ given by $p \cdot \xi = p(T_1, \dots, T_d)\xi$. In order to use results from algebra effectively, assume that the defect operator

$$\Delta := (I - TT^*)^{1/2} = \left(I - \sum_{i=1}^d T_i T_i^*\right)^{1/2}$$

has *finite rank*; and define $\text{rank}(T) := \text{rank}(\Delta)$. We also say that \mathcal{H} is a Hilbert module of the same rank.

Form a finitely generated module over $\mathbb{C}[z]$ as follows

$$M_{\mathcal{H}} := \{p \cdot \xi : p \in \mathbb{C}[z], \xi \in \Delta\mathcal{H}\}.$$

This space is *not closed*; so it is a finitely generated $\mathbb{C}[z]$ module algebraically. From commutative algebra (Hilbert's syzygy theorem), one obtains a free resolution

$$0 \rightarrow F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow M_{\mathcal{H}} \rightarrow 0.$$

Each F_i is a finitely generated free $\mathbb{C}[z]$ -module of rank $\beta_i < \infty$. One can define an invariant known as the Euler characteristic

$$\chi(\mathcal{H}) := \sum_{i=1}^n (-1)^{i+1} \beta_i$$

which is independent of the choice of resolution.

Motivated by an analogy with the Gauss-Bonnet theorem relating the curvature of a manifold to its topological invariants, Arveson also defines an analytic invariant that he calls *curvature*. Set

$$T(z) = \sum_{i=1}^d \bar{z}_i T_i = [T_1 \ \dots \ T_d][z_1 \ \dots \ z_d]^*.$$

This satisfies $\|T(z)\| \leq \|z\|_2$. Thus we may define a function on the ball by

$$F(z) = \Delta(I - T(z)^*)^{-1}(I - T(z))^{-1}\Delta|_{\Delta\mathcal{H}} \in \mathcal{B}(\Delta\mathcal{H}) \quad \text{for } z \in \mathbb{B}_d.$$

The boundary values, appropriately weighted, exist in the sense that

$$K_0(\zeta) = \lim_{r \rightarrow 1^-} (1 - r^2) \text{Tr} F(r\zeta) \quad \text{exists a.e. for } \zeta \in \partial\mathbb{B}_d.$$

Define the *curvature* of \mathcal{H} to be

$$K(\mathcal{H}) = \int_{\partial\mathbb{B}_d} K_0(\zeta) d\sigma(\zeta)$$

where σ is normalized Lebesgue measure on the sphere $\partial\mathbb{B}_d$.

Theorem 8.1 (Arveson 2000). *Let \mathcal{H} be a finite rank Hilbert module. Then*

$$K(\mathcal{H}) = d! \lim_{n \rightarrow \infty} \frac{\text{Tr}(I - \varphi^{n+1}(I))}{n^d}$$

and

$$\chi(\mathcal{H}) = d! \lim_{n \rightarrow \infty} \frac{\text{rank}(I - \varphi^{n+1}(I))}{n^d}.$$

It is immediate that $0 \leq K(\mathcal{H}) \leq \chi(\mathcal{H})$. Examples show that this inequality may be proper. However there is an important context in which they are equal. Say that \mathcal{H} is *graded* if there is a strongly continuous unitary representation Γ of the circle \mathbb{T} on \mathcal{H} so that

$$\Gamma(t)T_i\Gamma(t)^* = tT_i \quad \text{for all } 1 \leq i \leq d, t \in \mathbb{T}.$$

This is called a gauge group on \mathcal{H} . When \mathcal{H} is graded, Fourier series allows the decomposition of \mathcal{H} into a direct sum of subspaces $\mathcal{H}_n = \{\xi \in \mathcal{H} : \Gamma(t)\xi = t^n\xi\}$. When \mathcal{H} is finite rank, these are all finite dimensional. Arveson calls the following result his analogue of the Gauss-Bonnet theorem:

Theorem 8.2 (Arveson 2000). *Let \mathcal{H} be a graded finite rank Hilbert module. Then*

$$K(\mathcal{H}) = \chi(\mathcal{H}).$$

In particular, the curvature is always an integer in the graded case. Arveson hypothesized that it was always an integer. This conjecture was verified by Greene, Richter and Sundberg [62].

An important tool for multivariable operator theory is the Taylor spectrum and functional calculus [106, 107]. Arveson [25] builds a Dirac operator based on this theory. Let $Z = \mathbb{C}^d$ with basis e_1, \dots, e_d ; and let ΛZ be the exterior algebra over Z , namely

$$\Lambda Z = \Lambda^0 Z \oplus \Lambda^1 Z \oplus \dots \oplus \Lambda^d Z$$

where $\Lambda^k Z$ is spanned by vectors of the form $x_1 \wedge \dots \wedge x_k$. For $z \in Z$, there is a Clifford operator given by

$$C(z)(x_1 \wedge \dots \wedge x_k) = z \wedge x_1 \wedge \dots \wedge x_k.$$

Now form $\tilde{\mathcal{H}} = \mathcal{H} \otimes \Lambda Z$, and for $\lambda \in \mathbb{C}^d$ and $z \in Z$, define

$$B_\lambda = \sum_{i=1}^d (T_i - \lambda_i I) \otimes C(e_i), \quad D_\lambda = B_\lambda + B_\lambda^* \quad \text{and} \quad R(z) = I \otimes (C(z) + C(z)^*).$$

Arveson observes that λ is not in the Taylor spectrum of T if and only if D_λ is invertible. The pair (D_0, R) is called the Dirac operator for T .

Now the graded space $\tilde{\mathcal{H}}$ splits into its even and odd parts:

$$\tilde{\mathcal{H}}_+ = \sum_{i \text{ even}} \oplus (\mathcal{H} \otimes \Lambda^i Z) \quad \text{and} \quad \tilde{\mathcal{H}}_- = \sum_{i \text{ odd}} \oplus (\mathcal{H} \otimes \Lambda^i Z).$$

Note that D_0 maps $\tilde{\mathcal{H}}_+$ into $\tilde{\mathcal{H}}_-$ and vice versa. Let $D_+ = D_0|_{\tilde{\mathcal{H}}_+}$ considered as an operator in $\mathcal{B}(\tilde{\mathcal{H}}_+, \tilde{\mathcal{H}}_-)$. The relationship to curvature is the following index theorem:

Theorem 8.3 (Arveson 2002). *If T is a pure graded row contraction, then both $\ker D_+$ and $\ker D_+^*$ are finite dimensional, and*

$$(-1)^d K(\mathcal{H}) = \text{ind } D_+ = \dim \ker D_+ - \dim \ker D_+^*.$$

Stability of the Fredholm index leads to the corollary that two d -contractions T and T' , which are both Fredholm and are unitarily equivalent modulo the compact operators, have the same curvature. He observes that if the commutators $[T_i, T_j^*]$ are compact, then D_T is Fredholm.

Finally we discuss [27] in which Arveson considers pure graded finite rank d -contractions. By the dilation theorem for commuting row contractions, such a row contraction is obtained from a graded submodule M of $H_d^2 \otimes \mathbb{C}^n$, where n is finite. Graded means that M is spanned by homogeneous polynomials, and thus is generated by a finite set of homogeneous polynomials (Hilbert's basis theorem).

An easy calculation shows that the multipliers S_i on H_d^2 satisfy

$$[S_i, S_j^*] = S_i S_j^* - S_j^* S_i \in \mathfrak{S}_p \quad \text{for all } 1 \leq i, j \leq d, \quad p > d,$$

where \mathfrak{S}_p is the Schatten p -class of compact operators with s -numbers lying in ℓ^p . Examples suggest that this condition on the commutators $[T_i, T_j^*]$ should persist for graded submodules M , i.e. $T_i = S_i^{(n)}|_M$. Arveson establishes this for modules generated by monomials of the form $z_1^{k_1} \dots z_d^{k_d} \otimes \xi$. Moreover he makes the following conjecture:

Conjecture 8.4 (Arveson 2005). *If $T = [T_1, \dots, T_d]$ is a pure graded finite rank d -contraction, then $[T_i, T_j^*] \in \mathfrak{S}_p$ for all $1 \leq i, j \leq d$ and $p > d$.*

Douglas [54] further refines this conjecture first by enlarging the family of Hilbert modules considered, and more significantly by considering the ideal I of all polynomials annihilating the d -tuple T . Let Z be the zero set of this ideal. Then Douglas conjectures that the commutators $[T_i, T_j^*]$ should lie in \mathfrak{S}_p for all $p > \dim(Z \cap \mathbb{B}_d)$.

There are no known counterexamples, so both of these conjectures remain open. There has been a lot of recent interest. The best result so far, due to Guo and Wang [63], establishes Arveson's conjecture when $d = 2$ or 3 , and when M is singly generated.

Acknowledgements. The author was partially supported by an NSERC grant.

I thank Matthew Kennedy, Vern Paulsen, Gilles Pisier, David Pitts and Orr Shalit for reading the first draft of this paper and making many helpful suggestions that improved the final version.

REFERENCES

- [1] J. Agler, *An abstract approach to model theory*, in *Surveys of some recent results in operator theory*, Vol. II, pp. 1–23, Longman Sci. Tech., Harlow, 1988.
- [2] J. Agler and J. McCarthy, *Complete Nevanlinna-Pick kernels*, *J. Funct. Anal.* **175** (2000), 111–124.
- [3] J. Agler and J. McCarthy, *Pick interpolation and Hilbert function spaces*, *Graduate Studies in Mathematics* **44**, Amer. Math. Soc., Providence RI, 2002.
- [4] N. Andersen, *Compact perturbations of reflexive algebras*, *J. Funct. Anal.* **38** (1980), 366–400.
- [5] T. Ando, *On a pair of commuting contractions*, *Acta Sci. Math. (Szeged)* **24** (1963), 88–90.
- [6] A. Arias and G. Popescu, *Noncommutative interpolation and Poisson transforms*, *Israel J. Math.* **115** (2000), 205–234.
- [7] W. Arveson, *Prediction theory and group representations*, Ph.D. Thesis, University of California, Los Angeles, 1964.
- [8] W. Arveson, *Operator algebras and measure preserving automorphisms*, *Acta Math.* **118**, (1967), 95–109.
- [9] W. Arveson, *Analyticity in operator algebras*, *Amer. J. Math.* **89** (1967), 578–642.
- [10] W. Arveson, *A density theorem for operator algebras*, *Duke Math. J.* **34** (1967), 635–647.
- [11] W. Arveson, *Subalgebras of C^* -algebras*, *Acta Math.* **123** (1969), 141–224.
- [12] W. Arveson, *Unitary invariants for compact operators*, *Bull. Amer. Math. Soc.* **76** (1970), 88–91.
- [13] W. Arveson, *Subalgebras of C^* -algebras II*, *Acta Math.* **128** (1972), 271–308.
- [14] W. Arveson, *Operator algebras and invariant subspaces*, *Ann. Math.* **100** (1974), 433–532.
- [15] W. Arveson, *On groups of automorphisms of operator algebras*, *J. Funct. Anal.* **15** (1974), 217–243.
- [16] W. Arveson, *Interpolation problems in nest algebras*, *J. Funct. Anal.* **20** (1975), 208–233.
- [17] W. Arveson, *An invitation to C^* -algebras*, *Graduate Texts in Mathematics* **39**, Springer-Verlag, New York-Heidelberg, 1976.
- [18] W. Arveson, *Notes on extensions of C^* -algebras*, *Duke Math. J.* **44** (1977), 329–355.
- [19] W. Arveson, *Perturbation theory for groups and lattices*, *J. Funct. Anal.* **53** (1983), 22–73.
- [20] W. Arveson, *Ten lectures on operator algebras*, *CBMS Regional Conference Series in Mathematics*, **55**; Published for the Conference Board of the Mathematical Sciences, Washington, DC by the American Mathematical Society, Providence, RI, 1984.
- [21] W. Arveson, *Continuous analogues of Fock space*, *Mem. Amer. Math. Soc.* **80** (1989), no. 409.
- [22] W. Arveson, *Subalgebras of C^* -algebras III*, *Acta Math.* **181** (1998), 159–228.
- [23] W. Arveson, *The curvature invariant of a Hilbert module over $\mathbb{C}[z_1, \dots, z_d]$* , *J. Reine Angew. Math.* **522** (2000), 173–236.

- [24] W. Arveson, *A short course on spectral theory*, Graduate Texts in Mathematics **209**, Springer-Verlag, New York, 2002.
- [25] W. Arveson, *The Dirac operator of a commuting d -tuple*, J. Funct. Anal. **189** (2002), 53–79.
- [26] W. Arveson, *Noncommutative dynamics and E -semigroups*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2003.
- [27] W. Arveson, *p -summable commutators in dimension d* , J. Operator Theory **54** (2005), 101–117.
- [28] W. Arveson, *The noncommutative Choquet boundary*, J. Amer. Math. Soc. **21** (2008), 1065–1084.
- [29] W. Arveson and K. Josephson, *Operator algebras and measure preserving automorphisms II*, J. Funct. Anal. **4** (1969), 100–134.
- [30] J. Ball, T. Trent and V. Vinnikov, *Interpolation and commutant lifting for multipliers on reproducing kernel Hilbert spaces*, Operator theory and analysis (Amsterdam, 1997), 89–138, Oper. Theory Adv. Appl. **122**, Birkhauser, Basel, 2001.
- [31] D. Blecher and L. Labuschagne, *Characterizations of noncommutative H^∞* , Integral Equations Operator Theory **56** (2006), 301–321.
- [32] D. Blecher and L. Labuschagne, *Von Neumann algebraic H^p theory*, in Function spaces, pp. 89–114, Contemp. Math. **435**, Amer. Math. Soc., Providence, RI, 2007.
- [33] D. Blecher, and C. Le Merdy, *Operator algebras and their modules—an operator space approach*, LMS Monographs, new series **30**, Oxford University Press, Oxford, 2004.
- [34] D. Blecher, Z.J. Ruan and A. Sinclair, *A characterization of operator algebras*, J. Funct. Anal. **89** (1990), 188–201.
- [35] L. Brown, R. Douglas and P. Fillmore, *Unitary equivalence modulo the compact operators and extensions of C^* -algebras*, Proc. conference on Operator theory, Halifax, NS, Lect. Notes Math. **345**, Springer Verlag, Berlin, 1973.
- [36] L. Brown, R. Douglas and P. Fillmore, *Extensions of C^* -algebras and K -homology*, Ann. Math. **105** (1977), 265–324.
- [37] J. Bunce, *The similarity problem for representations of C^* -algebras*, Proc. Amer. Math. Soc. **81** (1981), 409–414.
- [38] J. Bunce, *Models for n -tuples of non-commuting operators*, J. Funct. Anal. **57** (1984), 21–30.
- [39] L. Carleson, *Interpolation by bounded analytic functions and the corona problem*, Ann. Math. (2) **76** (1962), 547–559.
- [40] M.D. Choi and E. Effros, *Separable nuclear C^* -algebras and injectivity*, Duke Math. J. **43** (1976), 309–322.
- [41] M.D. Choi and E. Effros, *The completely positive lifting problem for C^* -algebras*, Ann. Math. **104** (1976), 585–609.
- [42] M.D. Choi and E. Effros, *Nuclear C^* -algebras and injectivity: the general case*, Indiana Univ. Math. J. **26** (1977), 443–446.
- [43] M.D. Choi and E. Effros, *Injectivity and operator spaces*, J. Funct. Anal. **24** (1977), 156–209.

- [44] E. Christensen, *On nonselfadjoint representations of C^* -algebras*, Amer. J. Math. **103** (1981), 817–833.
- [45] A. Connes, *Une classification des facteurs de type III*, Ann. Sci. École Norm. Sup. **6** (1973), 133–252.
- [46] A. Connes, *Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$* , Ann. Math. **104** (1976), 73–115.
- [47] S. Costea, E. Sawyer and B. Wick, *The corona theorem for the Drury-Arveson Hardy space and other holomorphic Besov–Sobolev spaces on the unit ball in \mathbb{C}^n* , Anal. PDE **4** (2011), 499–550.
- [48] K. Davidson, *Similarity and compact perturbations of nest algebras*, J. Reine Angew. Math. **348** (1984), 286–294.
- [49] K. Davidson, *Nest Algebras*, Pitman Research Notes in Mathematics Series **191**, Longman Scientific and Technical Pub. Co., London, New York, 1988.
- [50] K. Davidson and E. Katsoulis, *Isomorphisms between topological conjugacy algebras*, J. Reine Angew. Math. **621** (2008), 29–51.
- [51] K. Davidson and D. Pitts, *Invariant subspaces and hyper-reflexivity for free semi-group algebras*, Proc. London Math. Soc. **78** (1999), 401–430.
- [52] K. Davidson and D. Pitts, *Nevanlinna-Pick interpolation for non-commutative analytic Toeplitz algebras*, Integral Eqtns. and Operator Theory **31** (1998), 321–337.
- [53] R. Douglas and V. Paulsen, *Hilbert modules over function algebras*, Pitman Research Notes in Math. **217**, Longman Scientific & Technical, Harlow, 1989.
- [54] R. Douglas, *A new kind of index theorem*, Analysis, geometry and topology of elliptic operators, pp. 369–382, World Scientific Publications, Hackensack, NJ, 2006.
- [55] M. Dritschel and S. McCullough, *Boundary representations for families of representations of operator algebras and spaces*, J. Operator Theory **53** (2005), 159–167.
- [56] S. Drury, *A generalization of von Neumann's inequality to the complex ball*, Proc. Amer. Math. Soc. **68** (1978), 300–304.
- [57] E. Effros and E. Lance, *Tensor products of operator algebras*, Adv. Math. **25** (1977), 1–34.
- [58] E. Effros and Z.J. Ruan, *Operator spaces*, LMS Monographs, new series **23**, Oxford University Press, New York, 2000.
- [59] A. Frazho, *Models for non-commuting operators*, J. Funct. Anal. **48** (1982), 1–11.
- [60] B. Fuglede and R. Kadison, *Determinant theory in finite factors*, Ann. Math. **55** (1952), 520–530.
- [61] J. Garnett, *Bounded analytic functions*, revised first edition, Graduate Texts in Math. **236**, Springer, New York, 2007.
- [62] D. Greene, S. Richter and C. Sundberg, *The structure of inner multipliers on spaces with complete Nevanlinna-Pick kernels*, J. Funct. Anal. **194** (2002), 311–331.
- [63] K. Guo and K. Wang, *Essentially normal Hilbert modules and K -homology*, Math. Annalen **340** (2008), 907–934.
- [64] U. Haagerup, *Solution of the similarity problem for cyclic representations of C^* -algebras*, Ann. Math. **118** (1983), 215–240.

- [65] D. Hadwin and T. Hoover, *Operator algebras and the conjugacy of transformations.*, J. Funct. Anal. **77** (1988), 112–122.
- [66] M. Hamana, *Injective envelopes of operator systems*, Publ. Res. Inst. Math. Sci. **15** (1979), 773–785.
- [67] K. Hoffman, *Analytic functions and logmodular Banach algebras*, Acta Math. **108** (1962), 271–317.
- [68] K. Hoffman and H. Rossi, *Function theory and multiplicative linear functionals*, Trans. Amer. Math. Soc. **116** (1965), 536–543.
- [69] M. Izumi, *E_0 -semigroups: around and beyond Arveson's work*, J. Operator Theory **68** (2012), xxx–xxx.
- [70] R. Kadison, *Derivations of operator algebras*, Ann. Math. **83** (1966), 280–293.
- [71] R. Kadison and I. Singer, *Triangular operator algebras*, Amer. J. Math. **82** (1960), 227–259.
- [72] L. Labuschagne, *A noncommutative Szegő theorem for subdiagonal subalgebras of von Neumann algebras*, Proc. Amer. Math. Soc. **133** (2005), 3643–3646.
- [73] E. Lance, *On nuclear C^* -algebras*, J. Funct. Anal. **12** (1973), 157–176.
- [74] D. Larson, *Nest algebras and similarity transformations*, Ann. Math. **121** (1985), 409–427.
- [75] R. Loebel and P. Muhly, *Analyticity and flows in von Neumann algebras*, J. Funct. Anal. **29** (1978), 214–252.
- [76] S. McCullough, *Carathéodory interpolation kernels*, Integral Equations Operator Theory **15** (1992), 43–71.
- [77] P. Muhly and B. Solel, *Hilbert modules over operator algebras*, Mem. Amer. Math. Soc. **117** (1995), no. 559.
- [78] P. Muhly and B. Solel, *An algebraic characterization of boundary representations*, Non-selfadjoint operator algebras, operator theory, and related topics, 189–196, Oper. Theory Adv. Appl. **104**, Birkhauser, Basel, 1998.
- [79] V. Müller and F. Vasilescu, *Standard models for some commuting multioperators*, Proc. Amer. Math. Soc. **117** (1993), 979–989.
- [80] S. Parrott, *Unitary dilations for commuting contractions*, Pacific J. Math. **34** (1970), 481–490.
- [81] V. Paulsen, *Completely bounded homomorphisms of operator algebras*, Proc. Amer. Math. Soc. **92** (1984), 225–228.
- [82] V. Paulsen, *Every completely polynomially bounded operator is similar to a contraction*, J. Funct. Anal. **55** (1984), 1–17.
- [83] V. Paulsen, *Completely bounded maps and dilations*, Pitman Research Notes in Mathematics **217**, Longman Scientific and Technical, Harlow, 1986.
- [84] V. Paulsen, *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Mathematics **78**, Cambridge University Press, Cambridge, 2002.
- [85] J. Peters, *Semicrossed products of C^* -algebras*, J. Funct. Anal. **59** (1984), 498–534.
- [86] G. Pick, *Über die Beschränkungen analytischer Funktionen, welche durch vorgegebene Funktionswerte bewirkt werden*, Math. Ann. **77** (1916), 7–23.

- [87] G. Pisier, *A polynomially bounded operator on Hilbert space which is not similar to a contraction*, J. Amer. Math. Soc. **10** (1997), 351–369.
- [88] G. Pisier, *Similarity problems and completely bounded maps*, 2nd, expanded edition, Lecture Notes in Mathematics **1618**, Springer-Verlag, Berlin, 2001.
- [89] G. Pisier, *Introduction to operator space theory*, LMS Lecture Note Series **294**, Cambridge University Press, Cambridge, 2003.
- [90] G. Popescu, *Isometric dilations for infinite sequences of noncommuting operators*, Trans. Amer. Math. Soc. **316** (1989), 523–536.
- [91] G. Popescu, *Multi-analytic operators on Fock spaces*, Math. Ann. **303** (1995), 31–46.
- [92] P. Quiggin, *For which reproducing kernel Hilbert spaces is Pick's theorem true?*, Integral Equations Operator Theory **16** (1993), 244–266.
- [93] H. Radjavi and P. Rosenthal, *On invariant subspaces and reflexive algebras*, Amer. J. Math. **91** (1969), 683–692.
- [94] J. Ringrose, *On some algebras of operators*, Proc London Math. Soc. (3) **15** (1965), 61–83.
- [95] J. Ringrose, *On some algebras of operators II*, Proc. London Math. Soc. (3) **16** (1966), 385–402.
- [96] Z.J. Ruan, *Subspaces of C^* -algebras*, J. Funct. Anal. **76** (1988), 217–230.
- [97] S. Sakai, *Derivations of W^* -algebras*, Ann. Math. **83** (1966), 273–279.
- [98] D. Sarason, *Generalized interpolation in H^∞* , Trans. Amer. Math. Soc. **127** (1967), 179–203.
- [99] T. Srinivasan and J.K. Wang, *Weak*-Dirichlet algebras*, in Function algebras, F. Birtel, ed., Scott Foresman and Co., 1966, pp. 216–249.
- [100] W. Stinespring, *Positive functions on C^* -algebras*, Proc. Amer. Math. Soc. **6** (1955), 211–216.
- [101] E. Størmer, *Positive linear maps of operator algebras*, Acta Math. **110** (1963), 233–278.
- [102] B. Sz. Nagy, *Sur les contractions de l'espace de Hilbert*, Acta Sci. Math. (Szeged) **15** (1953), 87–92.
- [103] B. Sz. Nagy and C. Foias, *Harmonic analysis of operators on Hilbert space*, North Holland Pub. Co., London, 1970.
- [104] B. Sz. Nagy, C. Foias, H. Bercovici and L. Kerchy, *Harmonic analysis of operators on Hilbert space*, 2nd ed., Springer Verlag, New York, 2010.
- [105] B. Sz. Nagy and C. Foias, *Dilatation des commutants d'opérateurs*, Comptes Rendus Paris **266** (1968), A493–495.
- [106] J. Taylor, *A joint spectrum for several commuting operators*, J. Funct. Anal. **6** (1970), 172–191.
- [107] J. Taylor, *The analytic functional calculus for several commuting operators*, Acta Math. **125** (1970), 1–38.
- [108] N. Varopoulos, *On an inequality of von Neumann and an application of the metric theory of tensor products to operators theory*, J. Funct. Anal. **16** (1974), 83–100.
- [109] D. Voiculescu, *A non-commutative Weyl-von Neumann Theorem*, Rev. Roum. Pures Appl. **21** (1976), 97–113.

[110] G. Wittstock, *Ein operatorwertiger Hahn-Banach Satz*, J. Funct. Anal. **40** (1981), 127–150.

KENNETH R. DAVIDSON, DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY
OF WATERLOO, WATERLOO, ON N2L 3G1, CANADA
E-mail address: krdauids@uwaterloo.ca

Received August 20, 2012.