

# MEET IRREDUCIBLE IDEALS AND REPRESENTATIONS OF LIMIT ALGEBRAS

KENNETH R. DAVIDSON, ELIAS KATSOULIS, AND JUSTIN PETERS

ABSTRACT. In this paper we give criteria for an ideal  $\mathcal{J}$  of a TAF algebra  $\mathcal{A}$  to be meet irreducible. We show that  $\mathcal{J}$  is meet irreducible if and only if the  $C^*$ -envelope of  $\mathcal{A}/\mathcal{J}$  is primitive. In that case,  $\mathcal{A}/\mathcal{J}$  admits a faithful nest representation which extends to a  $*$ -representation of the  $C^*$ -envelope for  $\mathcal{A}/\mathcal{J}$ . We also characterize the meet irreducible ideals as the kernels of bounded nest representations; this settles the question of whether the  $n$ -primitive and meet-irreducible ideals coincide.

## 1. INTRODUCTION

Representation theory of operator algebras is still in its infancy. While for  $C^*$ -algebras the fundamentals of representation theory have long been known, for nonselfadjoint algebras there are hardly any results of a general nature. For ‘triangular operator algebras’ (a term which we leave undefined), intuition suggests that the fundamental building blocks for representation theory should be nest representations. In the category of  $C^*$ -algebras, the nest representations are precisely the irreducible representations.

Recall that a *nest representation* is a representation for which the closed, invariant subspaces form a nest (i.e., are linearly ordered). In his study of nonselfadjoint crossed products, Lamoureux introduced the notion of  *$n$ -primitive ideal*. An ideal is  $n$ -primitive if it is the kernel of a nest representation. Lamoureux has shown that in various contexts in nonselfadjoint algebras the  $n$ -primitive ideals play a role analogous to the primitive ideals in  $C^*$ -algebras. Thus, one can give the set of  $n$ -primitive ideals the hull-kernel topology, and for every (closed, two-sided) ideal  $\mathcal{I}$  in the algebra,  $\mathcal{I}$  is the intersection of all  $n$ -primitive ideals containing  $\mathcal{I}$ ; in other words,  $\mathcal{I} = k(h(\mathcal{I}))$ .

---

1991 *Mathematics Subject Classification.* 47L80.

October 20, 2001 version.

First author partially supported by an NSERC grant.

Second author’s research partially supported by a grant from ECU.

An ideal  $\mathcal{J}$  of an algebra  $\mathcal{A}$  is *meet irreducible* if, for any ideals  $\mathcal{I}_1$  and  $\mathcal{I}_2$  containing  $\mathcal{J}$ , the relation  $\mathcal{I}_1 \cap \mathcal{I}_2 = \mathcal{J}$  implies that either  $\mathcal{I}_1 = \mathcal{J}$  or  $\mathcal{I}_2 = \mathcal{J}$ . In the case of  $T_n$ , the algebra of upper triangular  $n \times n$  matrices, meet irreducible ideals are gotten by ‘cutting a wedge’ from the algebra: let  $1 \leq i_0 \leq j_0 \leq n$ . The ideal

$$\mathcal{I} = \{(a_{ij}) : a_{ij} = 0, i_0 \leq i \leq j \leq j_0\}$$

is meet irreducible, and every meet irreducible ideal of  $T_n$  has this form.

The relationship between meet-irreducible and n-primitive ideals is studied in a variety of examples in [7], and in [3] meet-irreducible ideals in strongly maximal triangular AF-algebras are characterized by sequences of matrix units and also in terms of groupoids. In that paper it is shown that every meet-irreducible ideal is n-primitive. This is done by constructing a nest representation. The converse question, whether every n-primitive ideal is meet-irreducible, was left open.

In a recent work [2], the first two authors examined the  $C^*$ -envelope of a quotient  $\mathcal{A}/\mathcal{J}$  of a strongly maximal TAF algebra by an ideal  $\mathcal{J}$ . They showed the  $C^*$ -envelope is an AF  $C^*$  algebra, even though the quotient  $\mathcal{A}/\mathcal{J}$  is not in general a TAF algebra. It turns out that the  $C^*$ -envelope of  $\mathcal{A}/\mathcal{J}$  is sensitive enough to detect the meet irreducibility of  $\mathcal{J}$ . In Theorem 2.3 we show that  $\mathcal{J}$  is meet irreducible if and only if  $C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$ , the  $C^*$ -envelope of  $\mathcal{A}/\mathcal{J}$ , is primitive. The theory of  $C^*$ -envelopes provides the natural framework for studying results of this type. In Theorem 2.4 we show that for a meet irreducible ideal  $\mathcal{J}$ , there exists a faithful and irreducible  $*$ -representation of  $C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$ , whose restriction on  $\mathcal{A}/\mathcal{J}$  is a nest representation. Since the converse is easily seen to be true, Theorem 2.4 provides a characterization of meet irreducible ideals in terms of the representation theory for  $\mathcal{A}$ .

The question of whether the kernel of a nest representation is a meet-irreducible ideal emerged at the Ambelside, U.K. conference in summer, 1997. Subsequently some progress was made. In [4] a partial result was obtained: if the TAF algebra  $\mathcal{A}$  has totally ordered spectrum, or if the nest representation  $\pi$  has the property that the von Neumann algebra generated by  $\pi(\mathcal{A} \cap \mathcal{A}^*)$  contains an atom, then  $\ker(\pi)$  is meet-irreducible. The solution presented in Theorem 2.6 is self-contained and does not make use of the results of [3] or [4]. Thus the question is now settled for strongly maximal TAF algebras.

Despite the fact that evidence at hand is limited, it nonetheless seems worthwhile to ask

**Question 1.1.** Are there any operator algebras for which the n-primitive ideals, and the meet-irreducible ideals do not coincide?

We would like to thank Alan Hopenwasser for several suggestions, including one leading to the final version of Theorem 2.6.

## 2. THE MAIN RESULTS

We begin by recalling a result of Lamoureux [7].

**Lemma 2.1.** *Let  $\mathcal{I}$  be a closed, two-sided ideal in a separable  $C^*$ -algebra  $\mathcal{A}$ . Then the following are equivalent:*

- (i)  $\mathcal{I}$  is  $n$ -primitive
- (ii)  $\mathcal{I}$  is primitive
- (iii)  $\mathcal{I}$  is prime
- (iv)  $\mathcal{I}$  is meet-irreducible

One can actually characterize when an AF  $C^*$ -algebra is primitive in terms of its Bratteli diagram. Let  $\mathfrak{A} = \varinjlim(\mathfrak{A}_i, \varphi_i)$  be an AF  $C^*$ -algebra and assume that each  $\mathfrak{A}_i$  decomposes as a direct sum  $\mathfrak{A}_i = \oplus_j \mathfrak{A}_{i,j}$  of finite dimensional full matrix algebras  $\mathfrak{A}_{i,j}$ . A *path*  $\Gamma$  for  $\mathfrak{A} = \varinjlim(\mathfrak{A}_i, \varphi_i)$  is a sequence  $\{\mathfrak{A}_{i,j_i}\}_{i=1}^\infty$  so that for each pair of nodes  $((i, j_i), (i+1, j_{i+1}))$  there exist an arrow in the Bratteli diagram for  $\mathfrak{A} = \varinjlim(\mathfrak{A}_i, \varphi_i)$  which joins them. It is known that  $\mathfrak{A}$  is primitive iff there is a path  $\Gamma$  for  $\mathfrak{A} = \varinjlim(\mathfrak{A}_i, \varphi_i)$  so that each summand of  $\mathfrak{A}_i$  is eventually mapped into a member of  $\Gamma$ . We call such a path  $\Gamma$  an *essential path* for  $\mathfrak{A}$ .

Beyond  $C^*$ -algebras, a meet irreducible ideal need not be primitive. In [3], a description of all meet irreducible ideals of a TAF algebra was given in terms of matrix unit sequences.

**Definition 2.2.** Let  $\mathcal{A} = \varinjlim(\mathcal{A}_i, \varphi_i)$  be a TAF algebra. A sequence  $(e_i)_{i \geq N}$  of matrix units from  $\mathcal{A}$  will be called an *mi-chain* if the following two conditions are satisfied for all  $i \geq N$ :

- (A)  $e_i \in \mathcal{A}_i$ .
- (B)  $e_{i+1} \in \text{Id}_{i+1}(e_i)$ ,

where  $\text{Id}_{i+1}(e_i)$  denotes the ideal generated by  $e_i$  in  $\mathcal{A}_{i+1}$ .

If  $(e_i)_{i \geq N}$  is an mi-chain for  $\mathcal{A} = \varinjlim(\mathcal{A}_i, \varphi_i)$ , let  $\mathcal{J}$  be the join of all ideals which do not contain any matrix unit  $e_i$  from the chain. In [3, Theorem 1.2] it is shown that for a TAF algebra  $\mathcal{A} = \varinjlim(\mathcal{A}_i, \varphi_i)$ , given an mi-chain  $(e_i)_{i \geq N}$ , the ideal  $\mathcal{J}$  associated with  $(e_i)_{i \geq N}$  is meet irreducible. Conversely, every proper meet irreducible ideal in  $\mathcal{A} = \varinjlim(\mathcal{A}_i, \varphi_i)$  is induced by some mi-chain, chosen from some contraction of this representation.

In this paper we give a characterization of the meet irreducible ideals of TAF algebras in terms of  $C^*$ -envelopes of quotient algebras. We need to recall the notation and machinery from [2].

Let  $\mathfrak{A} = \varinjlim(\mathfrak{A}_i, \varphi_i)$  be the enveloping  $C^*$ -algebra for a TAF algebra  $\mathcal{A} = \varinjlim(\mathcal{A}_i, \varphi_i)$ . Let  $\mathcal{J} \subseteq \mathcal{A}$  be a closed ideal, and let  $\mathcal{J}_i := \mathcal{J} \cap \mathcal{A}_i$ . For each  $i \geq 1$ ,  $\mathcal{S}_i$  denotes the collection of all diagonal projections  $p$  which are semi-invariant for  $\mathcal{A}_i$ , are supported on a single summand of  $\mathfrak{A}_i$  and satisfy  $(p\mathcal{A}_i p) \cap \mathcal{J} = \{0\}$ . We form finite dimensional  $C^*$ -algebras

$$\mathfrak{B}_i := \sum_{p \in \mathcal{S}_i} \oplus \mathcal{B}(\text{Ran } p)$$

where  $\mathcal{B}(\text{Ran } p)$  denotes the bounded operators on  $\text{Ran } p$ . Of course,  $\mathcal{B}(\text{Ran } p)$  is isomorphic to  $\mathfrak{M}_{\text{rank } p}$ . Let  $\sigma_i$  be the map from  $\mathfrak{A}_i$  into  $\mathfrak{B}_i$  given by  $\sigma_i(a) = \sum_{p \in \mathcal{S}_i} p a p|_{\text{Ran } p}$ . The map  $\sigma_i|_{\mathcal{A}_i}$  factors as  $\rho_i q_i$  where  $q_i$  is the quotient map of  $\mathcal{A}_i$  onto  $\mathcal{A}_i/\mathcal{J}_i$  and  $\rho_i$  is a completely isometric homomorphism of  $\mathcal{A}_i/\mathcal{J}_i$  into  $\mathfrak{B}_i$ . Notice that  $\mathfrak{B}_i$  equals the  $C^*$ -algebra generated by  $\rho_i(\mathcal{A}_i/\mathcal{J}_i)$ .

We then consider unital embeddings  $\pi_i$  of  $\mathfrak{B}_i$  into  $\mathfrak{B}_{i+1}$  defined as follows. For each  $q \in \mathcal{S}_{i+1}$  we choose projections  $p \in \mathcal{S}_i$  which maximally embed into  $q$  under the action of  $\varphi_i$ . This way, we determine multiplicity one embeddings of  $\mathcal{B}(\text{Ran } p)$  into  $\mathcal{B}(\text{Ran } q)$ . Taking into account all such possible embeddings, we determine the embedding  $\pi_i$  of  $\mathfrak{B}_i$  into  $\mathfrak{B}_{i+1}$ .

Finally we form the subsystem of the directed limit  $\mathfrak{B} = \varinjlim(\mathfrak{B}_i, \pi_i)$  corresponding to all summands which are *never* mapped into a summand  $\mathcal{B}(\text{Ran } p)$  where  $p$  is a maximal element of some  $\mathcal{S}_i$ . Evidently this system is directed upwards. It is also hereditary in the sense that if every image of a summand lies in one of the selected blocks, then it clearly does not map into a maximal summand and thus already lies in our system. By [1, Theorem III.4.2], this system determines an ideal  $\mathfrak{I}$  of  $\mathfrak{B}$ . The quotient  $\mathfrak{B}' = \mathfrak{B}/\mathfrak{I}$  is the AF algebra corresponding to the remaining summands and the remaining embeddings; it can be expressed as a direct limit  $\mathfrak{B}' = \varinjlim(\mathfrak{B}'_i, \pi'_i)$ , with the understanding that  $\mathfrak{B}'_i = \oplus_j \mathfrak{B}_{i,j}$  for these remaining summands  $\mathfrak{B}_{i,j}$  of  $\mathfrak{B}_i$ . It can be seen that the quotient map is isometric on  $\mathcal{A}/\mathcal{J}$  and that  $\mathfrak{B}'$  is the  $C^*$ -envelope of  $\mathcal{A}/\mathcal{J}$ .

**Theorem 2.3.** *Let  $\mathcal{A}$  be a TAF algebra and let  $\mathcal{J} \subseteq \mathcal{A}$  be an ideal. Then  $\mathcal{J}$  is meet irreducible if and only if the algebra  $C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$  is primitive.*

**Proof.** Assume that  $\mathfrak{B}' = C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$  is primitive and let  $\Gamma = (\mathfrak{B}_{i,j_i})_{i=1}^\infty$  essential path for  $\mathfrak{B}'$ . Let  $e_i$  for  $\mathfrak{B}_{i,j_i}$  be the *characteristic matrix units* for  $\mathfrak{B}_{i,j_i}$ , i.e., the ones on the top right corner of  $\mathfrak{B}_{i,j_i}$ .

Assume that there exist ideals  $\mathcal{I}_1$  and  $\mathcal{I}_2$ , properly containing  $\mathcal{J}$ . Since  $\mathcal{I}_1$  and  $\mathcal{I}_2$  properly contain  $\mathcal{J}$ , there exist matrix units  $f_k \in \mathcal{I}_k$

with  $f_k \notin \mathcal{J}$ ,  $k = 1, 2$ . So the images of the  $f_k$  appear in the presentation for the  $C^*$ -envelope in perhaps different summands. However, the existence of an essential path  $\Gamma$  implies that some subordinates for the  $f_k$  will appear in some member of  $\Gamma$ , say  $\mathfrak{B}_{i j_i}$ , for  $i$  sufficiently large, and so  $e_i \in \mathcal{I}_1 \cap \mathcal{I}_2$ . However,  $e_i \notin \mathcal{J}$  and so  $\mathcal{J}$  is properly contained in  $\mathcal{I}_1 \cap \mathcal{I}_2$ . It follows  $\mathcal{J}$  is meet irreducible.

Conversely, assume that  $\mathcal{J}$  is meet irreducible. In light of Lemma 2.1 and the subsequent comments, it suffices to show that the trivial ideal  $\{0\}$  is meet irreducible in the  $C^*$ -envelope  $\mathfrak{B}'$ .

By way of contradiction assume that there are non-trivial ideals  $\mathcal{I}_1$  and  $\mathcal{I}_2$  of  $\mathfrak{B}'$  so that  $\mathcal{I}_1 \cap \mathcal{I}_2 = \{0\}$ . We claim that  $(\mathcal{A}/\mathcal{J}) \cap \mathcal{I}_k \neq \{0\}$ ,  $k = 1, 2$ . Indeed, any non-trivial summand of  $\mathcal{I}_k$  will eventually be mapped into a direct summand  $\mathfrak{B}_{i j_i}$  of  $\mathfrak{B}'$  corresponding to some maximal element of  $\mathcal{S}_i$ . Hence all matrix units in  $\mathfrak{B}_{i j_i}$  belong to  $\mathcal{I}_k$ , including the characteristic one. This one however also belongs to  $\mathcal{A}/\mathcal{J}$  and therefore in the intersection  $(\mathcal{A}/\mathcal{J}) \cap \mathcal{I}_k$ .

The claim shows that the zero ideal is not meet irreducible in  $\mathcal{A}/\mathcal{J}$ . By considering the pullback, this implies that  $\mathcal{J}$  is not meet irreducible in  $\mathcal{A}$ , which is the desired contradiction.  $\blacksquare$

Notice that the sequence  $(e_i)_{i=1}^\infty$  associated with the path  $\Gamma$  in the proof above satisfies the Conditions (A) and (B) of the Definition 2.2 and is therefore an mi-chain for the ideal  $\mathcal{J}$ .

**Theorem 2.4.** *If  $\mathcal{A}$  is a TAF algebra and  $\mathcal{J}$  an ideal of  $\mathcal{A}$ , then the following are equivalent:*

- (i) *There exists a faithful representation  $\tau : C_{\text{env}}^*(\mathcal{A}/\mathcal{J}) \longrightarrow \mathcal{B}(\mathcal{H})$  so that  $\tau(\mathcal{A}/\mathcal{J})$  is weakly dense in some nest algebra.*
- (ii)  *$\mathcal{J}$  is meet irreducible.*

**Proof.** Assume that (i) is valid and let  $\tau : C_{\text{env}}^*(\mathcal{A}/\mathcal{J}) \longrightarrow \mathcal{B}(\mathcal{H})$  be a faithful representation so that  $\tau(\mathcal{A}/\mathcal{J})$  is weakly dense in some nest algebra  $\text{Alg } \mathcal{N}$ . By way of contradiction assume that  $\mathcal{J}$  is not meet irreducible. Theorem 2.3 and Lemma 2.1 imply the existence of nonzero closed ideals  $\mathcal{I}_1$  and  $\mathcal{I}_2$  in  $C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$  so that  $\mathcal{I}_1 \mathcal{I}_2 = \{0\}$ . The non-zero subspaces  $[\tau(\mathcal{I}_i)\mathcal{H}]$  must be mutually orthogonal. However they are both invariant under  $\tau(\mathcal{A}/\mathcal{J})$ , and therefore belong to  $\mathcal{N}$ , a contradiction.

Conversely, assume that (ii) is valid and so, by Theorem 2.3, we know that  $C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$  is primitive. Retain the notation established in the paragraphs preceding Theorem 2.3. Hence

$$C_{\text{env}}^*(\mathcal{A}/\mathcal{J}) = \mathfrak{B}' = \varinjlim (\mathfrak{B}'_i, \pi'_i)$$

where  $\mathfrak{B}'_i = \oplus_j \mathfrak{B}_{ij}$  for the remaining summands  $\mathfrak{B}_{ij}$  of  $\mathfrak{B}_i$ . Let  $\Gamma = (\mathfrak{B}_{ij})_{i=1}^\infty$  be the essential path for  $\mathfrak{B}'$

Each  $\mathfrak{B}_{ij}$  is a full matrix algebra and therefore contains the algebra  $\mathcal{T}_{ij}$  of upper triangular matrices. Form the finite dimensional algebras  $\mathcal{T}'_i = \oplus_j \mathcal{T}_{ij}$  and consider the direct limit algebra

$$\mathcal{T}' = \varinjlim (\mathcal{T}'_i, \pi'_i),$$

where  $\pi'_i$  is as earlier. Clearly,  $\mathcal{T}'$  is a TAF algebra whose enveloping  $C^*$ -algebra is  $\mathfrak{B}'$ . Moreover,  $\mathcal{T}'$  contains  $\mathcal{A}/\mathcal{J}$ .

We define a state  $\omega$  on  $\mathfrak{B}'$  as follows. Let  $(p_i)_{i=1}^\infty$  be a sequence of diagonal projections with  $p_i \in \mathcal{T}_{ij_i}$  so that  $p_{i+1}$  is a subordinate of  $p_i$ ,  $i \in \mathbb{N}$ . We define  $\omega_i : \mathfrak{B}'_i \rightarrow \mathbb{C}$  to be the compression on  $p_i$  and we let  $\omega$  to be the direct limit  $\omega = \varinjlim \omega_i$ . Consider the GNS triple  $(\tau, \mathcal{H}, g)$  associated with the state  $\omega$ , i.e.,  $\tau$  is a representation of  $\mathfrak{B}'$  on  $\mathcal{H}$  and  $g \in \mathcal{H}$  so that  $\omega(a) = \langle \tau(a)g, g \rangle$ ,  $a \in \mathfrak{B}'$ . Since  $\omega$  is pure,  $\tau$  is irreducible. Moreover,  $p_i \in \mathfrak{B}_{ij_i}$ ,  $i \in \mathbb{N}$  and so  $\tau$  is also faithful.

An alternative presentation for  $(\tau, \mathcal{H}, g)$  was given in [8, Proposition II.2.2]. Since  $\omega$  is multiplicative on the diagonal  $\mathcal{T}' \cap (\mathcal{T}')^*$ , one considers  $\mathcal{H}$  to be  $L^2(\mathcal{X}, \mu)$ , where  $\mathcal{X}$  is the Gelfand spectrum of  $\mathcal{T}' \cap (\mathcal{T}')^*$  and  $\mu$  the counting measure on the orbit of  $\omega$  in  $\mathcal{X}$ . With these identifications, given any matrix unit  $e$ ,  $\tau(e)$  is the translation operator on  $\mathcal{X}$  defined in the paragraphs preceding [8, Theorem II.1.1].

In [8, Proposition II.2.2] it is shown that  $\tau$  maps  $\mathcal{T}'$  onto a weakly dense subset of some nest algebra. The proof of the theorem will follow if we show that the weak closure of  $\tau(\mathcal{A}/\mathcal{J})$  contains  $\tau(\mathcal{T}')$ .

A moment's reflection shows that given any contraction  $a \in \mathcal{T}'_{ij}$  and matrix units  $e_1, e_2, \dots, e_n$  and  $f_1, f_2, \dots, f_n$  in  $\mathfrak{B}'$ , there exists a contraction  $\hat{a} \in \mathcal{A}/\mathcal{J}$  so that

$$\omega(f_k^* \hat{a} e_k) = \omega(f_k^* a e_k)$$

and therefore

$$\langle \tau(\hat{a})\tau(e_k)g, \tau(f_k)g \rangle = \langle \tau(a)\tau(e_k)g, \tau(f_k)g \rangle$$

for all  $k = 1, 2, \dots, n$ . However the collection of all vectors of the form  $\tau(e)g$ , where  $e$  ranges over all matrix units of  $\mathfrak{B}'$ , forms a dense subset of  $\mathcal{H}$  and so the desired density follows.  $\blacksquare$

### Remarks 2.5.

- (1) The implication (i)  $\implies$  (ii) also follows from Theorem 2.6.
- (2) There exists a faithful representation  $\tau$  of  $C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$  in  $\mathcal{B}(\mathcal{H})$  so that  $\tau(\mathcal{A}/\mathcal{J})$  is weakly dense in a nest algebra if and only

if there is a faithful *irreducible* representation  $\tau$  of  $C_{\text{env}}^*(\mathcal{A}/\mathcal{J})$  in  $\mathcal{B}(\mathcal{H})$  so that  $\tau(\mathcal{A}/\mathcal{J})$  is weakly dense in a nest algebra.

**Theorem 2.6.** *Let  $\mathcal{A}$  be a strongly maximal TAF algebra, and let  $\pi$  be a bounded nest representation of  $\mathcal{A}$  on a Hilbert space  $\mathcal{H}$ . Then  $\ker(\pi)$  is a meet-irreducible ideal.*

**Proof.** Since a bounded representation of the diagonal masa  $\mathcal{A} \cap \mathcal{A}^*$  is completely bounded, c.f. [9, Theorem 8.7], and a completely bounded representation is similar to a completely contractive representation by [9, Theorem 8.1], we may assume that the restriction of  $\pi$  to the diagonal masa is completely contractive. It follows that the restriction of  $\pi$  to the diagonal masa is a star representation. Let  $\mathcal{J} = \ker(\pi)$ , and  $\mathcal{J}_1, \mathcal{J}_2$  be ideals in  $\mathcal{A}$  properly containing  $\mathcal{J}$ . We need to show that  $\mathcal{J}_1 \cap \mathcal{J}_2$  properly contains  $\mathcal{J}$ .

Since  $\pi$  is a nest representation, we have (after possibly interchanging  $\mathcal{J}_1$  and  $\mathcal{J}_2$ ) that

$$(0) \neq [\pi(\mathcal{J}_1)\mathcal{H}] \subseteq [\pi(\mathcal{J}_2)\mathcal{H}]$$

where  $[\mathcal{X}]$  denotes the closed subspace generated by  $\mathcal{X} \subset \mathcal{H}$ . Fix  $n \in \mathbb{N}$  and let  $u$  be a matrix unit in  $\mathcal{J}_1 \cap \mathcal{A}_n \setminus \mathcal{J}$ . Choose a vector  $h \in \mathcal{H}$  be such that  $\|\pi(u)h\| = 1$ . There exist  $m \geq n$ ,  $N \geq 1$ , matrix units  $v_t \in \mathcal{J}_2 \cap \mathcal{A}_m$  and vectors  $h_t \in \mathcal{H}$  for  $1 \leq t \leq N$  such that

$$\left\| \pi(u)h - \sum_{t=1}^N \pi(v_t)h_t \right\| < \frac{1}{4}.$$

In particular,  $\left\| \sum_{t=1}^N \pi(v_t)h_t \right\| > 3/4$ . We may assume that  $\pi(v_t) \neq 0$  for all  $t$ .

Define a projection  $E = \bigvee_{t=1}^N \pi(e_t) = \pi\left(\bigvee_{t=1}^N e_t\right)$ , where  $e_t = v_t v_t^*$  are diagonal matrix units (which need not be distinct). For all  $s, t$ ,

$$\pi(e_s)\pi(v_t) = \pi(v_s v_s^* v_t) = \begin{cases} \pi(v_t) & \text{if } v_s v_s^* = v_t v_t^* \\ 0 & \text{otherwise} \end{cases}$$

So we have  $E \sum \pi(v_t)h_t = \sum \pi(v_t)h_t$ . Now

$$\begin{aligned} \left\| E\pi(u)h - \sum_{t=1}^N \pi(v_t)h_t \right\| &= \left\| E\left(\pi(u)h - \sum_{t=1}^N \pi(v_t)h_t\right) \right\| \\ &\leq \left\| \pi(u)h - \sum_{t=1}^N \pi(v_t)h_t \right\| < \frac{1}{4}. \end{aligned}$$

Therefore  $\|E\pi(u)h\| > 1/2$ . In particular, there exists at least one  $t$ ,  $1 \leq t \leq N$ , such that  $\pi(e_t)\pi(u) \neq 0$ .

Embed  $u \in \mathcal{A}_n \hookrightarrow \mathcal{A}_m$  and decompose it as a sum  $u = \sum u_s$  of matrix units in  $\mathcal{A}_m$ . Then  $e_t u = u_s$ , for some  $s$ , and so it follows  $\pi(u_s) \neq 0$ , i.e.,  $u_s \notin \mathcal{J}$ . Thus we have identified matrix units  $u_s \in \mathcal{J}_1 \setminus \mathcal{J}$  and  $v_t \in \mathcal{J}_2 \setminus \mathcal{J}$  of  $\mathcal{A}_m$  with the same final projection. Say  $u_s = e_{ij}^{(m,r)}$  and  $v_t = e_{ik}^{(m,r)}$ . We now distinguish three cases:

- If  $j = k$ , then  $u_s = v_t \in \mathcal{J}_1 \cap \mathcal{J}_2 \setminus \mathcal{J}$ ;
- If  $j < k$ , then  $v_t = u_s e_{jk}^{(m,r)} \in \mathcal{J}_1 \cap \mathcal{J}_2 \setminus \mathcal{J}$ ;
- If  $j > k$ , then  $u_s = v_t e_{kj}^{(m,r)} \in \mathcal{J}_1 \cap \mathcal{J}_2 \setminus \mathcal{J}$ .

It follows that in all three cases  $\mathcal{J}_1 \cap \mathcal{J}_2$  properly contains  $\mathcal{J}$ . Thus  $\mathcal{J}$  is meet-irreducible. ■

## REFERENCES

- [1] K. Davidson, *C\*-algebras by example*, Fields Institute Monographs, American Mathematical Society, 1996.
- [2] K. Davidson and E. Katsoulis, *Primitive limit algebras and C\*-envelopes*, Adv. Math., to appear
- [3] A. Donsig, A. Hopenwasser, T. Hudson, M. Lamoureux and B. Solel, *Meet irreducible ideals in direct limit algebras* Math. Scand., **87**, 2000, 27–63.
- [4] Alan Hopenwasser, Justin Peters, Stephen Power, *Nest Representations of TAF Algebras*, Canad. J. Math. **52**, 2000, 1221–1234.
- [5] Michael Lamoureux, *Nest Representations and Dynamical Systems*, J. Funct. Anal. **114**, 1993, 345–376.
- [6] Michael Lamoureux, *Ideals in some continuous nonselfadjoint crossed product algebras*, J. Funct. Anal. **142**, 1996, 211–248.
- [7] Michael Lamoureux, *Some Triangular AF Algebras*, J. Operator Theory **37**, 1997, 91–109.
- [8] J. Orr and J. Peters, *Some representations of TAF algebras*, Pacific. J. Math. **167** (1995), 129–161.
- [9] V. I. Paulsen, *Completely Bounded Maps and Dilations*, Longman Scientific, New York, Wiley, 1986.
- [10] G. K. Pedersen, *C\*-Algebras and their automorphism groups*, London Mathematical Society Monograph, Academic Press, 1979.
- [11] Ș. Strătilă and D.V. Voiculescu, *Representations of AF-algebras and of the group  $U(\infty)$* , Springer Lect. Notes Math. **486**, Springer Verlag, Berlin, New York, 1975.
- [12] Michael Thelwall, *Dilation theory for subalgebras of AF algebras*, J. Operator Theory **25**, 1991, 275–282.



PURE MATH. DEPT., U. WATERLOO, WATERLOO, ON N2L-3G1, CANADA  
*E-mail address:* `krdavids@uwaterloo.ca`

MATH. DEPT., EAST CAROLINA UNIVERSITY, GREENVILLE, NC 27858, USA  
*E-mail address:* `KatsoulisE@mail.ecu.edu`

MATH. DEPT., IOWA STATE UNIVERSITY, AMES, IA 50011, USA  
*E-mail address:* `peters@iastate.edu`