ERRATUM

Erratum to: The algebraic structure of non-commutative analytic Toeplitz algebras

Kenneth R. Davidson · David R. Pitts

Published online: 25 February 2015 © Springer-Verlag Berlin Heidelberg 2015

Erratum to: Math. Ann. 311, 275–303 (1998) DOI 10.1007/s002080050188

The algebraic structure of the non-commutative analytic Toeplitz algebra \mathfrak{L}_n is developed in the original article. Some of the results fail for the case $n=\infty$, and this implies that certain other results are not established in this case. In Theorem 3.2 of the original article, we showed there is continuous surjection $\pi_{n,k}$ from $\operatorname{Rep}_k(\mathfrak{L}_n)$, the space of completely contractive representations of \mathfrak{L}_n into the $k\times k$ matrices \mathfrak{M}_k , onto the closed unit ball $\overline{\mathbb{B}_{n,k}}$ of $\mathcal{R}_n(\mathfrak{M}_k)$ by evaluation at the generators. It is further claimed that if $T=[T_1,\ldots,T_n]\in\mathcal{R}_n(\mathfrak{M}_k)$ with $\|T\|<1$, then there is a unique representation in $\pi_{n,k}^{-1}(T)$. Further information is obtained for k=1 in Theorem 3.3 of the original article. Our proof of these results is valid for $n<\infty$, however, for $n=\infty$ the uniqueness claim is incorrect. An example due to Michael Hartz (see [2, Example 2.4]) shows that $\pi_{\infty,1}^{-1}(0)$ is very large—it contains a copy of the $\beta\mathbb{N}\setminus\mathbb{N}$.

The difficulty in the proof of Theorems 3.2 and 3.3 of the original article stems from the use of the factorization A = WX used in Lemma 3.1 of the original article. In the case $n = \infty$, this factorization comes from Corollary 2.9. The problem is that the infinite sum in Corollary 2.9 converges in the strong topology, not the norm topology, so that when the representation Φ is not strongly continuous (or equivalently,

The online version of the original article can be found under doi:10.1007/s002080050188.

K. R. Davidson

Pure Math. Dept., U. Waterloo, Waterloo, ON N2L-3G1, Canada e-mail: krdavidson@math.uwaterloo.ca

D. R. Pitts (\boxtimes)

Math. & Stat. Dept., University of Nebraska, Lincoln, NE 68588, USA e-mail: dpitts@math.unl.edu



WOT-continuous), the calculation of the norm in the last lines of the proof of Lemma 3.1 is invalid.

Theorem 3.3 is used throughout Section 4 of the original article. The results of Section 4 are valid when $n < \infty$. However, we are no longer certain of the validity of the following results when $n = \infty$: Theorem 4.1, Proposition 4.3, Theorem 4.6, Theorem 4.7, and Corollary 4.12. Proposition 4.3 is used in Theorem 4.6 to establish that all automorphisms are WOT-continuous. When attention is restricted to the class of WOT-continuous automorphisms, our proofs of Theorem 4.1 and Theorem 4.7 remain valid when $n = \infty$. We do not know whether WOT-continuity is automatic.

It is worth noting that the results of Section 4 are valid for the class of *isometric* isomorphisms even when $n=\infty$. This is because any isometric automorphism is WOT-continuous. Indeed, by [3, Theorem 3.3], \mathcal{L}_n has a unique predual, \mathcal{L}_{n*} , which sits naturally inside the dual space \mathcal{L}_n^* . The uniqueness of the predual implies that Θ^* fixes the image of \mathcal{L}_{n*} , so there is a surjective isometry $\Theta_*: \mathcal{L}_{n*} \to \mathcal{L}_{n*}$ such that $\Theta = (\Theta_*)^*$. Hence Θ is weak-* continuous. By [1, Corollary 2.12], the weak-* and WOT topologies on \mathcal{L}_n coincide, so Θ is WOT-continuous.

References

- Davidson, K.R., Pitts, D.R.: Invariant subspaces and hyper-reflexivity for free semigroup algebras. Proc. Lond. Math. Soc. 78(2), 401–430 (1999)
- Davidson, K.R., Hartz, M., Shalit, O.M.: Multipliers of embedded disks. Complex Anal. Oper. Theory (2014). arXiv:1307.3204v3; doi:10.1007/s11785-014-0360-8
- Davidson, K.R., Wright, A.: Operator algebras with unique preduals. Canad. Math. Bull. 54(3), 411–421 (2011)

