

OPERATOR ALGEBRAS WITH UNIQUE PREDUALS

KENNETH R. DAVIDSON AND ALEX WRIGHT

ABSTRACT. We show that every free semigroup algebra has a (strongly) unique Banach space predual. We also provide a new simpler proof that a weak-* closed unital operator algebra containing a weak-* dense subalgebra of compact operators has a unique Banach space predual.

1. INTRODUCTION

A famous theorem of Sakai [21] showed that C^* -algebras which are dual spaces are von Neumann algebras, and the techniques showed in addition that the predual of a von Neumann algebra is unique (up to isometric isomorphism). This generalized a result of Grothendieck [14] that $L^\infty(\mu)$ has a unique predual. Ando [1] showed that the algebra H^∞ of bounded analytic functions on the unit disk also has a unique predual. More recently, Ruan [20] showed that an operator algebra with a weak-* dense subalgebra of compact operators has a unique operator space predual. He points out that some general Banach space methods of Godefroy [10, 11] in fact imply that such algebras have a unique Banach space predual. Also, Effros, Ozawa and Ruan [8] have shown that W^* TROs (corners of von Neumann algebras) have unique preduals as well.

In this note, we show that every free semigroup operator algebra has a unique predual. A free semigroup algebra is the WOT-closed unital algebra generated by n isometries with pairwise orthogonal ranges. The prototypes are the non-commutative analytic Toeplitz algebras, \mathfrak{L}_n , given by the left regular representation of the free semigroup \mathbb{F}_n^+ of words in an alphabet of n letters [18, 19, 5]. The case $n = 1$ is just H^∞ , which follows from Ando's Theorem. Our proof deals with $n \geq 2$. Once the result is established for \mathfrak{L}_n , the general case follows from the Structure Theorem for free semigroup algebras [4] and the result mentioned above of Effros, Ozawa and Ruan.

2000 *Mathematics Subject Classification.* 47L50, 46B04, 47L35.

Key words and phrases. unique predual, free semigroup algebras, CSL algebras.
First author partially supported by an NSERC grant.

It is an open problem whether $H^\infty(\Omega)$ has a unique predual when Ω is a domain in \mathbb{C}^n , even for the bidisk \mathbb{D}^2 or the unit ball \mathbb{B}_2 of \mathbb{C}^2 . In a number of ways, the algebras \mathfrak{L}_n have proven to be more tractable than their commutative counterparts when it comes to finding analogues of classical results for H^∞ in dimension one. This predual result is another case in point.

A weak-* closed operator algebra \mathfrak{A} is called *local* if the ideal of compact operators in \mathfrak{A} is weak-* dense in \mathfrak{A} . We also provide a new simpler proof that a local operator algebra has a unique predual which is inspired by Ando's proof for H^∞ . For $\mathcal{B}(\mathcal{H})$, the manipulations involving approximate identities can be omitted. So this provides an alternative to invoking Sakai's Theorem in this case. However we also provide another very simple proof for $\mathcal{B}(\mathcal{H})$ which relies neither on positivity (like Sakai) nor on the density of the compacts.

In Banach space theory, there is an extensive literature on the topic of unique preduals. We refer the reader to a nice survey paper of Godefroy [10]. For example, if X is a dual space which does not contain an isomorphic copy of ℓ^1 , then the predual is unique. Also smoothness conditions on the predual X_* such as a locally uniformly convex norm or the Radon–Nikodym property imply that it is the unique predual of X . These properties do not often apply to algebras of operators.

Another observation due to Godefroy and Talagrand [13] is that Banach spaces with property (X) have unique preduals. This technical condition will be defined in the next section. In the proofs of Sakai and Ando mentioned above, it is a property very close to this which is exploited to establish uniqueness. It implies, for example, that if X is an M-ideal in X^{**} , then X^* is the unique predual of X^{**} [15, p.148]. This is the case for operator algebras with a weak-* dense ideal of compact operators. Recently, Pfitzner [17] has generalized this by showing that if X_* is a separable L-summand in X^* , then X_* has property (X) and so is the unique predual of X . Another basic class with unique predual are the spaces of operators $\mathcal{B}(X, Y)$ where X and Y are Banach spaces with the Radon–Nikodym property due to Godefroy and Saphar [12]. This includes spaces which are separable dual spaces, and all reflexive Banach spaces.

Nevertheless, in operator algebras, the literature on unique preduals is rather limited, and the main results have all been mentioned above.

A Banach space X has a strongly unique predual if there is a unique subspace E of X^* for which $X = E^*$. All known examples of Banach spaces with unique predual actually have a strongly unique predual [10]. This is the case in our examples as well.

We thank Gilles Godefroy for some helpful comments on this paper.

2. BACKGROUND

If X is a dual Banach space, then any predual E sits in a canonical manner as a subspace of the dual X^* . Let $\sigma(X, E)$ denote the weak-* topology on X induced from E . E has two evident properties which are characteristic:

- (1) E norms X : $\sup\{|\varphi(x)| : \varphi \in E, \|\varphi\| \leq 1\} = \|x\|$.
- (2) The closed unit ball of X is compact in the $\sigma(X, E)$ topology.

The latter property is a consequence of the Banach-Alaoglu Theorem.

Conversely, if E is a subspace of X^* with these properties, then X sits isometrically as a subspace of E^* by (1). By (2), the closed ball $\overline{b_1(X)} := \overline{b_1(0)}$ of X is weak-* compact in E^* . Therefore by the Krein–Smulyan Theorem, X is weak-* closed in E^* . However, as E is a subspace of X^* , the annihilator of X in E is $\{0\}$. Hence $X = E^*$. Thus we see that these two properties characterize the preduals of X .

In any weak-* topology on X , closed balls $\overline{b_r(x)}$ are compact for $x \in X$ and $r \geq 0$. Also, addition is always weak-* continuous. So finite sums of closed balls are also *universally weak-* compact*. This can sometimes be used to show that certain functionals are *universally weak-* continuous*, meaning that they belong to every predual of X .

Example 2.1. It is very easy to see that ℓ^∞ has a unique predual, namely ℓ^1 . Let e_n denote the sequence with a 1 in the n th coordinate, and 0 elsewhere. And let δ_n be the element of ℓ^1 which evaluates the n th coordinate. Observe that

$$\overline{b_1(e_n)} \cap \overline{b_1(-e_n)} = \overline{b_1(\ker \delta_n)}.$$

Hence $\overline{b_1(\ker \delta_n)}$ is universally weak-* compact. By the Krein–Smulyan Theorem, $\ker \delta_n$ is universally weak-* closed. So δ_n lies in every predual of ℓ^∞ . But these functionals span ℓ^1 , and hence it is the unique predual.

Example 2.2. A similar, but somewhat more involved, argument shows that $\mathcal{B}(\mathcal{H})$ has a unique predual. Let \mathcal{H} be an infinite dimensional Hilbert space, and consider unit vectors $x, y \in \mathcal{H}$. Write xy^* for the rank one operator $xy^*(z) = \langle z, y \rangle x$. Observe that

$$\begin{aligned} \mathcal{C}_x^y &:= \overline{b_1(xy^*)} \cap \overline{b_1(-xy^*)} \\ &= \{T \in \mathcal{B}(\mathcal{H}) : Ty = T^*x = 0 \text{ and } \|T\| \leq 1\}. \end{aligned}$$

Indeed, $Ty \in \overline{b_1(x)} \cap \overline{b_1(-x)} = \{0\}$ and similarly, $T^*x = 0$; so the result follows. Pick a unit vector z orthogonal to both x, y . Then a simple

calculation shows that

$$(\mathcal{C}_x^y + \mathcal{C}_z^y + \mathcal{C}_x^z) \cap \overline{b_1(\mathcal{B}(\mathcal{H}))} = \{T \in \mathcal{B}(\mathcal{H}) : \langle Ty, x \rangle = 0 \text{ and } \|T\| \leq 1\}.$$

Arguing as before, the functional $(yx^*)(T) = \langle Ty, x \rangle$ is universally weak-* continuous. But these functionals span the trace class operators \mathfrak{S}_1 , the standard predual of $\mathcal{B}(\mathcal{H})$. Therefore \mathfrak{S}_1 is the unique predual of $\mathcal{B}(\mathcal{H})$.

Example 2.3. There are WOT-closed operator algebras which do not have unique preduals. The basic point is that being an operator algebra is not restrictive. If we put any WOT-closed subspace of $\mathcal{B}(\mathcal{H})$ in the 1, 2 entry of 2×2 matrices over $\mathcal{B}(\mathcal{H})$, then we have an operator algebra. Adding in the scalar operators (on the diagonal) will not essentially change the Banach space characteristics, but will yield a unital algebra. In particular, let's put ℓ^∞ into $\mathcal{B}(\mathcal{H})$ as the diagonal operators, and place it in the 1, 2 entry. Every dual space X^* with separable predual can be isometrically imbedded into ℓ^∞ as a weak-* closed subspace. The weak-* and WOT-topologies coincide on ℓ^∞ . So this procedure yields a WOT-closed algebra. If we do this for $X^* = \ell^1$, we obtain the desired example.

A series (x_n) in a Banach space X is universally weakly Cauchy if $\sum_{n \geq 1} |\varphi(x_n)| < \infty$ for every $\varphi \in X^*$. If $X = E^*$, define $C(E)$ to be the set of all functionals $\varphi \in X^* = E^{**}$ with the property that for every universally weakly Cauchy series (x_n) in X ,

$$\varphi(w^*\text{-}\lim \sum_{i=1}^n x_i) = \sum_{i=1}^{\infty} \varphi(x_i).$$

Evidently this contains E . But Godefroy and Talagrand [13] show that $C(E)$ contains every predual of X , and this space does not depend on the choice of the predual E . They say that X has property (X) if $C(E) = E$. Evidently this immediately implies that X has a unique predual.

A similar property was established by Sakai for a von Neumann algebra \mathfrak{M} . He shows that a state φ on \mathfrak{M} belongs to \mathfrak{M}_* if and only if it satisfies: whenever (P_n) are pairwise orthogonal projections in \mathfrak{M} such that $\text{sOT-}\sum P_n = I$, then $\sum \varphi(P_n) = 1$. In Section 4, we use a similar property to establish unique preduals for algebras with sufficiently many compact operators.

3. FREE SEMI-GROUP ALGEBRAS

A free semigroup algebra is a WOT-closed unital operator algebra generated by n isometries S_1, \dots, S_n with pairwise orthogonal range. We allow $n = \infty$. The prototype is obtained from the left regular representation of the free semigroup \mathbb{F}_n^+ of all words in an alphabet of n letters. The operators L_v , for $v \in \mathbb{F}_n^+$, act on the Fock space $\ell^2(\mathbb{F}_n^+)$, with orthonormal basis $\{\xi_w : w \in \mathbb{F}_n^+\}$, by $L_v \xi_w = \xi_{vw}$. The algebra \mathfrak{L}_n generated by L_1, \dots, L_n is called the *noncommutative analytic Toeplitz algebra* because the case $n = 1$ yields the analytic Toeplitz algebra isometrically isomorphic to H^∞ , and because these algebras share many similar properties (see [18, 19, 5, 6]). The standard predual of \mathfrak{L}_n is the space \mathfrak{L}_{n*} of all weak-* continuous linear functionals on \mathfrak{L}_n . See [3] for an overview of these algebras.

Let $|w|$ denote the length of the word w . Note that the operators $\{L_w : |w| = k\}$ are isometries with pairwise orthogonal ranges. Thus $\text{span}\{L_w : |w| = k\}$ is isometric to a Hilbert space.

An element $A \in \mathfrak{L}_n$ is determined by $A\xi_\emptyset = \sum a_w L_w$. We call the series $\sum a_w L_w$ the Fourier series of A . As in classical harmonic analysis, this series need not converge. However the Cesaro means do converge in the strong operator topology to A [5]. The functional $\varphi_w(A) = \langle A\xi_\emptyset, \xi_w \rangle$ reads off the w -th Fourier coefficient, and it is evidently weak-* continuous. The ideal $\mathfrak{L}_n^0 = \ker \varphi_\emptyset$ consists of all elements with 0 constant term, and is the WOT-closed ideal generated by L_1, \dots, L_n . The powers $(\mathfrak{L}_n^0)^k$ are the ideals of elements for which $a_w = 0$ for all $|w| < k$.

We first deal with the noncommutative analytic Toeplitz algebras, \mathfrak{L}_n , for $n \geq 2$. This first lemma is motivated by the fact that for a vector v in a Hilbert space \mathcal{H} , $\bigcap_{\lambda \in \mathbb{C}} \overline{b_{\sqrt{1+|\lambda|^2}}(\lambda v)} = \mathbb{C}v^\perp \cap \overline{b_1}$.

Lemma 3.1. *For $n \geq 2$ and $k \geq 0$, the ideal $(\mathfrak{L}_n^0)^k$ is universally weak-* closed in \mathfrak{L}_n .*

Proof. Fix a word $w \in \mathbb{F}_n^+$. Define $C_w = \bigcap_{\lambda \in \mathbb{C}} \overline{b_{\sqrt{1+|\lambda|^2}}(\lambda L_w)}$. By the remarks in the previous section, this is universally weak-* compact. We will first establish that

$$C_w = \{A \in \mathfrak{L}_n : \|A\| \leq 1 \text{ and } L_w^* A = 0\}.$$

That is, a contraction A belongs to C_w if and only if A and L_w have orthogonal ranges. Observe that a contraction A lies in C_w if and only

if for all $\lambda \in \mathbb{C}$ and all $\xi \in \ell^2(\mathbb{F}_n^+)$ with $\|\xi\| = 1$,

$$\begin{aligned} 1 + |\lambda|^2 &\geq \|\lambda L_w \xi - A\xi\|^2 \\ &= \|\lambda L_w \xi\|^2 - 2\operatorname{Re}\langle A\xi, \lambda L_w \xi \rangle + \|A\xi\|^2 \\ &= |\lambda|^2 - 2\operatorname{Re}\langle \bar{\lambda} L_w^* A\xi, \xi \rangle + \|A\xi\|^2. \end{aligned}$$

If $L_w^* A = 0$ this inequality is clearly satisfied, so A belongs to C_w .

Conversely, if $A \in C_w$, by picking the sign of λ appropriately, we obtain that

$$1 + |\lambda|^2 \geq |\lambda|^2 + 2|\lambda| |\langle L_w^* A\xi, \xi \rangle| + \|A\xi\|^2.$$

Letting $|\lambda|$ tend to ∞ , we see that $\langle L_w^* A\xi, \xi \rangle = 0$ for all ξ . By the polarization identity, $L_w^* A = 0$.

It follows that $D_{k,i} := \bigcap_{|w|=k-1} C_{wi}$ is universally weak-* compact for any i . If $A \in \mathfrak{L}_n$ has a Fourier series $A \sim \sum a_v L_v$ and lies in $D_{k,i}$, then we claim that $a_v = 0$ if $|v| < k$ or if v has the form wiv' for $|w| = k-1$. Indeed, if $|v| < k$, choose any word v' , possibly empty, so that $|vv'i| = k$. Then since the range of A is orthogonal to the range of $L_{vv'i}$,

$$0 = \langle A\xi_{v'i}, L_{vv'i}\xi_{\emptyset} \rangle = \left\langle \sum a_v \xi_{vv'i}, \xi_{vv'i} \right\rangle = a_v.$$

Similarly if $v = wiv'$ for $|w| = k-1$, then since the range of A is orthogonal to the range of L_{wi} ,

$$0 = \langle A\xi_{\emptyset}, L_{wi}\xi_{v'} \rangle = \left\langle \sum a_v \xi_v, \xi_v \right\rangle = a_v.$$

Conversely, it is evident that any such A has range orthogonal to all ranges L_{wi} for $|w| = k-1$. So if it is a contraction, it will lie in $D_{k,i}$.

Since addition is always weak-* continuous, we obtain that

$$D_k = \overline{b_1(\mathfrak{L}_n)} \cap (D_{k,1} + D_{k,2})$$

is also universally weak-* compact. We claim that $D_k = \overline{b_1((\mathfrak{L}_n^0)^k)}$. For A to lie in either $D_{k,i}$, the Fourier coefficients $a_v = 0$ for $|v| < k$. So this persists in the sum, and hence D_k is contained in $\overline{b_1((\mathfrak{L}_n^0)^k)}$.

Conversely, if $A \in \overline{b_1((\mathfrak{L}_n^0)^k)}$, by [4, Lemma 2.6] there is a factorization $A = \sum_{|w|=k} L_w A_w$ where $A_w \in \mathfrak{L}_n$. Moreover, this factors as LC where L is the row operator with coefficients L_w for $|w| = k$ and C is the column operator with coefficients A_w . Since L is an isometry, we have $\|C\| = \|A\| \leq 1$. Define

$$B_1 = \sum_{i \geq 2} \sum_{|w|=k-1} L_{wi} A_{wi} \quad \text{and} \quad B_2 = \sum_{|w|=k-1} L_{w1} A_{w1}.$$

Then it follows that both B_i are contractions in $(\mathfrak{L}_n^0)^k$. Moreover, $B_1 \in D_{k,1}$ and $B_2 \in D_{k,2}$. Therefore $A = B_1 + B_2$ belongs to D_k as claimed.

We have shown that $\overline{b_1((\mathfrak{L}_n^0)^k)}$ is universally weak-* compact. Thus by the Krein–Smulyan Theorem, $(\mathfrak{L}_n^0)^k$ is universally weak-* closed. ■

Corollary 3.2. *The functionals φ_w , for $w \in \mathbb{F}_n^+$, are universally weak-* continuous for all $n \geq 2$.*

Proof. If $\mathfrak{L}_n = E^*$, let $E_k = ((\mathfrak{L}_n^0)^k)^\perp$ be the annihilator of $(\mathfrak{L}_n^0)^k$ in E . Since $(\mathfrak{L}_n^0)^k$ is $\sigma(\mathfrak{L}_n, E)$ closed, $(\mathfrak{L}_n^0)^k = E_k^\perp$ and $(\mathfrak{L}_n^0)^k \simeq (E/E_k)^*$. Therefore $(\mathfrak{L}_n^0)^k/(\mathfrak{L}_n^0)^{k+1} \simeq (E_k/E_{k+1})^*$.

Now $(\mathfrak{L}_n^0)^k/(\mathfrak{L}_n^0)^{k+1}$ is isometrically isomorphic to the subspace $\text{span}\{L_w : |w| = k\}$. Indeed, the elements of this quotient have the form $\sum_{|w|=k} a_w L_w + (\mathfrak{L}_n^0)^{k+1}$. So the norm is bounded above by

$$\left\| \sum_{|w|=k} a_w L_w \right\| = \|(a_w)_{|w|=k}\|_2.$$

On the other hand, it is bounded below by

$$\inf_{A \in (\mathfrak{L}_n^0)^{k+1}} \left\| \left(\sum_{|w|=k} a_w L_w + A \right) \xi_\emptyset \right\| = \left\| \sum_{|w|=k} a_w \xi_w \right\| = \|(a_w)_{|w|=k}\|_2$$

Hence this quotient is a Hilbert space.

Since a Hilbert space is reflexive, its dual is E_k/E_{k+1} . Therefore $\mathfrak{L}_n/(\mathfrak{L}_n^0)^{k+1}$ is reflexive with dual E/E_{k+1} . Since the functionals φ_w for $|w| \leq k$ are continuous on this quotient, they are all $\sigma(\mathfrak{L}_n, E)$ continuous. ■

Theorem 3.3. *\mathfrak{L}_n has a unique predual for $n \geq 2$.*

Proof. Recall that the standard predual \mathfrak{L}_{n*} of \mathfrak{L}_n consists of the weak-* continuous linear functionals of the form $[xy^*]$. Clearly taking x and y to be in the algebraic span of $\{\xi_w : w \in \mathbb{F}_n^+\}$ is norm dense in the predual. However, $[\xi_v \xi_w^*] = [\xi_\emptyset (L_v^* \xi_w)^*]$. So the span of the functionals $[\xi_\emptyset \xi_w^*]$ is norm dense in the predual. Since $[\xi_\emptyset \xi_w^*] = \varphi_w$ is universally weak-* continuous by Corollary 3.2, it follows that the standard predual is universally weak-* continuous.

No two preduals are comparable; so it follows that \mathfrak{L}_{n*} is the strongly unique predual of \mathfrak{L}_n . ■

To deal with the case of a general free semigroup algebra, we require a simple lemma. The dual space of an operator algebra \mathfrak{A} is a bimodule over \mathfrak{A} with the natural action $(A\varphi B)(T) = \varphi(BTA)$.

Lemma 3.4. *Let P be an orthogonal projection in an operator algebra \mathfrak{A} . Then for $\varphi \in \mathfrak{A}^*$,*

$$\|\varphi\|^2 \geq \|P\varphi\|^2 + \|P^\perp\varphi\|^2.$$

Proof. Find $A = AP$ and $B = BP^\perp$ in \mathfrak{A} of norm 1 so that $\varphi(A)$ and $\varphi(B)$ are real, and we have the approximations

$$\varphi(A) = (P\varphi)(A) \approx \|P\varphi\| \quad \text{and} \quad \varphi(B) = P^\perp\varphi(B) \approx \|P^\perp\varphi\|.$$

Consider $T = \cos \theta A + \sin \theta B$. Note that

$$\|T\|^2 = \|TT^*\| = \|\cos^2 AA^* + \sin^2 BB^*\| \leq 1.$$

Compute

$$\varphi(T) = \cos \theta \varphi(A) + \sin \theta \varphi(B) = (\cos \theta, \sin \theta) \cdot (\varphi(A), \varphi(B)).$$

Choosing θ so $(\cos \theta, \sin \theta)$ is parallel to $(\varphi(A), \varphi(B))$, we obtain

$$\|\varphi\|^2 \geq |\varphi(T)|^2 = \varphi(A)^2 + \varphi(B)^2 \approx \|P\varphi\|^2 + \|P^\perp\varphi\|^2. \quad \blacksquare$$

Now the general free semigroup algebra case follows from the structure theory of these algebras.

Theorem 3.5. *Every free semigroup algebra \mathfrak{S} has a strongly unique predual.*

Proof. We invoke the Structure Theorem for free semigroup algebras [4]. If \mathfrak{S} is a von Neumann algebra, then the result follows from Sakai's Theorem [21]. Otherwise, the WOT-closed ideal \mathfrak{S}_0 generated by $\{S_1, \dots, S_n\}$ is proper. Let $\mathfrak{J} = \bigcap_{k \geq 1} \mathfrak{S}_0^k$. This is a WOT-closed ideal of \mathfrak{S} , and is a left ideal in the von Neumann algebra \mathfrak{W} generated by $\{S_1, \dots, S_n\}$. There is a projection $P \in \mathfrak{S}$ so that $\mathfrak{J} = \mathfrak{W}P$, $P^\perp \mathcal{H}$ is invariant for \mathfrak{S} , and $\mathfrak{S}|_{P^\perp \mathcal{H}}$ is completely isometrically isomorphic and weak-* homeomorphic to \mathfrak{L}_n .

Now, define $C_{P^\perp} = \bigcap_{\lambda \in \mathbb{C}} \overline{B_{\sqrt{1+|\lambda|^2}}(\lambda P^\perp)}$. We claim $C_{P^\perp} = \overline{b_1(\mathfrak{J})}$. By the calculation of C_\emptyset in \mathfrak{L}_n in Lemma 3.1, $P^\perp C_{P^\perp} P^\perp = C_\emptyset = \{0\}$. So $C_{P^\perp} \subset \mathfrak{J} \cap \overline{B_1(\mathfrak{L}_n)} = b_1(\mathfrak{J})$. Conversely,

$$\begin{aligned} \left\| \begin{bmatrix} A & 0 \\ B & \lambda I \end{bmatrix} \right\|^2 &= \left\| \begin{bmatrix} A & 0 \\ B & \lambda I \end{bmatrix} \begin{bmatrix} A^* & B^* \\ 0 & \overline{\lambda} I \end{bmatrix} \right\| \\ &\leq \left\| \begin{bmatrix} A & 0 \\ B & 0 \end{bmatrix} \begin{bmatrix} A^* & B^* \\ 0 & 0 \end{bmatrix} \right\| + \left\| \begin{bmatrix} 0 & 0 \\ 0 & |\lambda|^2 I \end{bmatrix} \right\| \\ &= \left\| \begin{bmatrix} A & 0 \\ B & 0 \end{bmatrix} \right\|^2 + |\lambda|^2. \end{aligned}$$

Thus, we see that C_{P^\perp} contains $\overline{b_1(\mathfrak{J})}$. Whence $C_{P^\perp} = \overline{b_1(\mathfrak{J})}$.

Now C_{P^\perp} is universally weak-* compact. By the Krein-Smulyan Theorem, $\mathfrak{J} = \text{span } C_{P^\perp}$ is universally weak-closed. Note, \mathfrak{J} is a W*TRO, and therefore has a strongly unique predual [8].

Let E be a predual of \mathfrak{S} . Then the predual of $\mathfrak{S}/\mathfrak{J}$ is

$$E_0 = \{\varphi \in E : \varphi|_{\mathfrak{J}} = 0\}.$$

Since $\mathfrak{S}/\mathfrak{J}$ is isomorphic to \mathfrak{L}_n , Theorem 3.3 implies that E_0 coincides with the weak-* continuous functionals on \mathfrak{L}_n . Because the isomorphism of $\mathfrak{S}|_{P^\perp \mathcal{H}}$ to \mathfrak{L}_n is a weak-* homeomorphism, E_0 coincide with the weak-* continuous functionals on $\mathfrak{S}|_{P^\perp \mathcal{H}}$.

The predual of \mathfrak{J} is E/E_0 . If $\varphi \in E$, then $\|\varphi + E_0\| = \|\varphi|_{\mathfrak{J}}\| = \|P\varphi\|$. Clearly every functional $\psi \in E_0$ has $\psi = P^\perp \psi$. Therefore by Lemma 3.4, we see that

$$\|P\varphi\|^2 = \|\varphi + E_0\|^2 \geq \|P\varphi\|^2 + \text{dist}(P^\perp \varphi, E_0)^2.$$

Hence $P^\perp \varphi \in E_0$ and so $P\varphi \in E$. It follows that $E_0 = P^\perp E$ and $E/E_0 \simeq PE$. Hence $\mathfrak{S}P^\perp = (PE)^\perp$ is also $\sigma(\mathfrak{S}_n, E)$ closed.

As PE is the unique predual of $\mathfrak{S}P$ and $P^\perp E$ is the unique predual of $\mathfrak{S}P^\perp$, both consisting of the weak-* continuous functionals, we deduce that E necessarily coincides with the weak-* continuous functionals on \mathfrak{S} . So there is a strongly unique predual. \blacksquare

4. OPERATOR ALGEBRAS WITH MANY COMPACT OPERATORS

Suppose that \mathfrak{A} is a local weak-* closed unital sub-algebra of $\mathcal{B}(\mathcal{H})$, meaning that $\mathfrak{A} \cap \mathfrak{K}$ is weak-* dense in \mathfrak{A} . We will provide a new proof that \mathfrak{A} has a unique predual using an argument modelled on Ando's argument [1] that H^∞ has a unique predual.

Observe that the weak-* density means that $(\mathfrak{A} \cap \mathfrak{K})^\perp = \mathfrak{A}_\perp$ in the space \mathfrak{S}_1 of trace class operators. Thus $(\mathfrak{A} \cap \mathfrak{K})^* \simeq \mathfrak{S}_1/\mathfrak{A}_\perp \simeq \mathfrak{A}_*$; and hence $(\mathfrak{A} \cap \mathfrak{K})^{**} = \mathfrak{A}$. Thus there is a canonical contractive projection \mathcal{P} of the triple dual, \mathfrak{A}^* , onto the dual space \mathfrak{A}_* given by restriction to $\mathfrak{A} \cap \mathfrak{K}$, and considered as a subspace of \mathfrak{A}^* . For $\varphi \in \mathfrak{A}^*$, we will write $\mathcal{P}\varphi =: \varphi_a$ and $(\text{id} - \mathcal{P})\varphi =: \varphi_s$.

In fact by [7], $\mathfrak{A} \cap \mathfrak{K}$ is an M-ideal in \mathfrak{A} . Therefore \mathcal{P} is an L-projection, meaning that $\|\varphi\| = \|\varphi_a\| + \|\varphi_s\|$. We will not require this fact in our proof.

By Goldstine's Theorem, the unit ball of $\mathfrak{A} \cap \mathfrak{K}$ is weak* dense in the unit ball of \mathfrak{A} . Since the closed convex sets in WOT and SOT* topologies coincide, the ball of compact operators is also SOT* dense in the ball of \mathfrak{A} . In particular, there is a net (a sequence when \mathcal{H} is separable) $K_n \in \mathfrak{A} \cap \mathfrak{K}$ with $\|K_n\| \leq 1$ and $\text{SOT}^*\text{-}\lim K_n = I$. Evidently, this is a contractive approximate identity for $\mathfrak{A} \cap \mathfrak{K}$.

We require a somewhat better approximate identity. It is shown in [7] that the existence of a bounded one-sided approximate identity implies the existence of a contractive two-sided approximate identity C_n with the additional property that $\limsup \|I - C_n\| \leq 1$. What we require here is similar, and the argument follows from tricks using the Riesz functional calculus.

For various interesting examples such as $\mathcal{B}(\mathcal{H})$ and atomic CSL algebras (see the end of this section for definitions), the compact operators in the algebra have a bounded approximate identity consisting of finite rank projections $\{P_n\}$. In this case, one may take $S_n = C_n = P_n$ in Lemma 4.1, and avoid all of the tricky calculations.

Lemma 4.1. *Let \mathfrak{A} be an operator algebra with a contractive approximate identity $\{K_n\}$. Then \mathfrak{A} has a contractive approximate identity $\{S_n\}$ and a bounded approximate identity $\{I - C_n\}$ so that*

$$\lim_{n \rightarrow \infty} \left\| \begin{bmatrix} S_n & C_n \end{bmatrix} \right\| = 1 = \lim_{n \rightarrow \infty} \left\| \begin{bmatrix} S_n \\ C_n \end{bmatrix} \right\|.$$

Proof. It is easy to see that for any fixed i , $\{K_n^i\}$ is a contractive approximate identity. So if p is a polynomial p with $p(0) = 0$ and $p(1) = 1$, then $\{p(K_n)\}$ is an approximate identity. By von Neumann's inequality, it is bounded by $\|p\|_\infty = \sup_{|z| \leq 1} |p(z)|$. Now if $f \in A(\mathbb{D})$ satisfies $f(0) = 0$ and $f(1) = 1$, then it can be uniformly approximated by such polynomials. So again von Neumann's inequality shows that $\{f(K_n)\}$ is an approximate identity bounded by $\|f\|_\infty$.

Now $\sin(\pi z/2)$ is analytic, and takes $(-1, 1)$ into itself. It is easy to check that there is a convex open set U containing $(-1, 1)$ on which $|\sin(\pi z/2)| < 1$. Let $U_\varepsilon = U \cap \{x + iy : |y| < \varepsilon\}$; and let f_ε be the conformal map of \mathbb{D} onto U_ε such that $f_\varepsilon(0) = 0$ and $f_\varepsilon(1) = 1$. We define

$$g_\varepsilon(z) = \sin\left(\frac{\pi}{2} f_\varepsilon(z)\right) \quad \text{and} \quad h_\varepsilon(z) = \cos\left(\frac{\pi}{2} f_\varepsilon(z)\right).$$

Define $S_n = g_\varepsilon(K_n)$ and $C_n = h_\varepsilon(K_n)$. Then $\{S_n\}$ is a contractive approximate identity, and $\{I - C_n\}$ is a bounded approximate identity for \mathfrak{A} . By the matrix version of von Neumann's inequality,

$$\left\| \begin{bmatrix} S_n & C_n \end{bmatrix} \right\| \leq \left\| \begin{bmatrix} g_\varepsilon & h_\varepsilon \end{bmatrix} \right\| = \left\| \sqrt{|g_\varepsilon|^2 + |h_\varepsilon|^2} \right\| \leq \left(1 + 2 \sinh^2\left(\frac{\pi\varepsilon}{2}\right)\right)^{1/2}.$$

Similarly,

$$\left\| \begin{bmatrix} S_n \\ C_n \end{bmatrix} \right\| \leq \left(1 + 2 \sinh^2\left(\frac{\pi\varepsilon}{2}\right)\right)^{1/2}.$$

Now use a diagonal argument to let ε go to 0 slowly relative to n so as to still be an approximate identity and obtain the desired norm limit. \blacksquare

We can now prove the uniqueness of preduals.

Theorem 4.2. *Let \mathfrak{A} be a weak-* closed unital subalgebra of $\mathcal{B}(\mathcal{H})$ with a weak-* dense subalgebra of compact operators. Then \mathfrak{A} has a strongly unique predual.*

Proof. By the remarks at the beginning of this section, $\mathfrak{A} \cap \mathfrak{K}$ has a contractive approximate identity. So by Lemma 4.1, we obtain approximate identities $\{S_n\}$ and $\{I - C_n\}$ for $\mathfrak{A} \cap \mathfrak{K}$ as described. In particular, S_n converges SOT* to I and C_n converges SOT* to 0.

Let E be a subspace of \mathfrak{A}^* which norms \mathfrak{A} , and so that the closed unit ball of \mathfrak{A} is $\sigma(\mathfrak{A}, E)$ compact. Fix $A \in \mathfrak{A}$ with $\|A\| = 1$. Then the sequence $C_n A C_n$ is bounded, and converges SOT* to 0. Since this is a bounded net, it has a subnet $C_\alpha A C_\alpha$ which converges in the $\sigma(\mathfrak{A}, E)$ topology to some element B in the ball of \mathfrak{A} . That is,

$$\lim \varphi(C_\alpha A C_\alpha) = \varphi(B) \quad \text{for all } \varphi \in E.$$

We will show that $B = 0$. Fix $n \geq 1$. Then for $\varphi \in E$ with $\|\varphi\| = 1$,

$$\begin{aligned} |\varphi(S_n \pm B)| &= \lim_\alpha |\varphi(S_n \pm C_\alpha A C_\alpha)| \\ &\leq \lim_\alpha \left\| \begin{bmatrix} S_\alpha & C_\alpha \end{bmatrix} \begin{bmatrix} S_n & 0 \\ 0 & \pm A \end{bmatrix} \begin{bmatrix} S_\alpha \\ C_\alpha \end{bmatrix} \right\| + \|S_n - S_\alpha S_n S_\alpha\| \\ &\leq 1 + \lim_\alpha \|S_n - S_\alpha S_n S_\alpha\| = 1. \end{aligned}$$

Since E norms \mathfrak{A} , we conclude that $\|S_n \pm B\| \leq 1$. Letting n go to infinity, this converges WOT to $I \pm B$. Hence $\|I \pm B\| \leq 1$. Therefore $B = 0$.

Fix $\varphi \in E$. We have the decomposition $\varphi = \varphi_a + \varphi_s$. Note that $A - C_\alpha A C_\alpha = (1 - C_\alpha)A + C_\alpha A(1 - C_\alpha)$ is compact because $1 - C_\alpha$ is compact, and this net converges WOT to A . Hence

$$\begin{aligned} \varphi(A) &= \varphi(A - B) = \lim \varphi(A - C_\alpha A C_\alpha) \\ &= \lim \varphi_a(A - C_\alpha A C_\alpha) = \varphi_a(A). \end{aligned}$$

It follows that $\varphi = \varphi_a$. This shows that E is contained in \mathfrak{A}_* . Since it separates points, $E = \mathfrak{A}_*$. \blacksquare

A nest algebra is the set of all operators that are upper triangular with respect to a fixed chain of invariant subspaces. All nest algebras have a dense subalgebra of compact operators [9]. More generally, a CSL algebra is a reflexive algebra of operators containing a masa. The

lattice of invariant subspaces is a sublattice of the projection lattice of the masa, hence the name commutative subspace lattice (CSL). The compact operators are weak-* dense precisely when the lattice is completely distributive [16]. When the masa is atomic, one can find an approximate identity for \mathfrak{K} consisting of finite rank projections in the masa. So in particular, the compact operators in atomic CSL algebras are weak-* dense. See [2] for more details. So we obtain:

Corollary 4.3. *Every completely distributive CSL algebra has a unique predual.*

The algebra $L^\infty(0, 1)$ is not completely distributive. But it still has a unique predual by Grothendieck's Theorem. We do not know whether every CSL algebra has a unique predual.

REFERENCES

- [1] T. Ando, *On the predual of H^∞* , Comment. Math. Special Issue **1** (1978), 33–40.
- [2] K.R. Davidson, *Nest Algebras*, Pitman Research Notes in Mathematics Series **191**, Longman Scientific and Technical Pub. Co., London, New York, 1988.
- [3] K.R. Davidson, *Free semigroup algebras, a survey*, Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000), 209–240, Oper. Theory Adv. Appl., **129**, Birkhauser, Basel, 2001.
- [4] K.R. Davidson, E. Katsoulis and D.R. Pitts, *The structure of free semigroup algebras*, J. reine angew. Math. (Crelle) **533** (2001), 99–125.
- [5] K.R. Davidson and D.R. Pitts, *Invariant subspaces and hyper-reflexivity for free semi-group algebras*, Proc. London Math. Soc. **78** (1999), 401–430.
- [6] K.R. Davidson and D.R. Pitts, *The algebraic structure of non-commutative analytic Toeplitz algebras*, Math. Ann. **311** (1998), 275–303.
- [7] K. Davidson and S. Power, *Best approximation in C^* -algebras*, J. Reine Angew. Math. **368** (1986), 43–62.
- [8] E.G. Effros, N. Ozawa, and Z.J. Ruan, *On injectivity for non-commuting operators*, Duke Math. J. **110** (2001), 489–522.
- [9] J.A. Erdos, *Operators of finite rank in nest algebras*, J. London Math. Soc. **43** (1968), 391–397.
- [10] G. Godefroy, *Existence and uniqueness of isometric preduals: a survey*, Banach space theory (Iowa City, IA, 1987), 131–193, Contemp. Math. **85**, Amer. Math. Soc., Providence, RI, 1989.
- [11] G. Godefroy and Li, *Banach spaces which are M -ideals in their bidual have property (u)*, Ann. Inst. Fourier (Grenoble) **39** (1989), 361–371.
- [12] G. Godefroy and P. Saphar, *Duality in spaces of operators and smooth norms on Banach spaces*, Illinois J. Math. **32** (1988), 672–695.
- [13] G. Godefroy and M. Talagrand, *Nouvelles classes d'espaces de Banach à predual unique*, Séminaire d'analyse fonctionnelle de l'Ecole Polytechnique (1980–81).
- [14] A. Grothendieck, *Une caracterisation vectorielle metrique des espaces L^1* , Canad. J. Math. **7** (1955), 552–561.

- [15] P. Harmand, D. Werner and W. Werner, *M-ideals in Banach spaces and Banach algebras*, Lecture Notes in Math. **1547**, Springer-Verlag, New York, 1993.
- [16] C. Laurie and W. Longstaff, *A note on rank one operators in reflexive algebras*, Proc. Amer. Math. Soc. **89** (1983), 293–297.
- [17] H. Pfitzner, *Separable L -embedded Banach spaces are unique preduals*, Bull. London Math. Soc. **39** (2007), 1039–1044.
- [18] G. Popescu, *Multi-analytic operators and some factorization theorems*, Indiana Univ. Math. J. **38** (1989), 693–710.
- [19] G. Popescu, *Multi-analytic operators on Fock spaces*, Math. Ann. **303** (1995), 31–46.
- [20] Z.J. Ruan, *On the predual of dual algebras*, J. Operator Theory **27** (1992), 179–192.
- [21] S. Sakai, *A characterization of W^* -algebras*, Pacific J. Math. **6** (1956), 763–773.

PURE MATH. DEPT., U. WATERLOO, WATERLOO, ON N2L-3G1, CANADA
E-mail address: krdavids@uwaterloo.ca

PURE MATH. DEPT., U. WATERLOO, WATERLOO, ON N2L-3G1, CANADA
E-mail address: alexmwright@gmail.com