

# Universal Algebra and Computational Complexity

## Lecture 3

Ross Willard

University of Waterloo, Canada

Třešť, September 2008

# Summary of Lecture 2

Recall from Tuesday:

$L$	$\subseteq$	$NL$	$\subseteq$	$P$	$\subseteq$	$NP$	$\subseteq$	$PSPACE$	$\subseteq$	$EXPTIME \dots$
$\Psi$		$\Psi$		$\Psi$		$\Psi$		$\Psi$		$\Psi$
$FVAL,$ $2COL$		$PATH,$ $2SAT$		$CVAL,$ $HORN-$ $3SAT$		$SAT,$ $3SAT,$ $3COL,$ $4COL,$ etc. $HAMPATH$		$1-CLO$		$CLO$

Today:

- Some decision problems involving finite algebras
- How hard are they?

# Encoding finite algebras: size matters

Let  $\mathbf{A}$  be a finite algebra (always in a finite signature).

How do we **encode**  $\mathbf{A}$  for computations? And what is its *size*?

Assume  $A = \{0, 1, \dots, n-1\}$ .

For each fundamental operation  $f$ : If  $\text{arity}(f) = r$ , then  $f$  is given by its *table*, having ...

- $n^r$  entries;
- each entry requires  $\log n$  bits.

The tables (as bit-streams) must be separated from each other by  $\#$ 's.

Hence the **size** of  $\mathbf{A}$  is

$$\|\mathbf{A}\| = \sum_{\text{fund } f} \left( n^{\text{arity}(f)} \log n + 1 \right).$$

# Size of an algebra

$$||\mathbf{A}|| = \sum_{fund\ f} \left( n^{\text{arity}(f)} \log n + 1 \right).$$

Define some parameters:

$R$  = maximum arity of the fundamental operations (assume  $> 0$ )

$T$  = number of fundamental operations (assume  $> 0$ ).

Then

$$n^R \log n \leq ||\mathbf{A}|| \leq T \cdot n^R \log n + T.$$

In particular, if we restrict our attention to algebras with some **fixed** number  $T$  of operations, then

$$||\mathbf{A}|| \sim n^R \log n.$$

# Some decision problems involving algebras

INPUT: a finite algebra  $\mathbf{A}$ .

- ① Is  $\mathbf{A}$  simple? Subdirectly irreducible? Directly indecomposable?
- ② Is  $\mathbf{A}$  primal? Quasi-primal? Maltsev?
- ③ Is  $\mathbf{V}(\mathbf{A})$  congruence distributive? Congruence modular?

INPUT: two finite algebras  $\mathbf{A}, \mathbf{B}$ .

- ④ Is  $\mathbf{A} \cong \mathbf{B}$ ?
- ⑤ Is  $\mathbf{A} \in \mathbf{V}(\mathbf{B})$ ?

INPUT: A finite algebra  $\mathbf{A}$  and two terms  $s(\vec{x}), t(\vec{x})$ .

- ⑥ Does  $s = t$  have a solution in  $\mathbf{A}$ ?
- ⑦ Is  $s \approx t$  an identity of  $\mathbf{A}$ ?

INPUT: an operation  $f$  on a finite set.

- ⑧ Does  $f$  generate a minimal clone?

How hard are these problems?

# Categories of answers

Suppose  $D$  is some decision problem involving finite algebras.

- ① Is there an “obvious” algorithm for  $D$ ? What is its complexity?
  - If an obvious algorithm obviously has complexity  $Y$ , then we call  $Y$  an **obvious upper bound** for the complexity of  $D$ .
- ② Do we know a clever (nonobvious) algorithm? Does it give a lesser complexity (relative to the spectrum  $L < NL < P < NP$  etc.)?
  - If so, call this a **nonobvious upper bound**.
- ③ Can we find a clever reduction of some  $X$ -complete problem to  $D$ ?
  - If so, this gives  $X$  as a **lower bound** to the complexity of  $D$ .

Ideally, we want to find an  $X \in \{L, NL, P, NP, \dots\}$  which is both an upper and a lower bound to the complexity of  $D$  ...

- ... i.e., such that  $D$  is  $X$ -complete.

# An easy problem: Subalgebra Membership (*SUB-MEM*)

## Subalgebra Membership Problem (*SUB-MEM*)

INPUT:

- An algebra  $\mathbf{A}$ .
- A set  $S \subseteq A$ .
- An element  $b \in A$ .

QUESTION: Is  $b \in \text{Sg}^{\mathbf{A}}(S)$ ?

How hard is *SUB-MEM*?

# An obvious upper bound for *SUB-MEM*

Algorithm:

INPUT:  $\mathbf{A}, S, b$ .

$S_0 := S$

For  $i = 1, \dots, n$  ( $:= |\mathbf{A}|$ )

$S_i := S_{i-1}$

For each operation  $f$  (of arity  $r$ )

For each  $(a_1, \dots, a_r) \in (S_{i-1})^r$

$c := f(a_1, \dots, a_r)$

$S_i := S_i \cup \{c\}$ .

Next  $i$ .

OUTPUT: whether  $b \in S_n$ .

$n$  loops

$T$  operations

$\leq n^r$  instances

Heuristics:

$$n \left( \sum_f n^{\text{ar}(f)} \right) \leq n \|\mathbf{A}\| \text{ steps}$$



# The Complexity of *SUB-MEM*

So  $SUB-MEM \in TIME(N^2)$ , or maybe  $TIME(N^{4+\epsilon})$ , or surely in  $TIME(N^{55})$ , and so we get the “obvious” upper bound:

$$SUB-MEM \in P.$$

Next questions:

- Can we obtain  $P$  as a *lower* bound for  $SUB-MEM$ ?
- What was that  $P$ -complete problem again?... ( $CVAL$  or  $HORN-3SAT$ )
- Can we show  $HORN-3SAT \leq_L SUB-MEM$ ?

Theorem (N. Jones & W. Laaser, '77)

Yes.

*In other words,  $SUB-MEM$  is  $P$ -complete.*

# A variation: 1-SUB-MEM

## 1-SUB-MEM

This is the restriction of *SUB-MEM* to **unary** algebras (all fundamental operations are unary). I.e.,

INPUT: A *unary* algebra  $\mathbf{A}$ , a set  $S \subseteq A$ , and  $b \in A$ .

QUESTION: Is  $b \in \text{Sg}^{\mathbf{A}}(S)$ ?

Here is a nondeterministic log-space algorithm showing  $1\text{-SUB-MEM} \in \text{NL}$ :

NALGORITHM: guess a sequence  $c_0, c_1, \dots, c_k$  such that

- $c_0 \in S$
- For each  $i < k$ ,  $c_{i+1} = f_j(c_i)$  for some fundamental operation  $f_j$
- $c_k = b$ .

Theorem (N. Jones, Y. Lien & W. Laaser, '76)

*1-SUB-MEM is NL-complete.*

# Some tractable problems about algebras

Using *SUB-MEM*, we can deduce that many more problems are tractable (in *P*).

- 1 Given  $\mathbf{A}$  and  $S \cup \{(a, b)\} \subseteq A^2$ , determine whether  $(a, b) \in \text{Cg}^{\mathbf{A}}(S)$ .
  - Easy exercise: show this problem is  $\leq_P \text{SUB-MEM}$ .
  - (Bonus: prove that it is in *NL*.)

- 2 Given  $\mathbf{A}$  and  $S \subseteq A$ , determine whether  $S$  is a subalgebra of  $\mathbf{A}$ .

$$S \in \text{Sub}(\mathbf{A}) \Leftrightarrow \forall a \in A (a \in \text{Sg}^{\mathbf{A}}(S) \rightarrow a \in S).$$

- 3 Given  $\mathbf{A}$  and  $\theta \in \text{Eqv}(A)$ , determine whether  $\theta$  is a congruence of  $\mathbf{A}$ .
- 4 Given  $\mathbf{A}$  and  $h : A \rightarrow A$ , determine whether  $h$  is an endomorphism.
- 5 Given  $\mathbf{A}$ , determine whether  $\mathbf{A}$  is simple.

$$\mathbf{A} \text{ simple} \Leftrightarrow \forall a, b, c, d [c \neq d \rightarrow (a, b) \in \text{Cg}^{\mathbf{A}}(c, d)].$$

- 6 Given  $\mathbf{A}$ , determine whether  $\mathbf{A}$  is abelian.

$$\mathbf{A} \text{ abelian} \Leftrightarrow \forall a, c, d [c \neq d \rightarrow ((a, a), (c, d)) \notin \text{Cg}^{\mathbf{A}^2}(0_A)].$$

## Clone Membership Problem (*CLO*)

INPUT: An algebra  $\mathbf{A}$  and an operation  $g : A^k \rightarrow A$ .

QUESTION: Is  $g \in \text{Clo } \mathbf{A}$ ?

Obvious algorithm: Determine whether  $g \in \text{Sg}^{\mathbf{A}^{(A^k)}}(pr_1^k, \dots, pr_k^k)$ .

The running time is bounded by a polynomial in  $\|\mathbf{A}^{(A^k)}\|$ .

Can show

$$\log \|\mathbf{A}^{(A^k)}\| \leq n^k \|\mathbf{A}\| \leq (\|g\| + \|\mathbf{A}\|)^2.$$

Hence the running time is bounded by the exponential of a polynomial in the size of the input  $(\mathbf{A}, g)$ . I.e.,  $CLO \in EXPTIME$ .

By reducing a known *EXPTIME*-complete problem to *CLO*, Friedman and Bergman *et al* showed:

### Theorem

*CLO* is *EXPTIME*-complete.

## The Primal Algebra Problem (*PRIMAL*)

INPUT: a finite algebra  $\mathbf{A}$ .

QUESTION: Is  $\mathbf{A}$  primal?

The obvious algorithm is actually a reduction to *CLO*.

For a finite set  $A$ , let  $g_A$  be your favorite binary Sheffer operation on  $A$ .

Define  $f : \text{PRIMAL}_{\text{inp}} \rightarrow \text{CLO}_{\text{inp}}$  by

$$f : \mathbf{A} \mapsto (\mathbf{A}, g_A).$$

Since

$$\mathbf{A} \text{ is primal} \iff g_A \in \text{Clo } \mathbf{A},$$

we have  $\text{PRIMAL} \leq_f \text{CLO}$ . Clearly  $f$  is  $P$ -computable, so

$$\text{PRIMAL} \leq_P \text{CLO}$$

which gives the obvious upper bound

$$\text{PRIMAL} \in \text{EXPTIME}.$$

But testing primality of algebras is special. Maybe there is a better, “nonobvious” algorithm?

(E.g., using Rosenberg's classification?)

## Open Problem 1.

Determine the complexity of *PRIMAL*.

- Is it in *PSPACE*? ( = *NPSPACE*)
- Is it *EXPTIME*-complete? (  $\Leftrightarrow CLO \leq_P PRIMAL$ )

## MALTSEV

INPUT: a finite algebra  $\mathbf{A}$ .

QUESTION: Does  $\mathbf{A}$  have a Maltsev term?

The obvious upper bound is *NEXPTIME*, since *MALTSEV* is a projection of

$$\{ (\mathbf{A}, p) : \underbrace{p \in \text{Clo } \mathbf{A}}_{EXPTIME} \text{ and } \underbrace{p \text{ is a Maltsev operation}}_P \},$$

a problem in *EXPTIME*.

But a slightly less obvious algorithm puts *MALTSEV* in *EXPTIME*. Use the fact that if  $x, y$  name the two projections  $A^2 \rightarrow A$ , then  $\mathbf{A}$  has a Maltsev term iff

$$(y, x) \in \text{Sg}^{(\mathbf{A}^{(A^2)})^2}((x, x), (x, y), (y, y))$$

(which is decidable in *EXPTIME*).

Similar characterizations give *EXPTIME* as an upper bound to the following:

### Some problems in *EXPTIME*

Given **A**:

- 1 Does **A** have a majority term?
- 2 Does **A** have a semilattice term?
- 3 Does **A** have Jónsson terms?
- 4 Does **A** have Gumm terms?
- 5 Does **A** have terms equivalent to  $\mathbf{V}(\mathbf{A})$  being congruence meet-semidistributive?
- 6 Etc. etc.

Are these problems easier than *EXPTIME*, or *EXPTIME*-complete?



# Freese & Valeriote's theorem

For some of these problems we have an answer:

## Theorem (R. Freese, M. Valeriote, '0?)

*The following problems are all EXPTIME-complete:*

*Given  $\mathbf{A}$ ,*

- ① *Does  $\mathbf{A}$  have Jónsson terms?*
- ② *Does  $\mathbf{A}$  have Gumm terms?*
- ③ *Is  $\mathbf{V}(\mathbf{A})$  congruence meet-semidistributive?*
- ④ *Does  $\mathbf{A}$  have a semilattice term?*
- ⑤ *Does  $\mathbf{A}$  have any nontrivial idempotent term?*
  - *idempotent* means “satisfies  $f(x, x, \dots, x) \approx x$ .”
  - *nontrivial* means “other than  $x$ .”

# Freese & Valeriote's theorem

## Proof.

Freese and Valeriote give a construction which, given an input  $\Gamma = (\mathbf{A}, g)$  to  $CLO$ , produces an algebra  $\mathbf{B}_\Gamma$  such that:

- $g \in \text{Clo } \mathbf{A} \Rightarrow$  there is a flat semilattice order on  $B_\Gamma$  such that  $(x \wedge y) \vee (x \wedge z)$  is a term operation of  $\mathbf{B}_\Gamma$ .
- $g \notin \text{Clo } \mathbf{A} \Rightarrow \mathbf{B}_\Gamma$  has no nontrivial idempotent term operations.

Moreover, the function  $f : \Gamma \mapsto \mathbf{B}_\Gamma$  is easily computed (in  $\mathbf{P}$ ).

Hence  $f$  is simultaneously a  $P$ -reduction of  $CLO$  to all the problems in the statement of the theorem. □

## Open Problem 2.

Are the following easier than  $EXPTIME$ , or  $EXPTIME$ -complete?

- Determining if  $\mathbf{A}$  has a majority operation.
- Determining if  $\mathbf{A}$  has a Maltsev operation ( $MALTSEV$ ).

If  $MALTSEV$  is easier than  $EXPTIME$ , then so is  $PRIMAL$ , since

## Theorem

$\mathbf{A}$  is primal iff:

- $\mathbf{A}$  has no proper subalgebras,
  - $\mathbf{A}$  is simple,
  - $\mathbf{A}$  is rigid,
  - $\mathbf{A}$  is not abelian, and
  - $\mathbf{A}$  is Maltsev.
- } in  $P$

Surprisingly, the previous problems become significantly easier when restricted to *idempotent* algebras.

### Theorem (Freese & Valeriote, '0?)

*The following problems for **idempotent** algebras are in P:*

- ① **A** has a majority term.
- ② **A** has Jónsson terms.
- ③ **A** has Gumm terms.
- ④  $V(\mathbf{A})$  is congruence meet-semidistributive.
- ⑤ **A** is Maltsev.
- ⑥  $V(\mathbf{A})$  is congruence  $k$ -permutable for some  $k$ .

### Proof.

Fiendishly nonobvious algorithms using tame congruence theory. □

## Variety Membership Problem (VAR-MEM)

INPUT: two finite algebras  $\mathbf{A}, \mathbf{B}$  in the same signature.

QUESTION: Is  $\mathbf{A} \in \mathbf{V}(\mathbf{B})$ ?

The obvious algorithm (J. Kalicki, '52): determine whether the identity map on  $A$  extends to a homomorphism  $\mathbf{F}_{\mathbf{V}(\mathbf{B})}(A) \rightarrow \mathbf{A}$ .

Theorem (C. Bergman & G. Slutzki, '00)

*The obvious algorithm puts VAR-MEM in 2-EXPTIME.*

$$2\text{-EXPTIME} \stackrel{\text{def}}{=} \bigcup_{k=1}^{\infty} \text{TIME}(2^{(2^{O(N^k)})})$$

$$\dots NEXPTIME \subseteq EXPSPACE \subseteq 2\text{-EXPTIME} \subseteq N(2\text{-EXPTIME}) \dots$$

What is the “real” complexity of *VAR-MEM*?

Theorem (Z. Székely, thesis '00)

*VAR-MEM* is NP-hard (i.e.,  $3SAT \leq_P VAR-MEM$ ).

Theorem (M. Kozik, thesis '04)

*VAR-MEM* is EXPSPACE-hard.

Theorem (M. Kozik, '0?)

*VAR-MEM* is 2-EXPTIME-hard and therefore 2-EXPTIME-complete.  
Moreover, there exists a specific finite algebra **B** such that the subproblem:

*INPUT: a finite algebra A in the same signature as B.*

*QUESTION: Is  $A \in V(B)$*

*is 2-EXPTIME-complete.*

## The Equivalence of Terms problem (*EQUIV-TERM*)

INPUT:

- A finite algebra  $\mathbf{A}$ .
- Two terms  $s(\vec{x}), t(\vec{x})$  in the signature of  $\mathbf{A}$ .

QUESTION: Is  $s(\vec{x}) \approx t(\vec{x})$  identically true in  $\mathbf{A}$ ?

It is convenient to name the *negation* of this problem:

## The Inequivalence of Terms problem (*INEQUIV-TERM*)

INPUT: (same)

QUESTION: Does  $s(\vec{x}) \neq t(\vec{x})$  have a solution in  $\mathbf{A}$ ?

How hard are these problems?

Obviously *INEQUIV-TERM* is in *NP*. (Any solution  $\vec{x}$  to  $s(\vec{x}) \neq t(\vec{x})$  serves as a certificate.)

On the other hand, and equally obviously,  $SAT \leq_P \text{INEQUIV-TERM}$ . (Map  $\varphi \mapsto (2_{BA}, \varphi, 0)$ .)

Hence *INEQUIV-TERM* is obviously *NP*-complete.

*EQUIV-TERM*, being its negation, is said to be **co-*NP***-complete.

## Definition

- **Co-*NP*** is the class of problems  $D$  whose negation  $\neg D$  is in *NP*.
- A problem  $D$  is **co-*NP*-complete** if its negation  $\neg D$  is *NP*-complete, or equivalently, if  $D$  is in the top  $\equiv_P$ -class of co-*NP*.

Done. End of story. Boring.



But WAIT!!!! There's more!!!!

For each fixed finite algebra  $\mathbf{A}$  we can pose the subproblem for  $\mathbf{A}$ :

### *EQUIV-TERM*( $\mathbf{A}$ )

INPUT: two terms  $s(\vec{x}), t(\vec{x})$  in the signature of  $\mathbf{A}$ .

QUESTION: (same).

The following are obviously obvious:

- *EQUIV-TERM*( $\mathbf{A}$ ) is in *co-NP* for any algebra  $\mathbf{A}$ .
- *EQUIV-TERM*( $\mathbf{2}_{BA}$ ) is *co-NP*-complete. (Hint:  $\varphi \mapsto (\varphi, 0)$ .)
- *EQUIV-TERM*( $\mathbf{A}$ ) is in *P* when  $\mathbf{A}$  is nice, say, a vector space or a set.

Problem: for which finite algebras  $\mathbf{A}$  is *EQUIV-TERM*( $\mathbf{A}$ ) *NP*-complete?

For which  $\mathbf{A}$  is it in *P*?

There are a huge number of publications in this area. Here is a sample:

Theorem (H. Hunt & R. Stearns, '90; S. Burris & J. Lawrence, '93)

Let  $R$  be a finite ring.

- If  $R$  is nilpotent, then  $EQUIV-TERM(R)$  is in  $P$ .
- Otherwise,  $EQUIV-TERM(R)$  is co-NP-complete.

Theorem (Burris & Lawrence, '04; G. Horváth & C. Szabó, '06; Horváth, Lawrence, L. Mériai & Szabó, '07)

Let  $G$  be a finite group.

- If  $G$  is nonsolvable, then  $EQUIV-TERM(G)$  is co-NP-complete.
- If  $G$  is nilpotent, or of the form  $Z_{m_1} \rtimes (Z_{m_2} \rtimes \cdots (Z_{m_k} \rtimes A) \cdots)$  with each  $m_i$  square-free and  $A$  abelian, then  $EQUIV-TERM(G)$  is in  $P$ .

And many partial results for **semigroups** due to e.g. Kisielewicz, Klíma, Pleshcheva, Popov, Seif, Szabó, Tesson, Therien, Vértesi, and Volkov.

# An outrageous scandal

## Theorem (G. Horváth & C. Szabó)

Consider the group  $\mathbf{A}_4$ .

- $EQUIV-TERM(\mathbf{A}_4)$  is in  $P$ .
- Yet there is an algebra  $\mathbf{A}$  with the same clone as  $\mathbf{A}_4$  such that  $EQUIV-TERM(\mathbf{A})$  is *co-NP-complete*.

This is either wonderful or scandalous.


In my opinion, this is evidence that  $EQUIV-TERM$  is the wrong problem.

## Definition

A **circuit** (in a given signature for algebras) is an object, similar to a term, except that repeated subterms need be written only once.

Example: Let  $t = ((x + y) + (x + y)) + ((x + y) + (x + y))$ .

A circuit for  $t$ :



Straight-line program:

$$v_1 = x + y$$

$$v_2 = v_1 + v_1$$

$$t = v_2 + v_2.$$

Note that circuits may be significantly shorter than the terms they represent.

# Equivalence of Terms Problem (correct version)

Fix a finite algebra  $\mathbf{A}$ .

The Equivalence of Circuits problem ( $EQUIV-CIRC(\mathbf{A})$ )

INPUT: two **circuits**  $s(\vec{x})$ ,  $t(\vec{x})$  in the signature of  $\mathbf{A}$ .

QUESTION: is  $s(\vec{x}) \approx t(\vec{x})$  identically true in  $\mathbf{A}$ ?

This is the correct problem.

- The input is presented “honestly” (computationally).
- It is invariant for algebras with the same clone.

## Open Problem 3.

For which finite algebras  $\mathbf{A}$  is  $EQUIV-CIRC(\mathbf{A})$  NP-complete? For which  $\mathbf{A}$  is it in  $P$ ?

# Two problems for relational structures

## Relational Clone Membership (*RCLO*)

INPUT:

- A finite relational structure  $\mathbf{M}$ .
- A finitary relation  $R \subseteq M^k$ .

QUESTION: Is  $R \in \text{Inv Pol}(\mathbf{M})$ ?

A slightly nonobvious characterization gives *NEXPTIME* as an upper bound. For a lower bound, we have:

## Theorem (W, '0?)

*RCLO* is *EXPTIME*-hard.

## Open Problem 4.

Is *RCLO* in *EXPTIME*? Is it *NEXPTIME*-complete?

Fix a finite relational structure  $\mathbf{B}$ .

Consider the following problem associated to  $\mathbf{B}$ :

### A problem

INPUT: a finite structure  $\mathbf{A}$  in the same signature as  $\mathbf{B}$ .

QUESTION: Is there a homomorphism  $h : \mathbf{A} \rightarrow \mathbf{B}$ ?

This problem is called  $CSP(\mathbf{B})$ .

Obviously  $CSP(\mathbf{B}) \in NP$  for any  $\mathbf{B}$ .

If  $\mathbf{K}_3$  is the triangle graph, then  $CSP(\mathbf{K}_3) = 3COL$ , so is  $NP$ -complete in this case. If  $\mathbf{G}$  is a bipartite graph, then  $CSP(\mathbf{G}) \in P$ .

### $CSP$ Classification Problem

For which finite relational structures  $\mathbf{B}$  is  $CSP(\mathbf{B})$  in  $P$ ? For which is it  $NP$ -complete?