

MODEL THEORY OF COMPACT COMPLEX MANIFOLDS WITH AN AUTOMORPHISM

MARTIN BAYS, MARTIN HILS, AND RAHIM MOOSA

ABSTRACT. Motivated by possible applications to meromorphic dynamics, and generalising known properties of difference-closed fields, this paper studies the theory CCMA of compact complex manifolds with a generic automorphism. It is shown that while CCMA does admit geometric elimination of imaginaries, it cannot eliminate imaginaries outright: a counterexample to 3-uniqueness in CCM is exhibited. Finite-dimensional types are investigated and it is shown, following the approach of Pillay and Ziegler, that the canonical base property holds in CCMA. As a consequence the Zilber dichotomy is deduced: finite-dimensional types of SU -rank one are either one-based or almost internal to the fixed field. In addition, a general criterion for stable embeddedness in TA (when it exists) is established, and used to determine the full induced structure of CCMA on projective varieties, simple nonalgebraic complex tori, and simply connected nonalgebraic strongly minimal manifolds.

CONTENTS

1. Introduction	1
2. CCM fails 3-uniqueness	4
3. Internalising finite covers in CCM	7
4. Generic automorphisms: imaginaries and stable embeddedness	10
5. Finite-dimensional types in CCMA	14
6. Difference-analytic jet spaces and the Zilber dichotomy	16
7. Minimal one-based types	19
Appendix A. Complex analytic jet spaces	21
References	23

1. INTRODUCTION

Underlying the applications of model theory to algebraic dynamics [8, 19] is the equivalence of the category of rational dynamical systems with that of finitely generated difference fields: To a rational dynamical system (V, f) over \mathbb{C} , where V is a projective algebraic variety and $f : V \rightarrow V$ is a dominant rational map, one associates the difference field $(\mathbb{C}(V), f^*)$ where $\mathbb{C}(V)$ is the rational function field

Date: February 24, 2015.

2010 *Mathematics Subject Classification.* Primary: 03C60; Secondary: 03C45, 03C65, 32J99.

Key words and phrases. Model Theory, Compact Complex Manifold, Generic Automorphism, Zilber Dichotomy, Canonical Base Property.

M. Hils was partially funded by the Agence Nationale de Recherche [ValCoMo, Projet ANR blanc ANR-13-BS01-0006].

R. Moosa was partially supported by an NSERC Discovery Grant.

and $f^* : \mathbb{C}(V) \rightarrow \mathbb{C}(V)$ is the endomorphism induced by precomposition with f . The theory of difference fields is tractable from a model-theoretic point of view because it admits a model companion, the theory ACFA of difference-closed fields, which is tame: it admits a certain quantifier reduction, eliminates imaginaries, is supersimple, and realises a Zilber dichotomy for types of SU -rank one.

What happens when (V, f) is a *meromorphic dynamical system*, that is, when V is a possibly nonalgebraic compact complex manifold and f is a dominant meromorphic map? The first challenge is to find a replacement for the rational function field. The meromorphic function field will not do as it only captures the algebraic part of V . For example, $\mathbb{C}(V) = \mathbb{C}$ when V is a simple nonalgebraic complex torus. An appropriate generalisation of the rational function field is suggested by model theory. Let \mathcal{A} denote the structure whose sorts are (reduced and irreducible) compact complex analytic spaces and where the basic relations are all complex analytic subsets. The first-order theory of \mathcal{A} , in this language L , is denoted by CCM. It is a stable theory that admits quantifier elimination, eliminates imaginaries, and whose sorts are of finite Morley rank — see [22] for a survey of this theory. Let \mathcal{A}' be an \aleph_0 -saturated elementary extension of \mathcal{A} . Then V is a sort and by saturation there exists $c \in V(\mathcal{A}')$ that is *generic* in that it is not contained in $X(\mathcal{A}')$ for any proper complex analytic subset $X \subset V$. What model theory suggests as an analogue to the rational function field is the L -substructure of \mathcal{A}' whose underlying set is the definable closure of c . Let us denote this L -structure, which is uniquely determined up to L -isomorphism, by $\mathcal{C}(V)$. Note that by quantifier elimination, $\mathcal{C}(V)$ can be identified with the set of all dominant meromorphic maps $g : V \rightarrow W$, as W ranges over all compact complex manifolds. (In contrast, the meromorphic function field corresponds to considering only $W = \mathbb{P}(\mathbb{C})$, the projective line.)

We can now imitate the difference-field construction. If (V, f) is a meromorphic dynamical system then f induces a map $f^* : \mathcal{C}(V) \rightarrow \mathcal{C}(V)$ by precomposition: an element of $\mathcal{C}(V)$ is of the form $g(c)$ for some dominant meromorphic map $g : V \rightarrow W$, and $f^*(g(c)) := g(f(c))$. Note that as f is dominant, $f(c)$ is again generic in $V(\mathcal{A}')$, and hence g is defined at $f(c)$. The dominance of f also ensures that f^* is injective. It is not hard to check that f^* is an L -embedding and, exactly extending the algebraic case, the association $(V, f) \mapsto (\mathcal{C}(V), f^*)$ is an equivalence of categories.

This motivates us to study substructures of models of CCM equipped with L -embeddings. That is, letting $L_\sigma := L \cup \{\sigma\}$, we are interested in the universal L_σ -theory $\text{CCM}_{V, \sigma} := \text{CCM}_V \cup \{\sigma\}$ is an L -embedding}. As was noted in [3], $\text{CCM}_{V, \sigma}$ has a model companion, denoted by CCMA. Various tameness properties for CCMA, in particular supersimplicity, follow automatically from the work of Chatzidakis and Pillay [9] on generic automorphisms. In this paper we begin a finer study of CCMA.

It may be worth emphasising that actual compact complex manifolds have no nontrivial L -automorphisms since every point is named in the language. It is the consideration of L -automorphisms of *nonstandard models* of compact complex manifolds that gives rise to CCMA.

Our first observation about CCMA is negative: while it admits geometric elimination of imaginaries,

CCMA *does not eliminate imaginaries*. (Theorem 4.2)

To prove this we use a characterisation of Hrushovski's which reduces the problem to proving that CCM does not satisfy the property of *3-uniqueness* introduced in [17]. The failure of 3-uniqueness in CCM, which is established in Section 2 below, is of independent interest because it distinguishes CCM from the stable theories that arise in the model theory of fields (ACF, DCF, SCF).

Since the complex field is definable in CCM, on the projective line, in CCMA we have a definable difference-closed field extending $(\mathbb{C}, +, \times, \text{id})$. One of the things we point out here is that in this way

ACFA is purely stably embedded in CCMA. (Theorem 4.8)¹

Contrast this with the situation for differentially closed fields with a generic automorphism, where ACFA appears as a proper reduct of the full structure induced on the field of constants. In a similar vein, we prove that

the full structure induced by CCMA on any simple nonalgebraic complex torus, or on any simply connected nonalgebraic strongly minimal compact complex manifold, is just the complex analytic structure together with σ . (Theorem 4.8)

In fact all of these results follow from a general characterisation (Proposition 4.5) for stable embeddability in TA , whenever the latter exists. The characterisation is that the sort in question should *internalise finite covers in T* , that is, relative to that sort in T almost internality should imply outright internality. That the projective line internalises finite covers in CCM is a (known) uniform version of the fact that a finite cover of a Moishezon space is again Moishezon. For simple nonalgebraic complex tori and for simply connected nonalgebraic strongly minimal compact complex manifolds, the condition is proved in Section 3.

But our primary interest is in the structure of the *finite-dimensional* types, that is, the types of tuples c such that the complex dimension of the locus of $(c, \sigma(c), \dots, \sigma^n(c))$ is uniformly bounded as n grows. For example, the types arising from meromorphic dynamical systems as described above are finite-dimensional. The converse is almost true too: every finite-dimensional type in CCMA comes from a pair (V, Γ) where V is a compact complex manifold and Γ is a finite-to-finite meromorphic self-correspondence on V . See Section 5 for details. Our main theorem about finite-dimensional types is that they enjoy the canonical base property:

the canonical base of a finite-dimensional type in CCMA is almost internal, over a realisation of the type, to the fixed field. (Theorem 6.4)

Our proof follows the approach of Pillay and Ziegler [30] for ACFA; we define an appropriate notion of *jet space* in the context of CCMA. As a consequence, we obtain that

a finite-dimensional type of SU-rank one in CCMA is either one based or almost internal to the fixed field. (Corollary 6.5)

We expect, but do not prove, that the finite-dimensionality assumption in the above Zilber dichotomy statement can be removed. We point out in Example 6.7 that infinite-dimensional types of SU-rank one exist.

In a short final section we explain how the methods of Hrushovski [16] and Chatzidakis-Hrushovski [7] extend to our setting to yield a characterisation of the finite-dimensional types of *SU-rank one* that are nontrivial and one-based.

¹Here, and throughout, by ACFA we really mean the complete theory of models of ACFA extending the trivial difference field $(\mathbb{C}, +, \times, \text{id})$, i.e., with the elements of \mathbb{C} named by constants.

1.1. Review of Conventions and Notations for CCM. By a *complex variety* we will mean a reduced and irreducible complex analytic space. We say that a property holds *generally* if it holds outside a proper complex analytic subset. Recall that \mathcal{A} is the structure where the sorts are the compact complex varieties and the basic relations are the complex analytic subsets of the cartesian products of the sorts. The first-order theory of \mathcal{A} is denoted by CCM. One of the sorts of \mathcal{A} is the projective line $\mathbb{P}(\mathbb{C})$. The complex field $(\mathbb{C}, +, \times)$ is definable in this sort, and in fact the full induced structure on $\mathbb{P}(\mathbb{C})$, which by Chow's Theorem is just that of the algebraic sets over \mathbb{C} , is bi-interpretable with the complex field.

We will usually work in a sufficiently saturated elementary extension \mathcal{A}' of \mathcal{A} . We will denote by $(K, +, \times)$ the corresponding elementary extension of the complex field, the interpretation in \mathcal{A}' of the complex field. It is up to definable isomorphism the only infinite field definable in \mathcal{A}' , see [21, Corollary 4.8].

The complex analytic sets induce a noetherian topology on the sorts of \mathcal{A}' too: if X is a compact complex variety then by a *closed* subset of $X(\mathcal{A}')$ is meant a set of the form $F := Y(\mathcal{A}')_s$ where Y is a complex analytic subset of $S \times X$ for some compact complex variety S , and $s \in S(\mathcal{A}')$. If s comes from a given set of parameters B , then we also say that F is *B-closed*. A *B-closed* set is *B-irreducible* if it cannot be written as the union of two proper *B-closed* sets. A *generic point of F over B* is an element $c \in F$ that is not contained in any proper *B-closed* subset, in this case we also say that F is the *locus of c over B*, denoted by $F = \text{loc}(c/B)$. We say F is *irreducible* if it is absolutely irreducible in the sense that it cannot be written as a union of two proper closed subsets (over any parameters). Even though the closed subsets of $X(\mathcal{A}')$ are not themselves complex analytic spaces, it still makes sense to talk about their “complex” *dimension* since in the standard model the complex dimension of a complex analytic set is definable in parameters. For proofs of the claims made here, and for more details on this “nonstandard Zariski” topology, see [21, §2].

1.2. Acknowledgements. This collaboration began during the Spring 2014 MSRI programme *Model Theory, Arithmetic Geometry and Number Theory*. The authors would like to thank MSRI for its hospitality and stimulating research environment. We are also grateful to Frédéric Campana, Bradd Hart, and Matei Toma, for useful discussions.

2. CCM FAILS 3-UNIQUENESS

The goal of this section is to prove the following theorem about the model theory of compact complex manifolds. Its significance to the subject of this paper is that it will imply that CCMA does not eliminate imaginaries, see Theorem 4.2 and Remark 4.3 below.

Theorem 2.1. *CCM does not satisfy 3-uniqueness in the sense of [17, §4]. That is, working in a sufficiently saturated $\mathcal{A}' \models \text{CCM}$, there exist elements b, x, y, z with x, y, z independent over b and such that*

$$\text{acl}(bxy) \cap \text{dcl}(\text{acl}(bxz), \text{acl}(byz)) \neq \text{dcl}(\text{acl}(bx), \text{acl}(by)).$$

Our witness to the failure of 3-uniqueness will use some of the theory of holomorphic line bundles on complex manifolds. We suggest [12, §1.1] for more details and as a general reference for this material.

In fact, it is convenient for us to work instead with the corresponding \mathbb{C}^* -bundles. Recall that a *holomorphic \mathbb{C}^* -bundle* on a complex manifold M is a complex manifold P with a holomorphic surjective map $P \rightarrow M$ for which there exists a (euclidean) open cover $\{U_i\}$ of M such that $P|_{U_i}$ is biholomorphic to $U_i \times \mathbb{C}^*$ over U_i , and such that the corresponding biholomorphisms $U_i \cap U_j \times \mathbb{C}^* \rightarrow U_i \cap U_j \times \mathbb{C}^*$ are of the form $(x, p) \mapsto (x, g_{ij}(x)p)$ for some (holomorphic) $g_{ij} : U_i \cap U_j \rightarrow \mathbb{C}^*$. One obtains a uniform holomorphic group action of \mathbb{C}^* on the fibres of $P \rightarrow M$. Since the action by multiplication of \mathbb{C}^* on \mathbb{C}^* extends uniquely and holomorphically to \mathbb{C} , an alternative description of \mathbb{C}^* -bundles is that they are holomorphic line bundles with their zero sections removed. Moreover, as that action extends also uniquely and holomorphically to $\mathbb{P}(\mathbb{C})$, we can embed P as a Zariski open subset of a projective line bundle over M , which we will denote by $P^{\text{cl}} \rightarrow M$. Hence, if M is compact then $P \rightarrow M$, as well as the action of \mathbb{C}^* on the fibres, is definable in CCM. The set of holomorphic \mathbb{C}^* -bundles over M , up to a natural notion of isomorphism, can be identified with the cohomology group $H^1(M, \mathcal{O}_M^*)$, where \mathcal{O}_M^* is the sheaf of nonvanishing holomorphic functions on M . We will denote the element of $H^1(M, \mathcal{O}_M^*)$ corresponding to P by $[P]$.

The following fact was pointed out to us by Matei Toma.

Fact 2.2. *There exists a strongly minimal simply connected smooth compact complex surface X with $H^1(X, \mathcal{O}_X^*) \cong \mathbb{Z}$.*

Proof. The examples come from K3 surfaces, which are certain simply connected smooth compact complex surfaces, details about which can be found in [2, Chapter VII]. In particular, for any K3 surface X , $H^1(X, \mathcal{O}_X) = 0$, and so the exponential short exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X^* \rightarrow 1$ gives rise to an embedding of $H^1(X, \mathcal{O}_X^*)$ into the discrete group $H^2(X, \mathbb{Z})$. That is, the Picard and Néron-Severi groups coincide. In particular we have a cup-product $H^1(X, \mathcal{O}_X^*) \times H^1(X, \mathcal{O}_X^*) \rightarrow H^4(X, \mathbb{Z}) \cong \mathbb{Z}$.

By [32, Theorem 1], for any $g \in \mathbb{Z}$ there are K3 surfaces X with $H^1(X, \mathcal{O}_X^*) = \mathbb{Z}\beta$ for some β with $(\beta, \beta) = 2g - 2$. Let X be such a surface with $g < 0$. All that remains to be checked is that X is strongly minimal. But if C were an irreducible curve in X then it would give rise to a nontrivial \mathbb{C}^* -bundle $P \rightarrow X$, with C the divisor of the associated line bundle. Now $[P] = n\beta$ for some $n \in \mathbb{Z} \setminus \{0\}$, and so $([P], [P]) = n^2(2g - 2) \leq -4$. But $([P], [P])$ is equal to the self-intersection number of C , which by [2, VII.3.7(ii)] is at least -2 on a K3 surface. Hence no such C exists, and X is strongly minimal. \square

Fix X as given by Fact 2.2.

Lemma 2.3. *For any nonempty Zariski open subset $X' \subseteq X$, the inclusion induces a natural isomorphism $H^1(X', \mathcal{O}_{X'}^*) \cong H^1(X, \mathcal{O}_X^*)$.*

Proof. This only uses the fact that X is strongly minimal and of dimension greater than one. By strong minimality $X \setminus X'$ is a finite set, and hence of codimension 2, so that by Hartog's theorem restriction of line bundles to X' is an isomorphism. \square

Lemma 2.4. *If $b \in X(\mathcal{A}')$ is generic then $\text{acl}(b) = \text{dcl}(b)$*

Proof. This only uses the fact that X is simply connected, smooth, and has no subvarieties of codimension 1. Let $c \in \text{acl}(b)$, $Y := \text{loc}(c, b)$, and $\pi : Y \rightarrow X$ the generically finite surjective holomorphic map given by the second co-ordinate projection. Consider the Stein factorisation of π

$$\begin{array}{ccc} Y & \xrightarrow{f} & \tilde{Y} \\ & \searrow \pi & \swarrow \tilde{\pi} \\ & X & \end{array}$$

so that f is bimeromorphic and $\tilde{\pi}$ is now everywhere finite-to-one. Next, precomposing with a normalisation $\hat{Y} \rightarrow \tilde{Y}$, we obtain a (still finite) surjective map $\hat{\pi} : \hat{Y} \rightarrow X$. Because \hat{Y} is normal and X is smooth we can apply the purity of branch locus theorem (see, for example, Theorem VII.1.6 of [11]) which tells us that the ramification locus of $\hat{\pi}$, namely the set of points in \hat{Y} where $\hat{\pi}$ is not locally biholomorphic, say E , is a complex analytic set that is either empty or of codimension 1. In the latter case, as $\hat{\pi}$ is finite, it would follow that $\hat{\pi}(E)$ is a curve in X , contradicting strong minimality of this surface. Hence $\hat{\pi}$ is unramified. But simple-connectedness of X then forces $\hat{\pi}$, and hence $\tilde{\pi} : \tilde{Y} \rightarrow X$, to be an isomorphism. It follows that the original $\pi : Y \rightarrow X$ is bimeromorphic, and so $c \in \text{dcl}(b)$, as desired. \square

Fix a nontrivial holomorphic \mathbb{C}^* -bundle $P \rightarrow X$ such that $[P]$ is a generator of $H^1(X, \mathcal{O}_X^*)$, given to us by Fact 2.2. Let $\pi : P^{\text{cl}} \rightarrow X$ be the ambient projective line bundle in which P lives as a Zariski open set.

Proposition 2.5. *If $a \in P^{\text{cl}}(\mathcal{A}')$ is generic then $\text{acl}(a) = \text{dcl}(a)$.*

Proof. We are grateful to Will Sawin for his answer to a mathoverflow question [33] which pointed us toward analysing the situation in terms of étale covers.

Let $z \in \text{acl}(a)$ and set $Z := \text{loc}(z)$. Arguing exactly as in the beginning of the proof of Lemma 2.4, after replacing z with some z' such that $z' \in \text{acl}(a)$ and $a, z \in \text{dcl}(z')$, we may assume that there is a finite surjective holomorphic map $f : Z \rightarrow P^{\text{cl}}$ whose ramification locus E is either empty or of pure codimension one.

We first argue that $f(E) \subseteq P^{\text{cl}} \setminus P$. Indeed, if $E \neq \emptyset$ then as f is finite, $f(E)$ is of pure codimension one in P^{cl} , and hence is of dimension 2. Let S be an irreducible component of $f(E)$. Since the fibres of π are one-dimensional, it follows that $\pi(S)$ is infinite in X , and so by strong minimality is all of X . Thus S projects generically finitely onto X , and so by Lemma 2.4, the projection is in fact generically one-to-one. That is, S is the image of a meromorphic section to $P^{\text{cl}} \rightarrow X$. So if $S \cap P \neq \emptyset$, i.e., if S is generically contained in P , then $P \rightarrow X$ has a meromorphic section and must therefore be bimeromorphically trivial: there is a nonempty Zariski open subset $X' \subseteq X$ such that $P|_{X'}$ is isomorphic to $X' \times \mathbb{C}^*$ over X' . Hence $[P|_{X'}] = 0$ in $H^1(X', \mathcal{O}_{X'}^*)$. By 2.3, this contradicts the nontriviality of $[P]$. So $f(E) \subseteq P^{\text{cl}} \setminus P$.

Hence, setting $Q := f^{-1}(P)$, $f|_Q : Q \rightarrow P$ is an unramified finite cover. We aim to prove that, after possibly removing finitely many points from X , Q is also a \mathbb{C}^* -bundle and that in the local trivialisations $f|_Q$ becomes $\text{id}_U \times [n] : U \times \mathbb{C}^* \rightarrow U \times \mathbb{C}^*$ for some $n > 0$, where $[n] : \mathbb{C}^* \rightarrow \mathbb{C}^*$ is the raising to the power n map.

To that end, note first of all that by Lemma 2.4 applied to $b := \pi(a) \in X(\mathcal{A}')$, $\text{tp}(z/b)$ is stationary, and hence the general fibres of $Z \rightarrow X$ are irreducible (see

Lemmas 2.7 and 2.11 of [21]). It follows that the general fibres of $Q \rightarrow X$ are irreducible, so in particular connected for the Zariski topology and thus for the euclidean topology, too. Let $X' \subseteq X$ be a nonempty Zariski open set over which this happens, and let $U \subseteq X'$ be an open ball such that $P|_U$ is isomorphic to $U \times \mathbb{C}^*$ over U . Then $Q|_U$ is connected and we have an induced holomorphic unramified finite covering map $\alpha : Q|_U \rightarrow U \times \mathbb{C}^*$ over U . Since $U \times \mathbb{C}^*$ has fundamental group \mathbb{Z} , any topological covering map is isomorphic as a topological covering map to some $\text{id}_U \times [n] : U \times \mathbb{C}^* \rightarrow U \times \mathbb{C}^*$. Now α is in particular a topological covering map, so there is a topological covering map isomorphism $\theta : Q|_U \rightarrow U \times \mathbb{C}^*$ such that $\alpha = (\text{id}_U \times [n]) \circ \theta$. But α and $[n]$ are locally biholomorphic, so it follows that θ is locally biholomorphic, and so is an isomorphism of complex manifolds. That is, $Q|_U \rightarrow P|_U$ is isomorphic to $\text{id}_U \times [n] : U \times \mathbb{C}^* \rightarrow U \times \mathbb{C}^*$.

What we have shown is that the \mathbb{C}^* -bundle structure on $P|_{X'}$ lifts to one on $Q|_{X'}$ and that in $H^1(X', \mathcal{O}_{X'}^*)$, $n[Q|_{X'}] = [P|_{X'}]$. It follows by Lemma 2.3 that $[P]$ is n -divisible in $H^1(X, \mathcal{O}_X^*)$. This contradicts the fact that $[P]$ generates $H^1(X, \mathcal{O}_X^*)$, unless $n = 1$. But then $Q|_{X'} \rightarrow P|_{X'}$ is an isomorphism, and hence $f : Z \rightarrow P^{\text{cl}}$ is bimeromorphic. So $z \in \text{dcl}(a)$. \square

Proof of Theorem 2.1. We have a uniformly definable action of \mathbb{C}^* on the fibres of $P \rightarrow X$. So if $b \in X(\mathcal{A}')$ is generic, P_b is a definable (principal) homogenous space for K^* . Let $a \in P_b$ be generic over b , and $\phi, \theta \in K^*$ independent generics over a . So $(a, \phi a, \theta a) \in P_b^3$ is an independent triple over b , and this will be our witness to non-3-uniqueness. Fix $n > 1$ and choose n th roots ϕ' and θ' , of ϕ and θ respectively. Then $\frac{\phi'}{\theta'} \in \text{acl}(\frac{\phi}{\theta}) \subseteq \text{acl}(\phi a, \theta a)$. On the other hand, $\theta' \in \text{acl}(\theta) \subseteq \text{acl}(a, \theta a)$. It follows that $\phi' \in \text{dcl}(\frac{\phi'}{\theta'}, \theta') \subseteq \text{dcl}(\text{acl}(\phi a, \theta a), \text{acl}(a, \theta a))$. So

$$\phi' \in \text{acl}(\phi a, a) \cap \text{dcl}(\text{acl}(\phi a, \theta a), \text{acl}(a, \theta a))$$

Suppose toward a contradiction that $\phi' \in \text{dcl}(\text{acl}(\phi a), \text{acl}(a))$. Since both a and ϕa are generic in $P^{\text{cl}}(\mathcal{A}')$, by Proposition 2.5, it follows that $\phi' \in \text{dcl}(\phi a, a) = \text{dcl}(a, \phi)$. But note that $\text{tp}(a/\phi)$ is stationary (since $\text{tp}(a)$ is and $a \perp \phi$) and so $\text{dcl}(a, \phi) \cap \text{acl}(\phi) = \text{dcl}(\phi)$. Hence $\phi' \in \text{dcl}(\phi)$, which contradicts $n > 1$. \square

3. INTERNALISING FINITE COVERS IN CCM

Given $T = T^{\text{eq}}$ a complete stable theory, and \mathbf{P} a set of sorts in T , and working in a sufficiently saturated model, recall that a stationary type $q(x) \in S(A)$ is \mathbf{P} -internal if for some $B \supseteq A$, and some $a \models q$ with $a \perp_A B$, we have $a \in \text{dcl}(B\mathbf{P})$. If we replace dcl by acl in this definition we get the notion of *almost \mathbf{P} -internal*.

Definition 3.1. We say that \mathbf{P} *internalises finite covers* if almost \mathbf{P} -internality implies \mathbf{P} -internality.

Example 3.2. The projective line in CCM internalises finite covers, while the field of constants in DCF_0 does not. For the former see Fact 3.1 of [23], and for the latter note that the generic type of $\delta(x^2 - b) = 0$ where b is a fixed non-constant is almost internal but not internal to the field of constants (see [28, Lemma 3.1]).

Remark 3.3. Working in a sufficiently saturated model of T , internalising finite covers has the following characterisation in terms of algebraic and definable closure: for any $M \models T$ and any finite tuple $a \in \mathbf{P}$,

$$(3.1) \quad \text{acl}(Ma) = \text{dcl}(M, \text{acl}(\mathbf{P}(M)a) \cap \mathbf{P}^{\text{eq}})$$

Proof. If $\text{tp}(b/A)$ is almost \mathbf{P} -internal then there is a model $M \supseteq A$ independent from b over A , and a finite tuple $a \in \mathbf{P}$, such that $b \in \text{acl}(Ma)$. Then (3.1) implies that $b \in \text{dcl}(M\mathbf{P})$, showing that $\text{tp}(b/A)$ is internal to \mathbf{P} .

Conversely, assume that \mathbf{P} internalises finite covers and suppose $b \in \text{acl}(Ma)$, where a is a finite tuple from \mathbf{P} . Then $\text{tp}(b/M)$ is almost \mathbf{P} -internal and hence \mathbf{P} -internal. As we are working over a model, b is interdefinable over M with a finite tuple c from \mathbf{P}^{eq} . So $b \in \text{dcl}(Mc)$. On the other hand, $c \in \text{dcl}(Mb) \subseteq \text{acl}(Ma)$. But as a and c are finite tuples from \mathbf{P}^{eq} , it follows by stability (namely, the fact that \mathbf{P} is stably embedded in T) that $c \in \text{acl}(\mathbf{P}(M)a)$. Hence $b \in \text{dcl}(M, \text{acl}(\mathbf{P}(M)a) \cap \mathbf{P}^{\text{eq}})$, as desired. \square

Proposition 3.4. *Suppose X is a simple nonalgebraic complex torus, viewed as a sort in CCM. Then X internalises finite covers.*

Proof. We work in a sufficiently saturated model $\mathcal{A}' \models \text{CCM}$.

See [31] for the basics of the model theory of complex tori.

Suppose $\text{tp}(a/A)$ is stationary and almost X -internal. Then there exists $B \supseteq A$ with $a \perp_A B$, and a tuple c from $X(\mathcal{A}')$ such that $a \in \text{acl}(Bc)$. As X is strongly minimal we may replace c by an acl-basis for c over B ; that is, we may assume c is a B -independent ℓ -tuple of generic points in X over B . We will show that $\text{tp}(ac/B)$ is X -internal – this will suffice as it implies that $\text{tp}(a/B)$ is X -internal, and $\text{tp}(a/B)$ is the nonforking extension of $\text{tp}(a/A)$.

Extending B we may assume that $\text{tp}(ac/B)$ is stationary. Replacing B by the canonical base of $\text{tp}(ac/B)$, we may assume that $B = b$ is a finite tuple. Let $S := \text{loc}(b)$ and $Y := \text{loc}(bac)$. Note that $\text{loc}(bc) = S \times X^\ell$. Co-ordinate projections yield the following commuting diagram of surjective morphisms:

$$\begin{array}{ccc} Y & \xrightarrow{\pi} & S \times X^\ell \\ & \searrow & \swarrow \\ & S & \end{array}$$

Note that π is generally finite-to-one since $a \in \text{acl}(bc)$. Since $\text{tp}(ac/b)$ is the generic type of the generic fibre of $Y \rightarrow S$, to prove that it is X -internal it will suffice to prove that, possibly after base change, there is a dominant meromorphic map back from $S \times X^\ell$ to Y over S . This is what we now do.

Consider the Stein factorisation of π ,

$$\begin{array}{ccccc} Y & \xrightarrow{f} & \tilde{Y} & \xrightarrow{\tilde{\pi}} & S \times X^\ell \\ & \searrow & \downarrow & \swarrow & \\ & & S & & \end{array}$$

where f is a bimeromorphism and $\tilde{\pi}$ is now finite-to-one everywhere. Next we take a normalisation of \tilde{Y} to get

$$\begin{array}{ccccc}
 & & \hat{Y} & & \\
 & & \downarrow & \searrow \hat{\pi} & \\
 Y & \xrightarrow{f} & \tilde{Y} & \xrightarrow{\tilde{\pi}} & S \times X^\ell \\
 & \searrow & \downarrow & \swarrow & \\
 & & S & &
 \end{array}$$

where \hat{Y} is normal and $\hat{\pi}$ remains a finite morphism.

Claim 3.5. *Possibly after a (finite) base change, the general fibres of $\hat{Y} \rightarrow S$ are complex tori with a uniformly definable group structure, and $\hat{\pi}$ restricted to these fibres is an isogeny.*

Proof of Claim 3.5. We are grateful to Frédéric Campana for some useful correspondence around these issues.

Fixing a general $s \in S$ we have that \hat{Y}_s is normal [4, Théorème 2] and connected, X^ℓ is smooth and $\hat{\pi}_s : \hat{Y}_s \rightarrow X^\ell$ is finite. It follows by the purity of branch locus theorem [11, Theorem VII.1.6] that the ramification locus of $\hat{\pi}_s$ is either empty or complex analytic of codimension 1 in \hat{Y}_s . In the latter case the image of this ramification locus would have to be a codimension 1 complex analytic subset of X^ℓ , but the fact that X is strongly minimal with $\dim X > 1$ makes this impossible (any complex analytic subset of X^ℓ has dimension a multiple of $\dim X$). Hence $\hat{\pi}_s$ is an unramified covering. It follows that \hat{Y}_s itself has the structure of a complex torus such that $\hat{\pi}_s : \hat{Y}_s \rightarrow X^\ell$ is an isogeny.

We have not yet proved that this group structure on \hat{Y}_s with respect to which $\hat{\pi}_s$ is an isogeny is *uniform* in s . Note, however, that if $\Gamma \leq (X^\ell)^3$ is the graph of the group structure on X^ℓ , then for each general $s \in S$, $\hat{\pi}_s^{-1}(\Gamma) \leq \hat{Y}_s^3$ is a possibly disconnected subtorus that is uniformly definable in s . Moreover, since $\hat{\pi}_s : \hat{Y}_s \rightarrow X^\ell$ is finite, $\dim \hat{\pi}_s^{-1}(\Gamma) = \dim \Gamma = 2\ell \dim X$. Now, the graph of the (not yet known to be uniformly definable) group operation on \hat{Y}_s is of dimension $2 \dim \hat{Y}_s = 2\ell \dim X$ also, and it is a connected subtorus of $\hat{\pi}_s^{-1}(\Gamma)$. It must therefore be the connected component of identity. This is still not enough because we do not yet know that the identity of \hat{Y}_s can be picked out definably in s . However, it does follow that each connected component of $\hat{\pi}_s^{-1}(\Gamma)$ is the graph of *some* group structure on \hat{Y}_s with respect to which $\hat{\pi}_s$ is an isogeny. Indeed, if H is any such connected component then it is a translate of the graph of the group operation on Y_s by some $(a, b, c) \in \hat{\pi}_s^{-1}(\Gamma)$. So H is itself the graph of a new group operation on Y_s , namely $x \oplus y := x + y - (a + b - c)$. That $\hat{\pi}_s$ remains an isogeny with respect to this new group operation follows from the fact that $\hat{\pi}_s(a) + \hat{\pi}_s(b) = \hat{\pi}_s(c)$.

So it suffices to show that after base change the connected components of $\hat{\pi}_s^{-1}(\Gamma)$ are uniformly definable. This we now point out. Let $Z \subset \hat{Y} \times_S \hat{Y} \times_S \hat{Y}$ be a maximal dimensional irreducible component of $\hat{\pi}^{-1}(S \times \Gamma)$ that projects onto S . Let

$$Z \longrightarrow T \xrightarrow{\theta} S$$

be the Stein factorisation of $Z \rightarrow S$. So for general $t \in T$, Z_t is a connected component of $\widehat{\pi}_s^{-1}(\Gamma)$ where $s = \theta(t)$. Taking the finite base change $\theta : T \rightarrow S$, we obtain $\widehat{\pi}_T : \widehat{Y}_T \rightarrow T \times X^\ell$. Then for general $t \in T$ we have that $\widehat{\pi}_t^{-1}(\Gamma)$ is connected, and so defines, uniformly in t , a group structure with respect to which $\widehat{\pi}_t$ is an isogeny, as required. \square

By the Claim, we may assume that the group structure on the general fibres \widehat{Y}_s with respect to which $\widehat{\pi}_s$ is an isogeny is uniformly definable over s . Let n be the size of the kernel of $\widehat{\pi}_s$ for $s \in S$ general. We can define $\rho : S \times X^\ell \rightarrow \widehat{Y}$ over S to be the meromorphic map such that for general $s \in S$, $\rho_s \circ \widehat{\pi}_s$ is multiplication by n on \widehat{Y}_s , by setting $\rho_s(x) := ny$ for any $y \in \widehat{\pi}_s^{-1}(x)$. Finally, looking at the diagram before the statement of Claim 3.5, we see that \widehat{Y} admits a dominant meromorphic map onto Y over S since f is bimeromorphic. We therefore have a dominant meromorphic map from $S \times X^\ell$ to Y over S , witnessing that $\text{tp}(a/A)$ is indeed internal to X . \square

We record here for future use the following much easier class of examples of compact complex manifolds which internalise finite covers in CCM.

Proposition 3.6. *Let X be a simply connected nonalgebraic strongly minimal compact complex variety. Then X internalises finite covers.*

Proof. The essential point is just that there are no finite covers to internalise. This is a uniform version of Lemma 2.4.

As in the first part of the proof of Proposition 3.4, it suffices to see that for any irreducible S , if $\widehat{\pi} : \widehat{Y} \rightarrow S \times X^\ell$ is a finite cover over S , and for general $s \in S$ the fibre \widehat{Y}_s is irreducible and the map $\widehat{\pi}_s : \widehat{Y}_s \rightarrow X^\ell$ is unramified, then $\widehat{\pi}$ has a meromorphic splitting.

But X^ℓ is simply connected since X is, so $\widehat{\pi}_s$ is an isomorphism for general $s \in S$. So π is a bimeromorphism. \square

4. GENERIC AUTOMORPHISMS: IMAGINARIES AND STABLE EMBEDDEDNESS

Suppose $T = T^{\text{eq}}$ is a complete stable theory admitting quantifier elimination in a language L . Consider the expanded language $L_\sigma := L \cup \{\sigma\}$ where σ is a unary function symbol, and in this language the universal theory

$$T_{\forall, \sigma} := T_\forall \cup \{\sigma \text{ is an } L\text{-embedding}\}.$$

Recall that by TA is meant the model companion of $T_{\forall, \sigma}$, when it exists. In this section we will review what is known about TA and give a characterisation for stable embeddedness in TA of sorts from L .

We will be mostly interested in the case when $T = \text{CCM}$. The first two authors and Gavrilovich showed in [3] that CCMA exists. In fact, they showed the existence of TA for T the theory of any ω_1 -compact noetherian topological structure with quantifier elimination and in which irreducibility is definable.

Suppose TA exists. We will use tp^- , acl^- , and \bigcup^T when dealing with the reduct $\mathbb{U} \models T$ of some $(\mathbb{U}, \sigma) \models TA$. We work generally in the real sorts of $(\mathbb{U}, \sigma) \models TA$, and not in $(\mathbb{U}, \sigma)^{\text{eq}}$ unless explicitly stated otherwise. The following properties of TA are due to Chatzidakis and Pillay [9, §3].

- (1) Quantifier reduction: $\text{tp}(a/A) = \text{tp}(b/A)$ if and only if there is an L_σ -isomorphism from $\text{acl}(Aa)$ to $\text{acl}(Ab)$ that takes a to b and fixes A . In particular, the completions of TA are determined by the isomorphism type of $(\text{acl}^-(\emptyset), \sigma)$.
- (2) For any set A , $\text{acl}(A)$ is the acl^- -closure of the inversive closure of A .
- (3) Every completion of TA is simple and

$$A \underset{E}{\downarrow} B \text{ if and only if } \text{acl}(A) \underset{\text{acl}(E)}{\overset{T}{\downarrow}} \text{acl}(B).$$

If T is superstable then every completion of TA is supersimple.

Here the *inversive closure* of A is by definition the set obtained from A by closing off under σ and σ^{-1} .

We now fix a sufficiently saturated model $(\mathbb{U}, \sigma) \models TA$.

What about imaginaries in models of TA (if it exists)? Recall that we assume $T = T^{\text{eq}}$, i.e. we concentrate on the issue of eliminating / classifying imaginaries in (\mathbb{U}, σ) relative to imaginaries from \mathbb{U} . For T strongly minimal such that $\text{acl}^-(\emptyset)$ is infinite, it is shown in [9] that every completion of TA eliminates imaginaries. This is not true in general. Indeed, in [17, Proposition 4.7] Hrushovski shows that for stable T , 3-uniqueness implies elimination of imaginaries in any completion of TA . Moreover, it follows from [17, Proposition 4.5] that the converse holds for T superstable. Indeed, any completion of TA is then supersimple, and so algebraically closed sets are amalgamation bases, which implies 3-existence in the terminology of that paper. However, as Bradd Hart pointed out to us, Hrushovski's method does give:

Fact 4.1. *Any completion of TA admits geometric elimination of imaginaries: every element of $(\mathbb{U}, \sigma)^{\text{eq}}$ is interalgebraic with a finite tuple from \mathbb{U} .*

Proof. The proof of Proposition 4.7 of [17] shows that if $e = a/E$ is in $(\mathbb{U}, \sigma)^{\text{eq}}$, where a is a finite tuple from the home sort and E is a definable equivalence relation, then there exists in the E -class of a an element, a' , such that $a' \underset{\text{acl}^{\text{eq}}(e) \cap \mathbb{U}}{\downarrow} a$. It follows

that $e \in \text{acl}^{\text{eq}}(a) \cap \text{acl}^{\text{eq}}(a') \subseteq \text{acl}^{\text{eq}}(\text{acl}^{\text{eq}}(e) \cap \mathbb{U})$, as desired. \square

Considering $T = \text{CCM}$ we obtain that CCMA is complete and satisfies properties (1) through (3) above, and it admits geometric elimination of imaginaries. However, since CCM fails 3-uniqueness by Theorem 2.1,

Theorem 4.2. *CCMA does not eliminate imaginaries.*

Remark 4.3. We can explicitly describe a finite imaginary which is not eliminated. Let $\pi : P \rightarrow X$ be as in Section 2. Let E be the relation on the fixed set $P^f := \{a \in P \mid \sigma(a) = a\}$ of P defined by

$$aEa' \iff \exists \theta \in K^* (a' = \theta^2 a \wedge \sigma(\theta) = \theta).$$

This is a \emptyset -definable equivalence relation, with two equivalence classes on each fibre P_b^f . Let $a \in P^f$ be generic in P ; such exists by the axioms for CCMA. Let $b := \pi(a)$. Let $\theta \in K^*$ be generic over a and such that $\sigma(\theta) = -\theta$. By Proposition 2.5 and Lemma 2.4, $\text{acl}(a) = \text{acl}^-(a) = \text{dcl}^-(a)$ and $\text{acl}(b) = \text{acl}^-(b) = \text{dcl}^-(b)$. Now $\theta^2 a \in P_b^f$ is also generic in P_b , and so it follows from the quantifier

reduction that $\text{tp}(a/\text{acl}(b)) = \text{tp}(\theta^2 a/\text{acl}(b))$. But $P_b^f/E = \{a/E, (\theta^2 a)/E\}$. So $a/E \in \text{acl}^{eq}(b) \setminus \text{acl}(b)$.

Next we address the issue of stable embeddedness in TA .

Fact 4.4. *Assume TA exists. Suppose \mathbf{P} is a set of sorts in \mathbb{U} , $L^{\mathbf{P}} \subseteq L$ is a language describing the induced structure on \mathbf{P} (over \emptyset), and $T^{\mathbf{P}} = \text{Th}_{L^{\mathbf{P}}}(\mathbf{P})$. Assume that $T^{\mathbf{P}}$ has QE. Then the following hold:*

- (i) $T^{\mathbf{P}}A$ exists.
- (ii) If $(M, \sigma) \models TA$, then $(\mathbf{P}(M), \sigma) \models T^{\mathbf{P}}A$.
- (iii) Conversely, for every $(P, \sigma) \models T^{\mathbf{P}}A$ there is a model $(M, \sigma) \models TA$ such that $(\mathbf{P}(M), \sigma) \succ (P, \sigma)$.

Proof. This is folklore. It follows from Chatzidakis and Pillay [9, 3.4 and 3.5]. As we could not find it explicitly stated, we give a proof. Closing off \mathbf{P} under \emptyset -definable quotients (which does not change the truth value of the statements (i), (ii) or (iii)), we may assume that $T^{\mathbf{P}}$ eliminates imaginaries.

Let $T_{\sigma} := T \cup \{\sigma \text{ is an } L\text{-automorphism}\}$.

Let $\kappa = |T|^+$, and let $\mathcal{C}_{T, \sigma}$ be the class of L_{σ} -structures (A, σ) , where A is an algebraically closed subset of a model of T and σ is an elementary permutation of A . A model (M, σ) of T_{σ} is said to be κ -generic if whenever $(A, \sigma) \subseteq (B, \sigma)$ are elements of $\mathcal{C}_{T, \sigma}$ of cardinality $< \kappa$, any embedding of (A, σ) into (M, σ) extends to an embedding of (B, σ) into (M, σ) .

A straightforward argument shows that if (M, σ) is κ -generic, then $(\mathbf{P}(M), \sigma)$ is κ -generic (for $\mathcal{C}_{T^{\mathbf{P}}, \sigma}$). Indeed, this follows from the fact that if $A = \text{dcl}(A) \subseteq M \models T$ and $b \in \mathbf{P}(M)$, then $\text{tp}(b/A \cap \mathbf{P}) \models \text{tp}(b/A)$ by stable embeddedness of \mathbf{P} . Moreover, on general grounds, if (M, σ) is κ -saturated, so is $(\mathbf{P}(M), \sigma)$.

By [9], TA exists if and only if every $(M_0, \sigma_0) \models T_{\sigma}$ embeds into some κ -generic $(M, \sigma) \models T_{\sigma}$ which is κ -saturated, and TA is then equal to the common theory of all κ -generic (M, σ) . (It then also follows that a model of TA is κ -saturated if and only if it is κ -generic.)

Now, any $(P, \sigma) \models T_{\sigma}^{\mathbf{P}}$ embeds into some $(M, \sigma) \models T_{\sigma}$ which is κ -generic for $\mathcal{C}_{T, \sigma}$ and κ -saturated, so $(P, \sigma) \subseteq (\mathbf{P}(M), \sigma)$ with $(\mathbf{P}(M), \sigma)$ κ -generic for $\mathcal{C}_{T^{\mathbf{P}}, \sigma}$ and κ -saturated. Thus, the existence of TA implies that of $T^{\mathbf{P}}A$, proving (i).

To prove (ii), one may assume that (M, σ) is κ -saturated. By the previous paragraph, $(\mathbf{P}(M), \sigma)$ is κ -generic, so in particular a model of $T^{\mathbf{P}}A$.

Part (iii) follows from the fact that every $(P, \sigma) \models T^{\mathbf{P}}A$ embeds into some $(M, \sigma) \models TA$. By (ii), $(\mathbf{P}(M), \sigma) \models T^{\mathbf{P}}A$, and so $(\mathbf{P}(M), \sigma) \succ (P, \sigma)$ by model-completeness of $T^{\mathbf{P}}A$. \square

Proposition 4.5. *Let \mathbf{P} and $L^{\mathbf{P}}$ be as above, such that $T^{\mathbf{P}}$ eliminates quantifiers. Assume TA exists.*

- (i) *If \mathbf{P} internalises finite covers² in T then \mathbf{P} is stably embedded in (\mathbb{U}, σ) , with induced structure given by $L_{\sigma}^{\mathbf{P}}$.*
- (ii) *Conversely, assume that \mathbf{P} is stably embedded in (\mathbb{U}, σ) with induced structure given by $L_{\sigma}^{\mathbf{P}}$. Assume in addition that σ restricted to $\text{acl}^-(\emptyset)$ is the identity. Then \mathbf{P} internalises finite covers in T .*

²See Definition 3.1.

Proof. Note that \mathbf{P}^{eq} internalises finite covers if and only if \mathbf{P} does, so we may close off \mathbf{P} under definable quotients and thus assume that $\mathbf{P} = \mathbf{P}^{\text{eq}}$, i.e., that $T^{\mathbf{P}}$ eliminates imaginaries.

To prove (i), suppose $(M, \sigma) \preceq (\mathbb{U}, \sigma)$, and let a_1, a_2 be finite tuples from \mathbf{P} such that $\text{tp}_{L_{\sigma}^{\mathbf{P}}}(a_1/\mathbf{P}(M)) = \text{tp}_{L_{\sigma}^{\mathbf{P}}}(a_2/\mathbf{P}(M))$. We have to show that $\text{tp}(a_1/M) = \text{tp}(a_2/M)$.

By assumption, there is an L_{σ} -automorphism of \mathbf{P} that fixes $\mathbf{P}(M)$ and takes a_1 to a_2 . For $i = 1, 2$, set

$$A_i := \text{acl}(\mathbf{P}(M)a_i) \cap \mathbf{P} = \text{acl}^{-}(\mathbf{P}(M)(\sigma^n a_i)_{n \in \mathbb{Z}}) \cap \mathbf{P}$$

We obtain an L_{σ} -isomorphism $f : A_1 \rightarrow A_2$ that fixes $\mathbf{P}(M)$ and takes a_1 to a_2 . By stable embeddedness of \mathbf{P} in \mathbb{U} and the choice of $L_{\sigma}^{\mathbf{P}}$, f extends further to an L -isomorphism $F : \text{dcl}^{-}(MA_1) \rightarrow \text{dcl}^{-}(MA_2)$ fixing M . Note that $M \cup A_i$, and hence $\text{dcl}^{-}(MA_i)$, are σ -invariant, and that F is in fact an L_{σ} -isomorphism (since f is). But

$$\text{dcl}^{-}(MA_i) = \text{acl}^{-}(M(\sigma^n a_i)_{n \in \mathbb{Z}}) = \text{acl}(Ma_i)$$

since \mathbf{P} internalises finite covers in T , see Remark 3.3, noting that it applies to infinite tuples by finite character of acl and dcl . Quantifier reduction in TA now implies that $\text{tp}(a_1/M) = \text{tp}(a_2/M)$.

To prove (ii), now assume that σ restricted to $\text{acl}^{-}(\emptyset)$ is the identity, and that \mathbf{P} is stably embedded in (\mathbb{U}, σ) with induced structure given by $L_{\sigma}^{\mathbf{P}}$.

Suppose for contradiction that \mathbf{P} does not internalise finite covers. By the characterisation given in Remark 3.3, there is $M \models T$ and a finite tuple $a \in \mathbf{P}$ such that $\text{acl}^{-}(Ma) \supsetneq \text{dcl}^{-}(M, \text{acl}^{-}(\mathbf{P}(M)a) \cap \mathbf{P})$. Using that $\sigma \upharpoonright_{\text{acl}^{-}(\emptyset)} = \text{id}$ and saturation of (\mathbb{U}, σ) we may find $a_1, a_2 \in \mathbf{P}$ such that, letting $A_i := \text{acl}^{-}(Ma_i)$ and $B_i := \text{dcl}^{-}(M, \text{acl}^{-}(\mathbf{P}(M)a_i) \cap \mathbf{P})$ for $i = 1, 2$, the following properties hold:

- $\text{tp}^{-}(a_i/M) = \text{tp}^{-}(a/M)$ for $i = 1, 2$;
- σ restricted to A_1 and to B_2 is the identity, and $\sigma \upharpoonright_{A_2} \neq \text{id}$;

Note that $T^{\mathbf{P}}A$ exists by Fact 4.4. Using quantifier reduction in $L_{\sigma}^{\mathbf{P}}$, it thus follows in particular that $\text{tp}_{L_{\sigma}^{\mathbf{P}}}(a_1/\mathbf{P}(M)) = \text{tp}_{L_{\sigma}^{\mathbf{P}}}(a_2/\mathbf{P}(M))$. Now let $(M, \text{id}) \subseteq (N, \sigma) \preceq (\mathbb{U}, \sigma)$, where N is small. By [9, Thm 3.7], as $(\mathbf{P}(M), \text{id}) \models T_{\sigma}^{\mathbf{P}}$, replacing (N, σ) by an isomorphic copy if necessary, we may assume that $\text{tp}_{L_{\sigma}^{\mathbf{P}}}(a_1/\mathbf{P}(N)) = \text{tp}_{L_{\sigma}^{\mathbf{P}}}(a_2/\mathbf{P}(N))$.

On the other hand, we clearly have $\text{tp}(a_1/M) \neq \text{tp}(a_2/M)$ by construction, and so in particular $\text{tp}(a_1/N) \neq \text{tp}(a_2/N)$, contradicting the assumption. \square

We expect that the following question has a positive answer, but we did not explore it in depth.

Question 4.6. Does Proposition 4.5(ii) hold without the assumption on $\sigma \upharpoonright_{\text{acl}(\emptyset)}$?

Specialising to the case when $T = \text{CCM}$ we obtain:

Corollary 4.7. *For a collection \mathbf{P} of sorts in CCM, the following are equivalent:*

- \mathbf{P} internalises finite covers in CCM;
- \mathbf{P} is stably embedded in CCMA, with induced structure given by $L_{\sigma}^{\mathbf{P}}$.

Theorem 4.8. *Suppose X is either a complex projective variety, a nonalgebraic simple complex torus, or a strongly minimal simply connected nonalgebraic compact complex manifold. Working in a sufficiently saturated model $(\mathcal{A}', \sigma) \models \text{CCMA}$, let*

\mathcal{X}' denote the structure induced on $X(\mathcal{A}')$ by \mathcal{A}' . Then (\mathcal{X}', σ) is purely stably embedded in (\mathcal{A}', σ) .

In particular, ACFA (with the elements of \mathbb{C} named) is purely stably embedded in CCMA.

Proof. This follows from the previous corollary by the fact that X internalises finite covers in CCM; for X a projective variety that is Fact 3.1 of [23], and the other cases are given by Propositions 3.4 and 3.6 above.

Let $(K, +, \times)$ be the interpretation of the complex field in \mathcal{A}' . The “in particular” clause is claiming that $(K, +, \times, \sigma)$ is purely stably embedded in (\mathcal{A}', σ) . This follows from the main statement because $(K, +, \times)$, with the elements of \mathbb{C} named, and \mathcal{X}' where $X = \mathbb{P}(\mathbb{C})$, are bi-interpretable without parameters. \square

Remark 4.9 (Bustamante [6, Prop. 2.3.5]). ACFA is also interpretable in the theory of differentially closed fields (of characteristic 0) with a generic automorphism, by considering the field of constants. However, ACFA is not in this way purely stably embedded in DCFA. Indeed, let $(K, \delta, \sigma) \models \text{DCFA}$ with $C = \text{const}(\delta)$, and $b \in \text{Fix}(\sigma) \setminus C$. Then $\{y \in C \mid \exists x [\sigma(x) = x \wedge x^2 - b = y]\}$ is not definable in $(C, +, \times, \sigma)$.

5. FINITE-DIMENSIONAL TYPES IN CCMA

Everything we do in this section could be done in the more general setting of TA for T the theory of an ω_1 -compact noetherian topological structure with quantifier elimination and in which irreducibility is definable. But we stick to CCMA for the sake of concreteness.

Fix a sufficiently saturated $(\mathcal{A}', \sigma) \models \text{CCMA}$.

Definition 5.1. By a CCM- σ -variety is meant a pair (F, G) where F is an irreducible closed set in \mathcal{A}' and $G \subseteq F \times F^\sigma$ is an irreducible closed subset whose projections to both F and F^σ are surjective and generically finite-to-one. This data gives rise to the following definable set in (\mathcal{A}', σ)

$$(F, G)^\sharp := \{a \in F : (a, \sigma a) \in G\}.$$

Note that $F^\sigma := \sigma(F)$ is again a closed set in \mathcal{A}' ; if $F = X_a$ where $X \subset S \times Z$ is a complex analytic subset of a product of compact complex varieties and $a \in S(\mathcal{A}')$, then $F^\sigma = X_{\sigma(a)}$.

We can associate to a meromorphic dynamical system (X, f) the CCM- σ -variety (F, G) where $F = X(\mathcal{A}')$ and G is the (set of \mathcal{A}' -points of the) graph of f . In fact, a CCM- σ -variety should be viewed as the generalisation of a meromorphic dynamical system where we allow finite-to-finite meromorphic correspondences in place of dominant meromorphic maps.

CCM- σ -varieties give rise to finite-dimensional types, in the following sense.

Definition 5.2. Given an inversive (i.e., closed under σ and σ^{-1}) set A , and a tuple a , we set

$$\dim_\sigma(a/A) := (\dim \text{loc}((a, \sigma a, \dots, \sigma^n a)/A))_{n < \omega}$$

where \dim and loc are meant in the sense of CCM. We say that $\text{tp}(a/A)$ is *finite-dimensional* if $\dim_\sigma(a/A)$ is eventually constant, in which case we often write $\dim_\sigma(a/A) = d$ where d is that eventual value.

Note that this dimension witnesses independence: for all inversive $B \supseteq A$,

$$(5.1) \quad a \downarrow_A B \iff \dim_\sigma(a/B) = \dim_\sigma(a/A)$$

This follows rather easily from (2) and (3) of Section 4, together with the fact that complex analytic dimension witnesses independence in CCM.

Lemma 5.3. *Suppose (F, G) is a CCM- σ -variety over inversive A . Then there exists $c \in (F, G)^\sharp$ that is CCM-generic in F over A . Such a point satisfies:*

- (i) $\text{tp}(c/A)$ is finite-dimensional with $\dim_\sigma(c/A) = \dim F$
- (ii) for all inversive $B \supseteq A$, $c \downarrow_A B$ if and only if $\text{loc}(c/B) = F$.

Proof. That we can find $c \in (F, G)^\sharp$ avoiding any particular proper A -closed subset of F follows from the axiomatisation of CCMA given in [3, Proposition 5.2]. By saturation we get a generic of F over A in $(F, G)^\sharp$. (Note that this does not use the fact that the projections are finite-to-one, only that they are surjective.)

For (i), note that c being generic in F and $(c, \sigma(c)) \in G$ implies that $\sigma(c) \in \text{acl}(Ac)$. Applying σ repeatedly gives $\sigma^n(c) \in \text{acl}(Ac)$ for all n . Hence $\dim_\sigma(c/A) = (\dim F, \dim F, \dots)$.

Part (ii) now follows. Indeed, from (5.1), we have that $c \downarrow_A B$ if and only if $\dim \text{loc}((c, \sigma c, \dots, \sigma^n c)/B) = \dim F$ for all n , if and only if $\dim \text{loc}(c/B) = \dim F$. By irreducibility of F , the latter is equivalent to $\text{loc}(c/B) = F$. \square

Remark 5.4. The (absolute) irreducibility of F and G in the definition of a CCM- σ -variety is essential. Suppose F_1 and F_2 are disjoint irreducible closed sets such that $F_1^\sigma = F_2$ and $F_2^\sigma = F_1$. Let $F = F_1 \cup F_2$ and let G be the union of the diagonals in $F_1 \times F_1$ and $F_2 \times F_2$. Then G projects generically finite-to-one onto both F and $F^\sigma = F$, but $(F, G)^\sharp = \emptyset$. If in addition F is A -definable but F_1 is not A -definable, then F and G will even be A -irreducible.

We conclude this section by pointing out that CCM- σ -varieties capture all the finite-dimensional types in CCMA.

Lemma 5.5. *Suppose A is acl-closed and c is such that $\dim_\sigma(c/A)$ is finite. Then there exists $N \geq 0$ and a CCM- σ -variety (F, G) over A , such that $(c, \sigma c, \dots, \sigma^N c)$ is generic in F over A and contained in $(F, G)^\sharp$.*

Proof. Let N and d be such that

$$d = \dim \text{loc}((c, \sigma c, \dots, \sigma^N c)/A) = \dim \text{loc}((c, \sigma c, \dots, \sigma^{N+1} c)/A)$$

Let $\bar{c} := (c, \sigma c, \dots, \sigma^N c)$, $F := \text{loc}(\bar{c}/A)$ and $G := \text{loc}((\bar{c}, \sigma \bar{c})/A) \subseteq F \times F^\sigma$. The assumption that $A = \text{acl}(A)$ ensures that F and G are irreducible. Note that $\dim G = \dim \text{loc}((c, \sigma c, \dots, \sigma^{N+1} c)/A) = d = \dim F = \dim F^\sigma$. Moreover, both projections of G are onto as \bar{c} is generic in F over A , and so $\sigma \bar{c}$ is generic in F^σ over A also. Hence these projections must be generically finite-to-one. \square

Remark 5.6. If furthermore we take N in Lemma 5.5 large enough so that $\text{mult}(\sigma^{N+1} c/A, c, \dots, \sigma^N c)$ is minimal, then the CCM quantifier-free type of c over A is determined by saying that $(c, \sigma c, \dots, \sigma^N c) \in (F, G)^\sharp$ and c is generic in F over A .

6. DIFFERENCE-ANALYTIC JET SPACES AND THE ZILBER DICHOTOMY

The goal of this section is to prove the Zilber dichotomy for finite-dimensional SU -rank one types in CCMA: either they are one-based or they are nonorthogonal to the fixed field (of the canonical interpretation of ACFA in CCMA). For ACFA itself this was done first by Chatzidakis and Hrushovski in [7], but then a much simpler proof was given by Pillay and Ziegler in [30]. It is this latter argument, which goes via an appropriate notion of “jet space” and actually proves something stronger than the dichotomy (namely the Canonical Base Property or the CBP), that we will follow here, and extend to all of CCMA.

In the appendix to this paper we have reviewed the uniformly definable construction of jet spaces in complex analytic geometry. Given a holomorphic map $\pi : X \rightarrow S$ of compact complex varieties, there exists a CCM-definable complex variety $\text{Jet}^n(X/S) \rightarrow X$ such that for all $x \in X$

$$\text{Jet}^n(X/S)_x = \text{Jet}^n(X_{\pi(x)})_x = \text{Hom}_{\mathbb{C}}(\mathfrak{m}_{X_{\pi(x)},x}/\mathfrak{m}_{X_{\pi(x)},x}^{n+1}, \mathbb{C})$$

and such that the vector space structure on these fibres is uniformly definable in CCM. Here $\mathfrak{m}_{X_{\pi(x)},x}$ denotes the maximal ideal of the local ring of germs of holomorphic functions on the fibre $X_{\pi(x)}$ at the point x . We point the reader to the appendix for further details.

The uniformity allows us to define jet spaces of nonstandard closed sets.

Definition 6.1. Suppose $F = X_a$ is a closed set in CCM where $X \subseteq S \times Z$ and $a \in S(\mathcal{A}')$. Then by the *n th jet space of F* , which we will denote by $\text{Jet}^n(F) \rightarrow F$, we will mean the restriction of $\text{Jet}^n(X/S)(\mathcal{A}') \rightarrow X(\mathcal{A}')$ to X_a .

The fibres of $\text{Jet}^n(F) \rightarrow F$ are therefore uniformly finite dimensional K -vector spaces, where recall that $(K, +, \times)$ is the interpretation of the complex field in \mathcal{A}' . Moreover, by the functoriality of the construction of jet spaces, if F is an irreducible closed subset of an irreducible closed set G , and $c \in F$, then $\text{Jet}^n(F)_c$ is naturally a K -linear subspace of $\text{Jet}^n(G)_c$. The main point of this construction is the following proposition that says these jet spaces effect a linearisation of irreducible closed sets.

Proposition 6.2. *Suppose S and Z are compact complex varieties and $X \subset S \times Z$ is a subvariety. There exists $n > 0$ such that for all $a, b \in S(\mathcal{A}')$, if X_a and X_b are irreducible, pass through some $c \in Z(\mathcal{A}')$, and have $\text{Jet}^n(X_a)_c = \text{Jet}^n(X_b)_c$ as K -subspaces of $\text{Jet}^n(Z)_c$, then $X_a = X_b$.*

Proof. The statement that is claimed by the proposition (for fixed S, Z, X) is first-order in the language of CCM (using the fact that irreducibility is definable), and hence it suffices to prove that it is true in the standard model. Now, working in the standard model, by Lemma A.3, it is the case that for any $s, s' \in S(\mathcal{A})$ if X_s and $X_{s'}$ are irreducible and pass through $z \in Z(\mathcal{A})$ with $\text{Jet}^n(X_s)_z = \text{Jet}^n(X_{s'})_z$ for all $n \geq 0$, then $X_s = X_{s'}$. To see that we can uniformly bound n is a straightforward compactness argument: For each $n > 0$, let $\phi_n(s, s', z)$ be the formula expressing that X_s and $X_{s'}$ are irreducible, pass through z , satisfy $\text{Jet}^n(X_s)_z = \text{Jet}^n(X_{s'})_z$ but $X_s \neq X_{s'}$. We are using here the definability of irreducibility and the fact that the jet spaces are uniformly definable in order to write down ϕ_n . Note that $\phi_n(s, s', z) \rightarrow \phi_m(s, s', z)$ for all $m \leq n$. What we need to prove is that for some n , ϕ_n is not satisfiable in \mathcal{A} . But by ω_1 -compactness of \mathcal{A} , if each ϕ_n were satisfiable then we would find s, s', z such that $X_s \neq X_{s'}$ but $\text{Jet}^n(X_s)_z = \text{Jet}^n(X_{s'})_z$ for all $n \geq 0$, contradicting what was previously established. \square

Now we pass to CCMA and define jet spaces there using the nonstandard complex analytic jet spaces described above, in very much the same way that Pillay and Ziegler use algebraic jet spaces to define jet spaces for finite-dimensional difference-algebraic varieties in [30, §4].

Work in a sufficiently saturated model $(\mathcal{A}', \sigma) \models \text{CCMA}$, and fix a CCM- σ -variety (F, G) defined over $B = \text{acl}(B)$. We can arrange things so that $F = X_a$ and $G = W_{(a, \sigma(a))} \subset X_a \times X_{\sigma(a)}$ where

- a is a tuple from B that is a generic point of a compact complex variety S ,
- $X \rightarrow S$ is a *fibre space*; that is, the general fibres are irreducible,
- $(a, \sigma(a))$ is a generic point of a subvariety $T \subseteq S^2$, and
- $W \subseteq X^2$ is a subvariety such that the induced map $W \rightarrow T$ is a fibre space.

The fact that G projects generically finite-to-one onto F and F^σ implies, by definability of dimension, that in the standard model, for general $t = (s_1, s_2) \in T(\mathcal{A})$, $W_t \subseteq X_{s_1} \times X_{s_2}$ projects generically finite-to-one onto each co-ordinate. It follows that for any $n > 0$ and for general $x_1 \in X_{s_1}(\mathcal{A})$ and $x_2 \in X_{s_2}(\mathcal{A})$, $\text{Jet}^n(W_t)_{(x_1, x_2)}$ is the graph of a \mathbb{C} -linear isomorphism from $\text{Jet}^n(X_{s_1})_{x_1}$ to $\text{Jet}^n(X_{s_2})_{x_2}$. (For this latter property of jet spaces, see for example Lemma 5.10 of [25], though in the algebraic setting.) As this is a definable property we get for $c \in (F, G)^\sharp$, with c generic in F over B , that $\text{Jet}^n(G)_{(c, \sigma(c))}$ is the graph of a (CCM-definable) K -linear isomorphism $g : \text{Jet}^n(F)_c \rightarrow \text{Jet}^n(F^\sigma)_{\sigma(c)}$.

Definition 6.3. Suppose (F, G) is a CCM- σ -variety over $B = \text{acl}(B)$, and $c \in (F, G)^\sharp$ is generic in F over B . By the *n th jet space of $(F, G)^\sharp$ at c* we mean the Bc -definable $\text{Fix}(K, \sigma)$ -vector subspace

$$\text{Jet}^n(F, G)_c^\sharp := (\text{Jet}^n(F)_c, \text{Jet}^n(G)_{(c, \sigma(c))})^\sharp = \{v \in \text{Jet}^n(F)_c : \sigma(v) = g(v)\}$$

Note that $g^{-1}\sigma : \text{Jet}^n(F)_c \rightarrow \text{Jet}^n(F)_c$ endows $\text{Jet}^n(F)_c$ with the structure of a σ -module over the difference field (K, σ) ; namely $g^{-1}\sigma(rv) = g^{-1}(\sigma(r)\sigma(v)) = \sigma(r)g^{-1}\sigma(v)$ for all $r \in K$ and $v \in \text{Jet}^n(F)_c$. This uses the fact that scalar multiplication on $\text{Jet}^n(F)_c$ is obtained by the restriction of the 0-definable scalar multiplication on $\text{Jet}^n X$, and hence commutes with σ . In any case, we have that $\text{Jet}^n(F, G)_c^\sharp$ is the fixed set of this σ -module, which is thus naturally a $\text{Fix}(K, \sigma)$ -vector subspace. Moreover, by [30, Lemma 4.2(ii)], the fact that (K, σ) is existentially closed implies that

$$\text{Jet}^n(F)_c = \text{Jet}^n(F, G)_c^\sharp \otimes_{\text{Fix}(K, \sigma)} K$$

In particular, the dimension of $\text{Jet}^n(F, G)_c^\sharp$ as a $\text{Fix}(K, \sigma)$ -vector space is equal to $\dim_K \text{Jet}^n(F)_c$.

We obtain the following form of the canonical base property, which should be viewed as a generalisation of [30, Theorem 1.2] from ACFA to CCMA. Indeed, our proof is modelled on the ACFA case. Canonical bases in simple theories were introduced as hyperimaginary elements in [15]. Details of the various properties they enjoy can be found there. Because CCMA is supersimple, it follows by [5] that types over acl^{eq} -closed sets are amalgamation bases, and that the canonical base of an amalgamation base is in fact interdefinable with a (possibly infinite) tuple of imaginary elements.

Theorem 6.4 (CBP for finite-dimensional types). *Suppose (F, G) is a CCM- σ -variety defined over $B = \text{acl}(B)$, and $c \in (F, G)^\sharp$ is generic in F over B . Let*

$B_1 \supseteq B$ be an acl^{eq} -closed set of parameters. If $e = \text{Cb}(c/B_1)$ then $\text{tp}(e/Bc)$ is almost $\text{Fix}(K, \sigma)$ -internal.

Proof. Let $B'_1 \supseteq B$ be the intersection of B_1 with the real sorts of (\mathcal{A}', σ) . By geometric elimination of imaginaries (Fact 4.1), $B_1 \subseteq \text{acl}^{\text{eq}}(B'_1)$.

Let $F_1 = \text{loc}(c/B'_1)$ and let d be a canonical parameter for F_1 in CCM. We first claim that d and e are interalgebraic over B in CCMA. Let α be an automorphism of (\mathcal{A}', σ) that is the identity on Be . Then $\alpha(c) \in (F, G)^\sharp$ is generic in F over B also, and by properties of canonical bases $\text{tp}(\alpha(c)/\alpha(B_1))$ is *parallel* to $\text{tp}(c/B_1)$, where parallelism is the transitive closure of the relation of having a common nonforking extension. So $\text{tp}(\alpha(c)/\alpha(B'_1))$ is parallel to $\text{tp}(c/B'_1)$ also. It follows by Lemma 5.3 that $\text{loc}(\alpha(c)/\alpha(B'_1)) = \text{loc}(c/B'_1)$. That is, $\alpha(F_1) = F_1$, and so $\alpha(d) = d$. This shows that $d \in \text{dcl}(Be)$. On the other hand, Lemma 5.3 also implies that $c \downarrow_d B'_1$, and so $e = \text{Cb}(c/B_1) \in \text{acl}^{\text{eq}}(d)$.

So it suffices to show that $\text{tp}(d/Bc)$ is almost $\text{Fix}(K, \sigma)$ -internal. In fact, we will show it is $\text{Fix}(K, \sigma)$ -internal.

Let $n > 0$ be as given by Proposition 6.2 applied to compact complex varieties $X \subset S \times Z$ where $F_1 = X_a$ for $a \in S(\mathcal{A}')$. Let A be a $\text{Fix}(K, \sigma)$ -basis for $\text{Jet}^n(F, G)_c^\sharp$, chosen so that $A \downarrow_{Bc} d$. We show that $d \in \text{dcl}(B, c, A, \text{Fix}(K, \sigma))$.

Setting $G_1 = \text{loc}((c, \sigma(c))/B_1)$, we have that (F_1, G_1) is a CCM- σ -variety, and so $\text{Jet}^n(G_1)_{(c, \sigma(c))}$ induces on $\text{Jet}^n(F_1)_c$ the structure of a σ -module over (K, σ) . Now $\text{Jet}^n(G_1)_{(c, \sigma(c))}$ is a K -subspace of $\text{Jet}^n(G)_{(c, \sigma(c))}$, so that $\text{Jet}^n(F_1)_c$ is a σ -submodule of $\text{Jet}^n(F)_c$. Hence $\text{Jet}^n(F_1, G_1)_c^\sharp$ is a $\text{Fix}(K, \sigma)$ -subspace of $\text{Jet}^n(F, G)_c^\sharp$.

Let α be an automorphism of (\mathcal{A}', σ) that fixes the sets $B, c, A, \text{Fix}(K, \sigma)$ pointwise. Then α fixes all of $\text{Jet}^n(F, G)_c^\sharp$ pointwise, and hence all of $\text{Jet}^n(F_1, G_1)_c^\sharp$. But, $\text{Jet}^n(F_1)_c = \text{Jet}^n(F_1, G_1)_c^\sharp \otimes_{\text{Fix}(K, \sigma)} K$, so that α preserves $\text{Jet}^n(F_1)_c$ setwise. That is, $\text{Jet}^n(F_1)_c = \text{Jet}^n(F_1^\alpha)_c$. So by Proposition 6.2, $F_1 = F_1^\alpha$. As d is a canonical parameter for F_1 in CCM, we have that $\alpha(d) = d$, as desired. \square

As was first observed by Pillay [27], from such a canonical base property one can deduce a Zilber dichotomy statement. Our's will apply to finite-dimensional types; recall that for B inductive, $\text{tp}(c/B)$ is *finite-dimensional* if $\dim \text{loc}((c, \dots, \sigma^n c)/B)$ is bounded independently of n .

Corollary 6.5 (Zilber dichotomy for finite-dimensional types). *Suppose B is an acl -closed set and $p(x) \in S(B)$ is a finite-dimensional type of SU -rank one. Then either p is one-based or it is almost internal to $\text{Fix}(K, \sigma)$.*

Proof. We follow the argumentation of [26, Corollary 6.19].

Suppose $p(x)$ is not one-based. Then there exists a finite tuple c of realisations of $p(x)$ and a model $M \supseteq B$ such that $e := \text{Cb}(c/M) \notin \text{acl}(Bc)$. Note that $\text{tp}(c/B)$ is also finite-dimensional so that, possibly after replacing c by $(c, \sigma c, \dots, \sigma^n c)$ for some n , we may assume by Lemma 5.5 that there is a CCM- σ -variety (F, G) over B and that c is generic in F over B and contained in $(F, G)^\sharp$. Hence, by Theorem 6.4 (with $B_1 = M$), we have that $\text{tp}(e/Bc)$ is almost $\text{Fix}(K, \sigma)$ -internal. So we have $e \in \text{acl}(A, B, c, d)$ for some tuple d from $\text{Fix}(K, \sigma)$ and some $A \downarrow_{Bc} e$. It follows that $e \not\downarrow_{ABc} d$. Now e is in the definable closure of a finite set of realisations of $p(x)$; this is because e is in the definable closure of a finite part of a Morley sequence in $\text{tp}(c/M)$, and c is a tuple of realisations of $p(x)$. Since $SU(p) = 1$, we can find ABc -independent realisations a_1, \dots, a_m of $p(x)$ such that $e \in \text{acl}(Ba_1, \dots, a_m)$.

Hence $(a_1, \dots, a_m) \not\perp_{ABc} d$. But then for some $i < m$, $a_{i+1} \in \text{acl}(a_1 \dots a_i ABcd)$, witnessing that $p(x)$ is almost internal to $\text{Fix}(K, \sigma)$. \square

Remark 6.6. On the face of it the above corollary is only about real types, as per our conventions we do not work in $(\mathcal{A}', \sigma)^{\text{eq}}$. But because CCMA admits geometric elimination of imaginaries (Fact 4.1), and because both one-basedness and almost internality to $\text{Fix}(K, \sigma)$ are preserved by interalgebraicity, we get the Zilber dichotomy for all finite-dimensional SU -rank one types in $(\mathcal{A}', \sigma)^{\text{eq}}$ also.

The Zilber dichotomy result of Corollary 6.5 applies only to finite-dimensional minimal types. We give an example to show that not all minimal types are finite-dimensional. Whether a Zilber dichotomy holds for these types we leave as an open question; we only give an example of a trivial infinite-dimensional minimal type.

Example 6.7. Let X be a simply connected strongly minimal compact complex variety of dimension greater than 1 and with trivial automorphism group; for example, almost all generic K3 surfaces have these properties [18, Theorem 3.6].

By Theorem 4.8, (\mathcal{X}, σ) is purely stably embedded in (\mathcal{A}', σ) , where \mathcal{X} denotes the structure on $X(\mathcal{A}')$ induced by \mathcal{A}' , which by [24, Proposition 2.3] is just the structure of equality.

So the full induced structure on (\mathcal{X}, σ) is that of an infinite set with a generic permutation σ , and with distinguished σ -fixed points for the complex points $X(\mathcal{A})$.

So the type in CCMA of an aperiodic point of X ,

$$p := \{x \in X\} \cup \{x \neq \sigma^n x \mid n \in \mathbb{N}\},$$

is complete, minimal, and has trivial geometry. If $b \models p$, then $\text{loc}(b, \dots, \sigma^{n-1}b) = X^n$, so p has infinite dimension.

We may also note that the types of aperiodic points of X provide examples of trivial minimal types of finite dimension which are orthogonal to ACFA. Indeed, the discussion above shows that X is *fully orthogonal* to ACFA in the sense that every tuple from X is independent of every tuple from K over any parameters.

7. MINIMAL ONE-BASED TYPES

The Zilber dichotomy theorem, Corollary 6.5, tells us that there are no new non-one-based finite-dimensional minimal types in CCMA; they all come from ACFA and in fact from the fixed field in ACFA. In Example 6.7, we saw that there are new trivial minimal types, both of finite and of infinite dimension. There are also new finite-dimensional minimal types that are nontrivial and one-based, as the following example shows.

$\text{Fix}(\mathcal{A}', \sigma) \models \text{CCMA}$ sufficiently saturated.

Example 7.1. There are definable groups in (\mathcal{A}', σ) that are finite-dimensional, SU -rank one, one-based, and fully orthogonal to ACFA.

Proof. If X is any nonalgebraic simple complex torus, then $\text{Fix}(X(\mathcal{A}'), \sigma)$ will be such an example. In fact, we will describe precisely which definable subgroups of $X(\mathcal{A}')$ will have the desired properties.

First of all, it follows from Theorem 4.8 that X is a one-based stable group in CCMA. Indeed, it is a one-based stable group in CCM, and so if \mathcal{X}' denotes the CCM-structure on $X(\mathcal{A}')$, then (\mathcal{X}', σ) is a one-based stable group by [14], and

by 4.8 the latter is the full structure induced on $X(\mathcal{A}')$. Secondly, as X is fully orthogonal to \mathbb{P} in CCM it follows from the characterisation of independence in CCMA – see (3) of Section 4 – that X and \mathbb{P} remain fully orthogonal in CCMA. Hence, one-basedness and full orthogonality to ACFA will come for free; what we require is simply a description of all the finite-dimensional definable subgroups of $X(\mathcal{A}')$ of SU -rank one. Hrushovski’s arguments in [16, §4.1], which are written for simple abelian varieties in ACFA but work also for simple complex tori in CCMA, show that the definable subgroups of $X(\mathcal{A}')$ of SU -rank one are precisely those of the form $\ker(g)$ where g is an element of $\text{End}(X)[\sigma]$ and is left-irreducible in $\text{End}_{\mathbb{Q}}(X)[\sigma, \sigma^{-1}]$. Here, $\text{End}(X)$ denotes the (noncommutative unitary) ring of holomorphic homomorphisms from X to itself, and $\text{End}_{\mathbb{Q}}(X) := \text{End}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$. Note that $\ker(g)$ will be finite-dimensional. Taking $g = \sigma - 1$ yields the example of $\text{Fix}(X(\mathcal{A}'), \sigma)$. \square

The following proposition essentially says that these are the only nontrivial one-based minimal types of finite dimension that don’t come from ACFA.

Proposition 7.2. *Suppose $p(x) \in S(B)$ is a finite-dimensional minimal nontrivial one-based type in (\mathcal{A}', σ) that is orthogonal to the projective line. There exist a commutative simple nonalgebraic \mathcal{A}' -compact \mathcal{A}' -meromorphic group G and a quantifier-free definable subgroup H of SU -rank one such that $p(x)$ is nonorthogonal to (all) the generic types of H .*

Remark 7.3. (i) In the case when $B = \emptyset$ and the sort of $p(x)$ is of Kähler-type, the group G in the conclusion can be taken to be (the interpretation in \mathcal{A}' of a) simple nonalgebraic complex torus. In general we need a nonstandard analogue of complex tori.

(ii) We mean \mathcal{A}' -meromorphic group in the sense of [1, Definition 4.3]; it is the natural nonstandard analogue of meromorphic group. By the usual proof of the Weil-vdDries-Hrushovski theorem, every group interpretable in \mathcal{A}' can be endowed with the structure of an \mathcal{A}' -meromorphic group. By \mathcal{A}' -compact we mean (somewhat oddly) that as an \mathcal{A}' -manifold G is the image of a closed set in \mathcal{A}' under an \mathcal{A}' -holomorphic map (again the notions of \mathcal{A}' -manifold and \mathcal{A}' -holomorphic are the natural nonstandard ones, see [1, Definition 4.1]). An \mathcal{A}' -compact group appears as the generic fibre of a definable family of meromorphic groups that are individually definably (though not necessarily uniformly) isomorphic to complex tori. In particular, \mathcal{A}' -compact groups are commutative. One would like to think of them as “nonstandard complex tori”, except that outside of the essentially saturated context there are some subtleties (and open questions) about doing so, see the discussion in the Introduction to [1].

Sketch of proof of 7.2. This is just the analogue for CCMA of the corresponding fact about minimal nontrivial one-based types in ACFA which is due to Chatzidakis-Hrushovski [7, Theorem 5.12]. Indeed, their proof goes through in this setting. We sketch some details.

By finite dimensionality, replacing $p(x)$ with something interdefinable with it, we can assume that if $a \models p(x)$ then $\sigma(a) \in \text{acl}^-(Ba)$. It follows, using geometric elimination of imaginaries for CCMA, that nontriviality of $p(x)$ gives rise to a group configuration in CCM made up of elements that are CCM-interalgebraic with realisations of $p(x)$. The group configuration theorem, along with the observation

that a stable connected group is abelian if $a * b * c \in \text{acl}(b, \text{acl}(a, c) \cap \text{acl}(b, a * b * c))$ for generic independent a, b, c , yields an abelian simple \mathcal{A}' -meromorphic group G and an SU -rank one type whose quantifier-free stabiliser H is a definable subgroup of G with generic types nonorthogonal to p .

Note that as p is assumed to be orthogonal to the projective line, G must be nonalgebraic. It remains to see that G is \mathcal{A}' -compact. Meromorphic groups in \mathcal{A} were given a Chevalley-type characterisation in [29], this was extended to strongly minimal \mathcal{A}' -meromorphic groups in [1], and then to all \mathcal{A}' -meromorphic groups in [34]. That characterisation says that G is the extension of an \mathcal{A}' -compact group by a linear algebraic group. Nonalgebraicity and simplicity thus force G to be \mathcal{A}' -compact. \square

APPENDIX A. COMPLEX ANALYTIC JET SPACES

In this appendix we review a particular construction of higher order tangent spaces in complex analytic geometry. The exposition given here is based on the third author's (more detailed) unpublished survey [20] of related constructions and their use in model theory.

A.1. Linear spaces. Suppose X is a complex variety. A *linear space over X* is a complex variety over X , $L \rightarrow X$, whose fibres are uniformly equipped with complex vector space structure. That is, there are holomorphic maps for addition $+: L \times_X L \rightarrow L$, scalar multiplication $\lambda: \mathbb{C} \times L \rightarrow L$, and zero section $z: X \rightarrow L$, all over X , satisfying the usual axioms. For example, $X \times \mathbb{C}^n$ is a linear space over X , it is the *trivial* linear space of rank n .

There is a natural notion of *homomorphism* between linear spaces L and L' over X , the set of which is denoted by $\text{Hom}_X(L, L')$ and has canonically the structure of an $\mathcal{O}_X(X)$ -module. See [10, §1.4] for details.

Given a coherent analytic sheaf \mathcal{F} on X , there is a *linear space over X associated to \mathcal{F}* , denoted by $L(\mathcal{F}) \rightarrow X$ which has the property that for all open U in X

$$\mathcal{F}(U) \cong \text{Hom}_U(L(\mathcal{F})_U, U \times \mathbb{C})$$

as $\mathcal{O}_X(U)$ -modules. Locally $L(\mathcal{F}) \rightarrow X$ can be described as follows: let $U \subset X$ be a (euclidean) open subset for which there exists a resolution of \mathcal{F}_U as

$$\mathcal{O}_U^p \xrightarrow{\alpha} \mathcal{O}_U^q \longrightarrow \mathcal{F}_U \longrightarrow 0$$

Represent α as a $q \times p$ matrix $M = (m_{ij}(x))$ with entries in \mathcal{O}_U . Then $L(\mathcal{F})_U$ is the subspace of the trivial linear space $U \times \mathbb{C}^q$ defined by the equations

$$m_{1i}(x)y_1 + \cdots + m_{qi}(x)y_q = 0$$

for $i = 1, \dots, p$.

Fact A.1 (Section 1.8 of [10]). *If \mathcal{F} is a coherent analytic sheaf on X and $x \in X$ then there is a canonical isomorphism $L(\mathcal{F})_x \cong \text{Hom}_{\mathbb{C}}(\mathcal{F}_x \otimes_{\mathcal{O}_{X,x}} \mathbb{C}, \mathbb{C})$*

What about definability of these objects in CCM? We can relativise the usual compactification of \mathbb{C}^n as $\mathbb{P}_n(\mathbb{C})$ as follows. If in the above local construction of $L(\mathcal{F})$ we treat $(y_1 : \cdots : y_q)$ as homogeneous co-ordinates, then the equations cut out a complex analytic subset of $U \times \mathbb{P}_{q-1}(\mathbb{C})$, and one obtains by gluing what is called the *projective linear space associated to \mathcal{F}* , denoted by $\mathbb{P}(\mathcal{F}) \rightarrow X$. Applying

the construction to the sheaf $\mathcal{F} \times \mathcal{O}_X$ we see that the linear space $L(\mathcal{F}) \rightarrow X$ embeds as a Zariski open subset of the projective linear space $\mathbb{P}(\mathcal{F} \times \mathcal{O}_X) \rightarrow X$ in such a way that linear structure extends meromorphically to the projective linear space. In particular, if X is compact then $L(\mathcal{F}) \rightarrow X$, together with the uniform vector space structure on the fibres, is definable in CCM.

A.2. Relative differentials. Suppose $\pi : X \rightarrow S$ is a holomorphic map of complex varieties and consider the diagonal embedding $d : X \rightarrow X \times_S X$. Recall that the inverse image sheaf $d^{-1}\mathcal{O}_{X \times_S X}$ is by definition the sheaf of rings on X which assigns to any open set U the direct limit of the rings $\mathcal{O}_{X \times_S X}(V)$ as V ranges among open neighbourhoods of $d(U)$ in $X \times_S X$. There is an induced homomorphism of sheaves $d^{-1}\mathcal{O}_{X \times_S X} \rightarrow \mathcal{O}_X$ which is surjective because d is a closed immersion. Let \mathcal{I} be the (ideal sheaf) kernel of this homomorphism. The *sheaf of relative n -differentials* is the sheaf $\underline{\Omega}_{X/S}^n := \mathcal{I}/\mathcal{I}^{n+1}$, viewed as a sheaf of \mathcal{O}_X -modules using the first co-ordinate projection.

Fact A.2 (Corollaire 2.7 of [13]). *If $\pi : X \rightarrow S$ is a holomorphic map of complex varieties and $x \in X$ then there is a canonical isomorphism*

$$\underline{\Omega}_{X/S,x}^n \otimes_{\mathcal{O}_{X,x}} \mathbb{C} \cong \mathfrak{m}_{X_{\pi(x)},x} / \mathfrak{m}_{X_{\pi(x)},x}^{n+1}$$

where $\mathfrak{m}_{X_{\pi(x)},x}$ is the maximal ideal of the local ring of the fibre $X_{\pi(x)}$ at the point x .

A.3. Relative jet spaces. The word “jet” is used in various ways in the literature. The reader should be warned that our jet spaces are different from those of Buium, which among model theorists usually go by the term “prolongation spaces”.

Given a point y on a complex variety Y , by the *n th jet space of Y at y* we mean the finite dimensional complex vector space $\text{Jet}^n(Y)_y := \text{Hom}_{\mathbb{C}}(\mathfrak{m}_{Y,y} / \mathfrak{m}_{Y,y}^{n+1}, \mathbb{C})$. We view this as a kind of higher order tangent space.

Lemma A.3. *Suppose A and B are irreducible complex analytic subsets of a complex variety Y and $y \in A \cap B$. If $\text{Jet}^n(A)_y = \text{Jet}^n(B)_y$ for all $n \geq 1$, then $A = B$.*

Proof. Note that in the statement we are viewing $\text{Jet}^n(A)_y$ and $\text{Jet}^n(B)_y$ canonically as subspaces of $\text{Jet}^n(Y)_y$. We will show that $A \subseteq B$ and conclude by symmetry that $A = B$. As $A \cap B$ is an analytic subset of A , and as A is irreducible, it suffices to show that in some non-empty (euclidean) open subset of Y containing y , say U , $(A \cap U) \subseteq (B \cap U)$. This in turn reduces to showing that the defining ideal of the germ of B at y is contained in that of the germ of A .

To that end, suppose $f \in \mathcal{O}_{Y,y}$ is a germ of a holomorphic function at y which vanishes on the germ of B . It follows that every linear functional in $\text{Jet}^n(B)_y$ vanishes on f , and so as $\text{Jet}^n(A)_y = \text{Jet}^n(B)_y$, the same is true of all functionals in $\text{Jet}^n(A)_y$. This implies that the image of f in $\mathcal{O}_{A,y}$ is contained in $\mathfrak{m}_{A,y}^{n+1}$ for all n . As $\bigcap_n \mathfrak{m}_{A,y}^{n+1} = 0$, we have that the image of f in $\mathcal{O}_{A,y}$ is zero. \square

What is particularly important for us is that when $\pi : X \rightarrow S$ is a holomorphic map of complex varieties, the n th jet spaces of the fibres of π vary uniformly.

Definition A.4. Suppose $\pi : X \rightarrow S$ is a holomorphic map of complex varieties. By the *n th jet space of X relative to S* , which we will denote by $\text{Jet}^n(X/S) \rightarrow X$, we will mean the linear space over X associated to $\underline{\Omega}_{X/S}^n$.

A consequence of Facts A.1 and A.2 is that the fibres of $\text{Jet}^n(X/S) \rightarrow X$ are

$$\text{Jet}^n(X/S)_x = \text{Jet}^n(X_{\pi(x)})_x = \text{Hom}_{\mathbb{C}}(\mathfrak{m}_{X_{\pi(x)},x}/\mathfrak{m}_{X_{\pi(x)},x}^{n+1}, \mathbb{C})$$

for all $x \in X$. Moreover, it follows from our discussion of linear spaces and their compactification in projective linear spaces that when X and S are compact, $\text{Jet}^n(X/S) \rightarrow X$ as well as the vector space structure on the fibres is (uniformly) definable in CCM.

REFERENCES

- [1] M. Aschenbrenner, R. Moosa, and T. Scanlon. Strongly minimal groups in the theory of compact complex spaces. *Journal of Symbolic Logic*, 71(2):529–552, 2006.
- [2] W. Barth, C. Peters, and A. Van de Ven. *Compact complex surfaces*. Springer-Verlag, Berlin, 1984.
- [3] M. Bays, M. Gavrilovich, and M. Hils. Some definability results in abstract Kummer theory. *International Mathematics Research Notices*, 2013. doi:10.1093/imrn/rnt057.
- [4] C. Bănică. Le lieu réduit et le lieu normal d’un morphisme. In *Romanian-Finnish Seminar on Complex Analysis (Proc. Bucharest, 1976)*, volume 743 of *Lecture Notes in Math*. Springer, Berlin, 1979.
- [5] S. Buechler, A. Pillay, and F. Wagner. Supersimple theories. *Journal of the American Mathematical Society*, 14(1):109–124, 2001.
- [6] R. Bustamante. *Théorie de modèles des corps différentiellement clos avec un automorphisme générique*. PhD thesis, Université Paris 7, 2005.
- [7] Z. Chatzidakis and E. Hrushovski. Model theory of difference fields. *Transactions of the American Mathematical Society*, 351(8):2997–3071, 1999.
- [8] Z. Chatzidakis and E. Hrushovski. Difference fields and descent in algebraic dynamics. I. *J. Inst. Math. Jussieu*, 7(4):653–686, 2008.
- [9] Z. Chatzidakis and A. Pillay. Generic structures and simple theories. *Ann. Pure Appl. Logic*, 95(1-3):71–92, 1998.
- [10] G. Fischer. *Complex Analytic Geometry*. Springer-Verlag, Berlin, 1976.
- [11] H. Grauert, T. Peternell, and R. Remmert, editors. *Several Complex Variables VII*, volume 74 of *Encyclopedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1994.
- [12] P. Griffiths and J. Harris. *Principles of algebraic geometry*. John Wiley & Sons, Inc., New York, 1994. Reprint of the 1978 original.
- [13] A. Grothendieck. Techniques de construction en géométrie analytique VII. Étude locale des morphisme: éléments de calcul infinitésimal. *Séminaire Henri Cartan*, 13(14), 1960/61.
- [14] H.B. Gute and K.K. Reuter. The last word on elimination of quantifiers in modules. *Journal of Symbolic Logic*, 55(2):670–673, 1990.
- [15] B. Hart, B. Kim, and A. Pillay. Coordinatisation and canonical bases in simple theories. *The Journal of Symbolic Logic*, 65(1):293–309, 2000.
- [16] E. Hrushovski. The Manin-Mumford conjecture and the model theory of difference fields. *Annals of Pure and Applied Logic*, 112(1):43–115, 2001.
- [17] E. Hrushovski. Groupoids, imaginaries and internal covers. Preprint, arXiv:math/0603413v3, 2011.
- [18] E. Macrì and P. Stellari. Automorphisms and autoequivalences of generic analytic $K3$ surfaces. *J. Geom. Phys.*, 58(1):133–164, 2008.
- [19] A. Medvedev and T. Scanlon. Invariant varieties for polynomial dynamical systems. *Ann. of Math. (2)*, 179(1):81–177, 2014.
- [20] R. Moosa. Jet spaces in complex analytic geometry: An exposition. E-print available at <http://arxiv.org/abs/math.LO/0405563>, 2003.
- [21] R. Moosa. A nonstandard Riemann existence theorem. *Transactions of the American Mathematical Society*, 356(5):1781–1797, 2004.
- [22] R. Moosa. The model theory of compact complex spaces. In *Logic Colloquium ’01*, volume 20 of *Lect. Notes Log.*, pages 317–349. Assoc. Symbol. Logic, Urbana, IL, 2005.
- [23] R. Moosa and A. Pillay. On canonical bases and internality criteria. *Illinois Journal of Mathematics*, 52(3):901–917, 2008.

- [24] R. Moosa and A. Pillay. \aleph_0 -categorical strongly minimal compact complex manifolds. *Proc. Amer. Math. Soc.*, 140(5):1785–1801, 2012.
- [25] R. Moosa and T. Scanlon. Jet and prolongation spaces. *Journal de l’Institut de Mathématiques de Jussieu*, 9(2):391–430, 2010.
- [26] R. Moosa and T. Scanlon. Model theory of fields with free operators in characteristic zero. Preprint, 2013.
- [27] A. Pillay. Model-theoretic consequences of a theorem of Campana and Fujiki. *Fundamenta Mathematicae*, 174(2):187–192, 2002.
- [28] A. Pillay. Remarks on algebraic D -varieties and the model theory of differential fields. In *Logic in Tehran*, volume 26 of *Lect. Notes Log.*, pages 256–269. Assoc. Symbol. Logic, La Jolla, CA, 2006.
- [29] A. Pillay and T. Scanlon. Meromorphic groups. *Transactions of the American Mathematical Society*, 355(10):3843–3859, 2003.
- [30] A. Pillay and M. Ziegler. Jet spaces of varieties over differential and difference fields. *Selecta Math. (N.S.)*, 9(4):579–599, 2003.
- [31] Anand Pillay. Some model theory of compact complex spaces. In *Hilbert’s tenth problem: relations with arithmetic and algebraic geometry (Ghent, 1999)*, volume 270 of *Contemp. Math.*, pages 323–338. Amer. Math. Soc., Providence, RI, 2000.
- [32] J. Le Potier. Simple connexité des surfaces $K3$. *Astérisque*, (126):79–89, 1985. Geometry of $K3$ surfaces: moduli and periods (Palaiseau, 1981/1982).
- [33] W. Sawin. Answer to a mathoverflow question, <http://mathoverflow.net/questions/96680/etale-covers-of-line-bundles-on-an-abelian-variety>, 2012.
- [34] T. Scanlon. Nonstandard meromorphic groups. In *Proceedings of the 12th Workshop on Logic, Language, Information and Computation (WoLLIC 2005)*, volume 143 of *Electron. Notes Theor. Comput. Sci.*, pages 185–196 (electronic). Elsevier, Amsterdam, 2006.

MARTIN BAYS, MCMASTER UNIVERSITY, DEPARTMENT OF MATHEMATICS AND STATISTICS, 1280 MAIN STREET WEST, HAMILTON, ONTARIO L8S 4L8, CANADA
E-mail address: `mbays@math.mcmaster.ca`

MARTIN HILS, UNIV PARIS DIDEROT, SORBONNE PARIS CITÉ, INSTITUT DE MATHÉMATIQUES DE JUSSIEU–PARIS RIVE GAUCHE, UMR 7586, CNRS, SORBONNE UNIVERSITÉS, UPMC UNIV PARIS 06, F-75013, PARIS, FRANCE
E-mail address: `hils@math.univ-paris-diderot.fr`

RAHIM MOOSA, UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, 200 UNIVERSITY AVENUE WEST, WATERLOO, ONTARIO N2L 3G1, CANADA
E-mail address: `rmoosa@uwaterloo.ca`