

Name:

ID:

---

PMath 330

Assignment 8

Mark \_\_\_\_\_

The following output was generated by S.O.B.B:

```


14  (x · y) + (x · z)      =  x · (y + z)
100 (x · y) + (z · x)      =  x · (y + z)
127 x · (y · (z + x))      =  y · x
628 x · (y + (z · (u + x))) =  x · (y + z) [para_from 127.1.1, 14.1.1.2; demod 100 ;flip.1]
```

Give the details of how S.O.B.B. found equation 628. (Follow the format in the supplementary page on Critical Pairs and Equational Theorem Proving.)

DETAILS:

Structures


Fill in the table for the following directed graph:



	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
<i>a</i>				
<i>b</i>				
<i>c</i>				
<i>d</i>				

Draw the directed graph for the following table:

	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
<i>a</i>	1	0	1	0
<i>b</i>	0	1	1	0
<i>c</i>	1	0	0	1
<i>d</i>	1	0	1	0



Herbrand Universe

Suppose that a first-order language has only a constant symbol  $a$ , a binary function symbol  $f$ , and a binary relation symbol  $r$ .

List the elements of the second level  $S_2$  of the Herbrand universe:

$S_2$  = \_\_\_\_\_

List all of the ground clauses that one can make from the symmetry clause  $\{\neg rxy, ryx\}$  using just the portion of the Herbrand universe called  $S_1$ .

ANSWER:

**Clauses**

Let **S** be the structure on  $\{0, 1, 2\}$  with a unary operation defined by  $fx = x + 1 \pmod 3$ . The relation  $<$  is the usual, that is,  $0 < 1 < 2$ . Given the literals

$L_1 = \neg(fx < y) \qquad L_2 = x < fy$

determine all pairs  $(a, b)$  that satisfy the clause  $C = \{L_1, L_2\}$  in **S** by filling in the following:

$x$	$y$	$fx$	$fy$	$fx < y$	$\neg(fx < y)$	$x < fy$	$C$
0	0						
0	1						
0	2						
1	0						
1	1						
1	2						
2	0						
2	1						
2	2						

Does **S** satisfy the clause **C**? \_\_\_\_\_

**Opp-unification:** Suppose our first-order language has a unary relation symbol  $r$  and a binary operation symbol  $g$ . Show that the two clauses  $\{rgxgyx, rggyzw\}$  and  $\{\neg rgxggyzw\}$  are opp-unifiable, and find the most general opp-unifier  $(\sigma_1, \sigma_2)$ . [Change  $x, y, z, w$  in the second clause to  $X, Y, Z, W$ .]







Thus  $\sigma_1 = \left( \begin{array}{l} x \leftarrow \\ y \leftarrow \\ z \leftarrow \\ w \leftarrow \end{array} \right)$  and  $\sigma_2 = \left( \begin{array}{l} x \leftarrow \\ y \leftarrow \\ z \leftarrow \\ w \leftarrow \end{array} \right)$

Resolution Theorem Proving

For  $f$  a unary function symbol and  $r$  a binary relation symbol fill in the reasons for the following resolution derivation:

1.	$\{ rxfx \}$	<u>given</u>
2.	$\{ \neg rf00 \}$	<u>given</u>
3.	$\{ \neg rxy, ryx \}$	<u>given</u>
4.	$\{ rfx \}$	
5.	$\{ \}$	

The following uses ground clauses from the given clauses 1–3 of the previous problem. Identify the source of each ground clause (e.g., clause 3) that is a ground instance of one of the above given clauses, and give reasons (i.e., line numbers) for the resolution steps:

1.	$\{ r0f0 \}$	
2.	$\{ \neg r0f0, rf00 \}$	
3.	$\{ \neg rf00 \}$	
4.	$\{ \neg r0f0 \}$	
5.	$\{ \}$	