

The following output was generated by S.O.B.B:

```

14  (x · y) + (x · z)      =  x · (y + z)
100 (x · y) + (z · x)      =  x · (y + z)
127 x · (y · (z + x))      =  y · x
628 x · (y + (z · (u + x))) =  x · (y + z) [para_from 127.1.1, 14.1.1.2; demod 100 ;flip.1]
```

Give the details of how S.O.B.B. found equation 628. (Follow the format in the supplementary page on Critical Pairs and Equational Theorem Proving.)

DETAILS: We want to find the critical pair equation for the following pair of equations by unifying the boxed terms:

$$\begin{array}{lcl}
 127 & \boxed{x \cdot (y \cdot (z + x))} & = y \cdot x \\
 14 & (x \cdot y) + \boxed{(x \cdot z)} & = x \cdot (y + z)
 \end{array}$$

The first thing to do is to make the variables disjoint:

$$\begin{array}{lcl}
 127 & \boxed{x \cdot (y \cdot (z + x))} & = y \cdot x \\
 14 & (X \cdot Y) + \boxed{(X \cdot Z)} & = X \cdot (Y + Z)
 \end{array}$$

The most general unifier of the boxed terms is

$$\mu = \left(\begin{array}{l} X \leftarrow x \\ Z \leftarrow y \cdot (z + x) \end{array} \right)$$

Applying this to 127 and 14 gives

$$\begin{array}{lcl}
 127 & \boxed{x \cdot (y \cdot (z + x))} & = y \cdot x \\
 14 & (x \cdot Y) + \boxed{(x \cdot (y \cdot (z + x)))} & = x \cdot (Y + (y \cdot (z + x)))
 \end{array}$$

From this we have the Critical Pair Equation:

$$(**) \quad (x \cdot Y) + (y \cdot x) = x \cdot (Y + (y \cdot (z + x)))$$

By applying 100 to the left side of (**) we obtain:


$$x \cdot (Y + y) = x \cdot (Y + (y \cdot (z + x)))$$

Flip, and change the variables, to get

$$x \cdot (y + (z \cdot (u + x))) = x \cdot (y + z)$$

Structures


Fill in the table for the following directed graph:



	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
<i>a</i>	1	1	0	0
<i>b</i>	1	0	0	1
<i>c</i>	1	1	1	0
<i>d</i>	1	0	1	0

Draw the directed graph for the following table:

	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
<i>a</i>	1	0	1	0
<i>b</i>	0	1	1	0
<i>c</i>	1	0	0	1
<i>d</i>	1	0	1	0



Herbrand Universe

Suppose that a first-order language has only a constant symbol *a*, a binary function symbol *f*, and a binary relation symbol *r*.

List the elements of the second level S_2 of the Herbrand universe:

$$S_2 = \underline{\{a, faa, ffaaa, fafaa, ffaafaa\}}$$

List all of the ground clauses that one can make from the symmetry clause $\{\neg rxy, ryx\}$ using just the portion of the Herbrand universe called S_1 .

ANSWER:

$$\begin{aligned} &\{\neg raa, raa\} \\ &\{\neg rafaa, rfaaa\} \\ &\{\neg rfaaa, rafaa\} \\ &\{\neg rfafaa, rfaafaa\} \end{aligned}$$

Clauses

Let \mathbf{S} be the structure on $\{0, 1, 2\}$ with a unary operation defined by $fx = x + 1 \pmod{3}$. The relation $<$ is the usual, that is, $0 < 1 < 2$. Given the literals

$$L_1 \text{ is } \neg(fx < y) \quad L_2 \text{ is } x < fy$$

determine all pairs (a, b) that satisfy the clause $C = \{L_1, L_2\}$ in \mathbf{S} by filling in the following:

x	y	fx	fy	$fx < y$	$\neg(fx < y)$	$x < fy$	C
0	0	1	1	0	1	1	1
0	1	1	2	0	1	1	1
0	2	1	0	1	0	0	0
1	0	2	1	0	1	0	1
1	1	2	2	0	1	1	1
1	2	2	0	0	1	0	1
2	0	0	1	0	1	0	1
2	1	0	2	1	0	0	0
2	2	0	0	1	0	0	0

Does \mathbf{S} satisfy the clause C ? NO

Opp-unification: Suppose our first-order language has a unary relation symbol r and a binary operation symbol g . Show that the two clauses $\{rgxgyx, rggyzw\}$ and $\{\neg rgxggysz\}$ are opp-unifiable, and find the most general opp-unifier (σ_1, σ_2) . [Change x, y, z, w in the second clause to X, Y, Z, W .]

r	g	x	g	y	x											
r	g	g	y	z	w											
r	g	X	g	g	Y	Z	W									

↓ $(x \leftarrow X)$

r	g	X	g	y	X											
r	g	g	y	z	w											
r	g	X	g	g	Y	Z	W									

↓ $(X \leftarrow gyz)$

r	g	g	y	z	g	y	g	y	z							
r	g	g	y	z	w											
r	g	g	y	z	g	g	Y	Z	W							

↓ $(w \leftarrow ggyyz)$

r	g	g	y	z	g	y	g	y	z							
r	g	g	y	z	g	y	g	y	z							
r	g	g	y	z	g	g	Y	Z	W							

↓ $(y \leftarrow gYZ)$

r	g	g	g	Y	Z	z	g	g	Y	Z	g	g	Y	Z	z	
r	g	g	g	Y	Z	z	g	g	Y	Z	g	g	Y	Z	z	
r	g	g	g	Y	Z	z	g	g	Y	Z	W					

↓ $(W \leftarrow ggYZz)$

r	g	g	g	Y	Z	z	g	g	Y	Z	g	g	Y	Z	z	
r	g	g	g	Y	Z	z	g	g	Y	Z	g	g	Y	Z	z	
r	g	g	g	Y	Z	z	g	g	Y	Z	g	g	Y	Z	z	

so $x \leftarrow ggYZz$, $w \leftarrow ggYZggYZz$, $X \leftarrow ggYZz$, $y \leftarrow gYZ$, $W \leftarrow ggYZz$.

Thus, with a, b, c being *any* three distinct variables,

$$\sigma_1 = \begin{pmatrix} x \leftarrow ggabc \\ y \leftarrow gab \\ z \leftarrow c \\ w \leftarrow ggabggabc \end{pmatrix} \quad \text{and} \quad \sigma_2 = \begin{pmatrix} x \leftarrow ggabc \\ y \leftarrow a \\ z \leftarrow b \\ w \leftarrow ggabc \end{pmatrix}$$

Resolution Theorem Proving

For f a unary function symbol and r a binary relation symbol fill in the reasons for the following resolution derivation:

1. $\{rxfx\}$	<u>given</u>
2. $\{\neg rf00\}$	<u>given</u>
3. $\{\neg rxy, ryx\}$	<u>given</u>
4. $\{rfxx\}$	<u>Res 1, 3</u>
5. $\{ \}$	<u>Res 2, 4</u>

The following uses ground clauses from the given clauses 1–3 of the previous problem. Identify the source of each ground clause (e.g., clause 3) that is a ground instance of one of the above given clauses, and give reasons (i.e., line numbers) for the resolution steps:

1. $\{r0f0\}$	<u>Clause 1</u>
2. $\{\neg r0f0, rf00\}$	<u>Clause 3</u>
3. $\{\neg rf00\}$	<u>Clause 2</u>
4. $\{\neg r0f0\}$	<u>Res 2, 3</u>
5. $\{ \}$	<u>Res 1, 4</u>