

1.	4.
2.	5.
3.	6.

Name _____
ID _____

Total = _____/90

PMath 330

MIDTERM

Friday, October 27, 2000

There are 4 pages, with 6 problems.

Problem 1

**12
marks**

Give the **mood** and **figure** of the syllogism to the right, and fill in its Venn diagram (using Modern Standards, as in the text). State whether the syllogism is valid under **Modern Standards** as well as under **Aristotelian Standards**.


No M is S
All P is M

Some S is not P.

Mood: AEO Figure: 4

Valid (Modern): NO

Valid (Aristotelian): YES



Problem 2

**13
marks**

Apply unit resolution to determine if the given set of clauses is satisfiable:

1. $\{P\}$	<u>given</u>	6. $\{Q, \neg R\}$	<u>1, 3</u>
2. $\{P, \neg Q, \neg R\}$	<u>given</u>	7. $\{\neg Q, R\}$	<u>1, 5</u>
3. $\{\neg P, Q, \neg R\}$	<u>given</u>	8. $\{P, \neg R\}$	<u>2, 4</u>
4. $\{Q\}$	<u>given</u>	9. $\{\neg P, R\}$	<u>4, 5</u>
5. $\{\neg P, \neg Q, R\}$	<u>given</u>	10. $\{R\}$	<u>1, 9</u>
		11. $\{P, \neg Q\}$	<u>2, 10</u>
		12. $\{\neg P, Q\}$	<u>3, 10</u>

ANSWER: (Reasons) YES. Cannot derive the empty clause.

Problem 3

**10
marks**

Given the following combined truth table for five formulas

	P	Q	R	F_1	F_2	F_3	F_4	F_5			
1.	1	1	1	0	1	1	1	0			
2.	1	1	0	1	0	1	1	1			
3.	1	0	1	0	1	0	1	1			
4.	1	0	0	1	1	0	1	0			
5.	0	1	1	1	1	1	1	0			
6.	0	1	0	1	0	1	1	1			
7.	0	0	1	0	0	0	0	1			
8.	0	0	0	1	0	0	0	0			

(a) What is the Conjunctive Normal Form of $F_2 \rightarrow F_3$?

$$(\neg P \vee Q \vee \neg R) \wedge (\neg P \vee Q \vee R)$$

(b) Is the set $\{F_1 \rightarrow F_2, F_2 \rightarrow F_3, F_3 \rightarrow F_1\}$ satisfiable? (Why?)

YES, by line 7.

(c) Is the argument $F_2, F_3, F_4, F_5 \therefore F_1$ valid? (Why?)

YES, as the premisses are not satisfiable.

Problem 4

**15
marks**

Given that $F(X) = AX \cup BX'$, where A and B are subsets of a given universe U , prove: there is an $X \subseteq U$ such that $F(X) = 0$ if and only if $AB = 0$.

PROOF:

First let us suppose that $AX \cup BX' = 0$, for some X . Then one has $AX = 0$ and $BX' = 0$. Thus $A \subseteq X'$ and $B \subseteq X$. But then $AB \subseteq X'X = 0$, so $AB = 0$. This proves one direction.


Next let us suppose that $AB = 0$. Then $F(B) = AB \cup BB' = 0$.

Problem 5

**20
marks**

You are given the premisses $A'B = 0$, $CD' = 0$, and $AD = 0$.

(a) Use Lewis Carroll's tree method to show that $BC = 0$ is a valid conclusion.



(b) Use Boole's elimination method to show that $BC = 0$ is the most general conclusion for the classes B and C .

SOLUTION:

The premisses are equivalent to the single equation $E(A, B, C, D) = 0$ where

$$E(A, B, C, D) = A'B \cup CD' \cup AD.$$

The most general conclusion for the classes B and C is $F(B, C) = 0$ where

$$\begin{aligned}
 F(B, C) &= E(1, B, C, 1)E(1, B, C, 0)E(0, B, C, 1)E(0, B, C, 0) \\
 &= (1)(C)(B)(B \cup C) \\
 &= BC.
 \end{aligned}$$

Thus $BC = 0$ is the most general conclusion for the classes B and C .

Problem 6

20

(a) Assuming just that the set of standard connectives (including the standard constants) is adequate, show that $\mathcal{C} = \{\rightarrow, 0\}$ is adequate. **marks**

SOLUTION:

$$\begin{aligned} 1 &\sim 0 \rightarrow 0 \\ \neg P &\sim P \rightarrow 0 \\ P \vee Q &\sim (P \rightarrow 0) \rightarrow Q \\ P \wedge Q &\sim ((P \rightarrow (Q \rightarrow 0)) \rightarrow 0) \\ P \leftrightarrow Q &\sim (P \rightarrow (Q \rightarrow 0)) \rightarrow (((P \rightarrow 0) \rightarrow Q) \rightarrow 0) \end{aligned}$$

(b) Give the truth table for the Sheffer stroke $|$. Then use the fact that \mathcal{C} in (a) is adequate to show that the Sheffer stroke is adequate.

SOLUTION:

P	Q	$P Q$
1	1	0
1	0	1
0	1	1
0	0	1

Since

$$\begin{aligned} 0 &\sim (P|(P|P))|(P|(P|P)) \\ P \rightarrow Q &\sim P|(P|Q) \end{aligned}$$

and \mathcal{C} is adequate, it follows that the Sheffer stroke is adequate.