


Parallels between
Additive Number Systems
and
Multiplicative Number Systems

Stanley Burris

Pure Mathematics Number Theory Seminar

November, 2002

Parallels




n	— — — — —	$\log n$
local	— — — — —	global
RT	— — — — —	RV

Review

Power Series

$$\mathbf{A}(z) = \sum_{n \geq 0} a(n)z^n$$

Radius of Convergence




Cauchy Formula

$$a(n) = \frac{1}{2\pi i} \int_{\mathcal{C}} \mathbf{A}(z) \frac{dz}{z^{n+1}}$$

Dirichlet Series

$$\mathbf{A}(s) = \sum_{n \geq 1} a(n)n^{-s}$$

Abscissa of Convergence



Perron Formula

$$A(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \mathbf{A}(s) \frac{x^s}{s} ds$$

if x is not an integer

Number Systems

Additive	Multiplicative
$(A, P, +, 0, \ \)$	$(A, P, \cdot, 1, \ \)$
A = the set of “numbers”	
P = the set of indecomposable “numbers”	
$\ \ $ is additive	$\ \ $ is multiplicative

Basic Requirements:

- only finitely many numbers have a given norm
- each number is uniquely decomposable as a sum of indecomposables.

The **Fundamental Identity** of a number system:

Additive Case

$$\begin{aligned} \mathbf{A}(x) &:= \sum_{n \geq 0} a(n)x^n = \prod_{n \geq 1} (1 - x^n)^{-p(n)} \\ &= \exp\left(\sum_{m \geq 1} \mathbf{P}(x^m)/m\right) \end{aligned}$$

where

$$\mathbf{P}(x) := \sum_{n \geq 1} p(n)x^n.$$

Multiplicative Case

$$\begin{aligned} \mathbf{A}(x) &:= \sum_{n \geq 1} a(n)n^{-x} = \prod_{n \geq 2} (1 - n^{-x})^{-p(n)} \\ &= \exp\left(\sum_{m \geq 1} \mathbf{P}(mx)/m\right) \end{aligned}$$

where

$$\mathbf{P}(x) := \sum_{n \geq 2} p(n)n^{-x}.$$

Radius/Abscissa of convergence

We have

$$0 \leq \rho \leq 1$$

$$0 \leq \alpha \leq \infty$$

$\rho = 0$ fast growth

$\alpha = \infty$ fast growth

$\rho > 0$ slow growth

$\alpha < \infty$ slow growth

Generating Functions of subsets of A

For $B \subseteq A$

Additive

Multiplicative

$$\mathbf{B}(x) = \sum_{n \geq 0} b(n)x^n$$

$$\mathbf{B}(x) = \sum_{n \geq 1} b(n)n^{-x}$$

where $b(n)$ counts the number of elements of B of size n .

Finitely Generated Number Systems show parallel behaviour

Let $r = \sum p(n)$.

Additive

$$a(n) \sim \left(\prod m^{-p(m)} \right) \cdot \frac{n^{r-1}}{(r-1)!}$$

Multiplicative

$$A(n) \sim \left(\prod (\log m)^{-p(m)} \right) \cdot \frac{(\log n)^r}{r!}$$

Another Parallel Result

**Knopfmacher₂ +
Warlimont**

$$p(n) = a\beta^n + O(\gamma^n)$$

implies

$$a(n) \sim C\beta^n \frac{e^{2\sqrt{an}}}{n^{3/4}}$$

Oppenheim +

$$p(n) = an^\beta + O(n^\gamma)$$

implies

$$A(x) \sim Cx^{\beta+1} \frac{e^{2\sqrt{a \log x}}}{(\log x)^{3/4}}$$

Asymptotic Density of a subset B of A

Local Density

Global Density

$$\delta(B) = \lim_{n \rightarrow \infty} \frac{b(n)}{a(n)} \quad \Delta(B) = \lim_{x \rightarrow \infty} \frac{B(x)}{A(x)}$$

Dirichlet Density of a subset B of A

$$\partial(B) = \lim_{x \rightarrow \rho^-} \frac{\mathbf{B}(x)}{\mathbf{A}(x)} \quad \partial(B) = \lim_{x \rightarrow \alpha^+} \frac{\mathbf{B}(x)}{\mathbf{A}(x)}$$

need $\rho > 0$

need $\alpha < \infty$

Results that look as though they should have been proved long ago, but are actually very recent

Additive	Multiplicative
<p>[Bell] [Bell, . . . , Richmond]</p> <p>$\delta(P) = 0$ implies</p> $\begin{cases} \rho > 0 \text{ and} \\ \mathbf{A}(\rho) = \infty \end{cases}$	<p>[Warlimont]</p> <p>$\Delta(P) = 0$ implies</p> $\begin{cases} \alpha < \infty \\ \mathbf{A}(\alpha) = \infty \end{cases}$ <p>[Ruzsa] has a new proof.</p>

Ratio Test and Regular Variation at Infinity

Definition A sequence $s(n)$ is in RT_ρ if it is eventually positive and

$$\lim_{n \rightarrow \infty} \frac{s(n-1)}{s(n)} = \rho.$$

Definition A function $S(x)$ is in RV_α if it is eventually defined, eventually positive and

$$\lim_{t \rightarrow \infty} \frac{S(tx)}{S(t)} = x^\alpha$$

for $x > 0$. We say $S(x)$ has **regular variation at infinity of index α** .

A function in RV_0 is **slowly varying at infinity**.

The parallels in the concepts are clearer if we write:

$$\frac{s(n-k)}{s(n)} \rightarrow \rho^k \quad \text{and} \quad \frac{S(t/x)}{S(t)} \rightarrow x^{-\alpha}$$

One Reason we like RT and RV

All sets of the form

$$b + A$$

have asymptotic density iff RT_ρ holds.

If all sets of the form

$$b \cdot A$$

have asymptotic density then either RV_α holds or the system is discrete.

RV_α implies all sets of the form bA have asymptotic density.

The cases RT_1 and RV_0 were studied first (they yield 0–1 laws).

<p>[Bell]</p> <p>$p(n) = O(n^c)$ implies RT_1</p>	<p>[Bell]</p> <p>$P(x) = O((\log x)^c)$ implies RV_0</p>
<p>[Bell]</p> <p>$p(n) \in RT_1$ implies $a(n) \in RT_1$</p>	<p>[Bell] (Conj II)</p> <p>$P(x) \in RV_0$ implies $A(x) \in RV_0$</p>
<p>[Stewart]</p> <p>$\mathcal{A}_i \in RT_1$ implies $\mathcal{A}_1 + \mathcal{A}_2 \in RT_1$</p>	<p>[Odlyzko]</p> <p>$\mathcal{A}_i \in RV_0$ implies $\mathcal{A}_1 + \mathcal{A}_2 \in RV_0$</p>

[Karen Yeats] has a common generalization that gives the precise conditions for the sum/product of two number systems to preserve the existence of asymptotic density for all partition sets.

Schur's Theorem

[Schur 1918]

If

- $\mathbf{R}(x) = \mathbf{S}(x)\mathbf{T}(x)$
- $t(n) \in \mathbf{RT}_\rho$
- $\rho_{\mathbf{S}} > \rho$

then

$$\lim_{n \rightarrow \infty} \frac{r(n)}{t(n)} = \lim_{x \rightarrow \rho} \frac{\mathbf{R}(x)}{\mathbf{T}(x)}$$

[Burris/Yeats 2001]

If

- $\mathbf{R}(x) = \mathbf{S}(x)\mathbf{T}(x)$
- $T(x) \in \mathbf{RV}_\alpha$
- $\alpha_{\mathbf{S}} < \alpha$

then

$$\lim_{x \rightarrow \infty} \frac{R(x)}{T(x)} = \lim_{x \rightarrow \alpha} \frac{\mathbf{R}(x)}{\mathbf{T}(x)}$$

For example these can be used on the alternate form of the fundamental identities

$$\mathbf{A}(x) = e^{\sum_{m \geq 1} \mathbf{P}(x^m)/m}$$

$$\mathbf{A}(x) = e^{\sum_{m \geq 1} \mathbf{P}(mx)/m}$$

to focus attention on the asymptotics of the count function of $\boxed{e^{\mathbf{P}(x)}}$.

Applying Schur to get the asymptotics for the count function of a number system.

Using Schur:

$$\begin{aligned} & \mathbf{A}(x) \\ &= e^{\sum_{m \geq 1} \mathbf{P}(x^m)/m} \\ &= e^{\mathbf{P}(x)} \cdot e^{\sum_{m \geq 2} \mathbf{P}(x^m)/m} \end{aligned}$$

to get

$$a(n) \sim C \cdot [x^n] e^{\mathbf{P}(x)}$$

where

$$C = e^{\sum_{m \geq 2} \mathbf{P}(\rho^m)/m}$$

Using Schur Analog:

$$\begin{aligned} & \mathbf{A}(x) \\ &= e^{\sum_{m \geq 1} \mathbf{P}(mx)/m} \\ &= e^{\mathbf{P}(x)} \cdot e^{\sum_{m \geq 2} \mathbf{P}(mx)/m} \end{aligned}$$

to get

$$A(x) \sim C \cdot \sum_{n \leq x} [n^{-u}] e^{\mathbf{P}(u)}$$

where

$$C = e^{\sum_{m \geq 2} \mathbf{P}(m\alpha)/m}$$

What is a Partition Set?

Let P_1, \dots, P_k be a partition of the set P of indecomposables into finitely many sets.

Let γ_i be in one of the forms

$$\gamma_i = m_i \quad \gamma_i = (\leq m_i) \quad \gamma_i = (\geq m_i).$$

Then a subset B of the form

$$P_1^{\gamma_1} \dots P_k^{\gamma_k}$$

is a partition set.

In other words, B consists of all elements of A which have exactly

γ_1 indecomposables from P_1 ,

...

γ_k indecomposables from P_k .

Let (\diamond) be the property

All **partition sets** have asymptotic density.

Then

(\diamond) implies RT_ρ	(\diamond) implies discrete or RV_α
(\diamond) implies $\delta(P) = 0$	(\diamond) implies $\Delta(P) = 0$
RT_1 implies (\diamond)	RV_0 implies (\diamond)

On the next slide we look at **the most powerful conditions** known to imply that (\diamond) holds when $\rho < 1$, resp. $\alpha > 0$.

[Compton]

If $a(n) \in RT_\rho$ and

$$\frac{a(n-m)}{a(n)} \leq C\rho^m$$

for $K \leq m \leq n$
then (\diamond) holds.

[Sárközy]

If $A(x) \in RV_\alpha$ and

$$\frac{A(x/m)}{A(x)} \leq Cm^{-\alpha}$$

for $K \leq m \leq n$
then (\diamond) holds.

[Bell]

If $p(n) \in RT_\rho$ and

$$\limsup_{n \rightarrow \infty} np(n)\rho^n > 1$$

then Compton's conditions for (\diamond) hold.

[Bell] (Conj III)

If

$$P(x) \sim x^\alpha P_0(x) / \log x,$$

$P_0(x) \in RV_0$, eventually increasing and

$$\lim_{x \rightarrow \infty} P_0(x) \in (1/\alpha, \infty)$$

then Sárközy's conditions for (\diamond) hold.

Admissible Functions

These are functions for which one can use the famous saddle-point method to determine asymptotics for the count function of the series expansion.

[Compton]

If $A(x)$ is **Hayman-admissible** then Compton's conditions for (\diamond) hold.

[Burris, Warlimont, Yeats]

If $A(x)$ is **admissible** then Sárközy's conditions for (\diamond) hold.

A **Tenenbaum admissible** Dirichlet series is admissible.

Admissible Power Series

Suppose

$$\mathbf{A}(z) := \sum_{n \geq 0} a(n)z^n = e^{\mathbf{H}(z)}$$

is Hayman-admissible. Let the Taylor series expansion of $\mathbf{H}(re^{i\theta})$ about $\theta = 0$ be

$$\mathbf{H}(re^{i\theta}) = \mathbf{H}(r) + ia(r)\theta - b(r)\theta^2/2 + \dots$$

Theorem.

$$a(n) = \frac{\mathbf{A}(r)}{r^n \sqrt{2\pi b(r)}} \left(\exp\left(\frac{-(a(r) - n)^2}{2b(r)}\right) + R(r, n) \right)$$

where $R(r, n) \rightarrow 0$ as $r \rightarrow \rho-$, **uniformly** for $n \geq 0$.

The **saddle point** is r_n , the solution r of

$$a(r) - n = 0$$

Admissible Dirichlet Series

Suppose

$$\mathbf{A}(s) := \sum_{n \geq 1} a(n)/n^s = e^{\mathbf{H}(s)}$$

is an admissible Dirichlet series. Let the Taylor series expansion of $\mathbf{H}(\sigma + it)$ about $t = 0$ be

$$\mathbf{H}(\sigma + it) = \mathbf{H}(\sigma) + ia(\sigma)t - b(\sigma)t^2/2 + \dots$$

Theorem.

$$\widehat{A}(x) = \frac{x^{\sigma+1} \mathbf{A}(\sigma)}{\sigma(\sigma+1)\sqrt{2\pi b(\sigma)}} \left(\exp\left(\frac{-(a(\sigma) + \log x)^2}{2b(\sigma)}\right) + R(\sigma, x) \right)$$

where $R(\sigma, x) \rightarrow 0$ as $\sigma \rightarrow \alpha+$, **uniformly** for $x > 0$.

The **saddle point** is σ_x , the solution σ of

$$a(\sigma) + \log x = 0.$$