



T is degradable by the degrading map:



T extends N : (say $S_i \circ T_i = N_i$, $W_i = \text{iso ext of } S_i$)



Finally, apply Thm 2 to T :

$$Q^{(1)}(T) = Q(T) \leq p_1 Q^{(1)}(\bar{T}_1) + p_2 Q^{(1)}(\bar{T}_2)$$

By Thm 1, $Q(N) \leq Q^{(1)}(T)$

$$\leq Q(pN_0 + (1-p)N_1) \leq p Q^{(1)}(T_0) + (1-p) Q^{(1)}(\bar{T}_1)$$

T_0 deg ext of N_0

For the depolarizing channel =

① Consider the depolar channel $N(p) = (1-p)p + p Z p Z$

$$\text{let } U_1 Z U_1^\dagger = X, U_1 X U_1^\dagger = Y, U_1 Y U_1^\dagger = Z$$

Then the depolarizing channel =

$$N_p = \underbrace{\frac{1}{3} N}_{\text{depolarizing w/ } Z} + \underbrace{\frac{1}{3} U_1 \circ N \circ U_1^\dagger}_{\text{depolarizing w/ } X} + \underbrace{\frac{1}{3} U_1^2 \circ N \circ U_1^{\dagger 2}}_{\text{depolarizing w/ } Y}$$

Each of $N, U_1 \circ N \circ U_1^\dagger, U_1^2 \circ N \circ U_1^{\dagger 2}$ degradable

$$Q^{(1)} \text{ of each} = 1 - H(p)$$

$$\therefore Q(N_p) \leq 1 - H(p)$$

(a) recovering Rain's bdd

(b) the upper for each N_p comes from a deg extension

(2) Repeat above w/ $N(p) = \text{amplitude damping channel}$

$$Q(N_p) \leq H\left(\frac{1-\gamma}{2}\right) - H\left(\frac{\gamma}{2}\right)$$

$$\text{Where } \gamma = 4\sqrt{1-p}(1-\sqrt{1-p}) \quad (A3)$$

③ Consider $N_p = (1-4p)I + 4p N_{\frac{1}{4}}$ ($p \in [0, \frac{1}{4}]$)

Both I & $N_{\frac{1}{4}}$ have additive ext

[trivial for I , and use Thm 3 for $N_{\frac{1}{4}}$
& the fact it is known
to be anti degenerate]

$\therefore$ Thm 4 applies &

$$Q(N_p) \leq (1-4p) \cdot 1 + 4p \cdot 0 = 1-4p$$

(a) recovering Ravi's bdd, (b) each N_p has deg
ext giving the bdd

Putting everything together:

Let $f(p) = \max$ convex fcn upper bounded by
 $1 - H(p)$, $1 - 4p$, $H\left(\frac{1-x}{2}\right) - H\left(\frac{x}{2}\right)$

due to thm 4 & that every thing's derived
 from a deg ext, $Q(N) \leq f(p)$

