

**SPECIAL K**  
**Saturday November 3, 2012**  
**10:00 am - 1:00 pm**

**1:** Let  $f(x) = x^4 + 2x^3$ . Find the equation of a line which is tangent to the curve  $y = f(x)$  at two distinct points.

**2:** Find the area of the region

$$R = \{(x, y) \in \mathbf{R}^2 \mid (x^2 + y^2)^2 \leq 4x^2 \text{ and } x(x^2 + y^2) \leq 2\sqrt{3}xy\}.$$

**3:** Let  $x_n$  be the number of  $2 \times n$  matrices with entries in  $\{0, 1\}$  which do not contain the  $2 \times 2$  block  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ . Find  $\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n}$ .

**4:** Let  $k \geq 3$  be an integer. Let  $n = \frac{k(k+1)}{2}$ . Let  $S \subseteq \mathbf{Z}_n$  with  $|S| = k$ . Show that  $S + S \neq \mathbf{Z}_n$ . Note that  $|S|$  denotes the cardinality of  $S$  and  $S + S = \{x + y \mid x \in S, y \in S\}$ .

**5:** Let  $f : \mathbf{R} \rightarrow \mathbf{R}$ . Suppose that  $\lim_{x \rightarrow 0} f(x) = f(0) = 0$  and  $\lim_{x \rightarrow 0} \frac{f(2x) - f(x)}{x} = 0$ . Show that  $f$  is differentiable at 0 with  $f'(0) = 0$ .

**6:** Let  $\mathbf{Z}^+$  be the set of positive integers. Show that there exists a bijection  $f : \mathbf{Z}^+ \rightarrow \mathbf{Z}^+$  with the property that  $\prod_{k=1}^n f(k)$  is an  $n^{\text{th}}$  power for every  $n \in \mathbf{Z}^+$ .

**BIG E**  
**Saturday November 3, 2012**  
**10:00 am - 1:00 pm**

**1:** Find the volume of the solid

$$S = \{(x, y, z) \in \mathbf{R}^3 \mid (x^2 + y^2 + z^2)^2 \leq 4x^2 \text{ and } x(x^2 + y^2) \leq xz^2\}.$$

**2:** Find the number of  $3 \times n$  matrices with entries in  $\{0, 1\}$  which do not contain the  $2 \times 2$  block  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ .

**3:** Let  $k \geq 3$  be an integer. Let  $n = \frac{k(k+1)}{2}$ . Let  $S \subseteq \mathbf{Z}_n$  with  $|S| = k$ . Show that  $S + S \neq \mathbf{Z}_n$ . Note that  $|S|$  denotes the cardinality of  $S$  and  $S + S = \{x + y \mid x \in S, y \in S\}$ .

**4:** Let  $f : \mathbf{R}^2 \rightarrow \mathbf{R}$ . Suppose that  $f$  is continuous and that  $\int_0^1 f(a + tu) dt = 0$  for every point  $a \in \mathbf{R}^2$  and every unit vector  $u \in \mathbf{R}^2$ . Show that  $f$  is constant.

**5:** Let  $\mathbf{Z}^+$  be the set of positive integers. Show that there exists a bijection  $f : \mathbf{Z}^+ \rightarrow \mathbf{Z}^+$  with the property that  $\prod_{k=1}^n f(k)$  is an  $n^{\text{th}}$  power for every  $n \in \mathbf{Z}^+$ .

**6:** Let  $A$  be an  $n \times n$  matrix. Let  $u$  be an eigenvector of  $A$  for the eigenvalue 1. Suppose that all of the entries of  $A$  and all of the entries of  $u$  are positive. Show that the eigenspace for the eigenvalue 1 is 1-dimensional.