

1 Low-Rank Matrix Completion  
2 using  
3 Nuclear Norm Minimization  
4 and  
5 Facial Reduction\*

6 Shimeng Huang<sup>†</sup> Henry Wolkowicz<sup>‡</sup>

7 Tuesday 23<sup>rd</sup> May, 2017, 23:01, REVISION 2

8 **Abstract**

9 Minimization of the nuclear norm, **NNM**, is often used as a surrogate (convex relaxation)  
10 for finding the minimum rank completion (recovery) of a *partial matrix*. The minimum nuclear  
11 norm problem can be solved as a trace minimization semidefinite programming problem, **SDP**.  
12 Interior point algorithms are the current methods of choice for this class of problems. This  
13 means that it is difficult to: solve large scale problems; exploit sparsity; and get high accuracy  
14 solutions.

15 The **SDP** and its dual are regular in the sense that they both satisfy strict feasibility. In  
16 this paper we take advantage of the structure at optimality for the **NNM**. We show that even  
17 though strict feasibility holds, the facial reduction framework used for problems where strict  
18 feasibility fails can be successfully applied to obtain a proper face that contains the optimal  
19 set. This can dramatically reduce the size of the final **NNM** problem, while simultaneously  
20 guaranteeing a low-rank solution. This can be compared to identifying the active set in general  
21 nonlinear programming problems.

22 We include numerical tests for both exact and noisy cases.

23 **Keywords:** Low-rank matrix completion, matrix recovery, semidefinite programming (**SDP**),  
24 facial reduction, bicliques, Slater condition, nuclear norm, compressed sensing.

25 **AMS subject classifications:** 65J22, 90C22, 65K10, 52A41, 90C46

26 **Contents**

27 **1 Introduction**

**3**

---

\*Presented as Part of tutorial at DIMACS Workshop on Distance Geometry: Theory and Applications, July 26-29, 2016, [24]

<sup>†</sup>Department of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1. Research supported by the Undergraduate Student Research Awards Program, Natural Sciences and Engineering Research Council of Canada.

<sup>‡</sup>Department of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1; Research supported by The Natural Sciences and Engineering Research Council of Canada.

28	1.1	Outline . . . . .	3
29	<b>2</b>	<b>Background on LRMC, NNM, SDP</b>	<b>3</b>
30	2.1	Models . . . . .	4
31	2.2	Graph Representation of the Problem . . . . .	5
32	<b>3</b>	<b>Facial Reduction, Biclques, Exposing Vectors</b>	<b>7</b>
33	3.1	Preliminaries on faces . . . . .	7
34	3.2	Structure at Optimum . . . . .	9
35	3.3	Biclques, Weights and Final Exposing Vector . . . . .	11
36	<b>4</b>	<b>Noiseless Case</b>	<b>14</b>
37	4.1	Numerics Noiseless Case . . . . .	15
38	<b>5</b>	<b>Noisy Case</b>	<b>17</b>
39	5.1	Refinement Step with Dual Multiplier . . . . .	18
40	5.2	Numerics Noisy Case . . . . .	19
41	<b>6</b>	<b>Conclusion</b>	<b>19</b>
42		<b>Acknowledgment</b>	<b>20</b>
43		<b>Index</b>	<b>22</b>
44		<b>Bibliography</b>	<b>23</b>

## 45 List of Tables

46	1	<u>noiseless</u> : $r = 2$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	16
47	2	<u>noiseless</u> : $r = 3$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	16
48	3	<u>noiseless</u> : $r = 5$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	16
49	4	<u>noiseless</u> : $r = 6$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	17
50	5	<u>noiseless</u> : $r = 8$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	17
51	6	<u>noiseless</u> : $r = 3$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	17
52	7	<u>noiseless</u> : $r = 4$ ; 100% recovered; nullspace with eigs; mean 5 instances. . . . .	17
53	8	<u>noisy</u> : $r = 2$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	19
54	9	<u>noisy</u> : $r = 3$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	19
55	10	<u>noisy</u> : $r = 4$ ; $m \times n$ size; density $p$ ; mean 20 instances. . . . .	20

## 56 List of Algorithms

57	3.1	(finding the final exposing vector) . . . . .	13
----	-----	---	----

# 1 Introduction

We consider the intractable *low-rank matrix completion problem*, **LRMC**, i.e., the problem of finding the missing elements of a given *partial matrix* so that the completion has low-rank. This problem can be relaxed using the nuclear norm that can be then solved using a *semidefinite programming*, **SDP**, model. Though the resulting **SDP** and its dual satisfy strict feasibility, we show that it is implicitly highly degenerate and amenable to *facial reduction*, **FR**. This is done by taking advantage of the special structure *at the optimum* and by using the *exposing vector* approach, see [4]. The exposing vector approach is particularly amenable to the noisy case. Moreover, from **FR** we get a significant reduction in the size of the variables and a corresponding decrease in the possible rank of the solution. If the data is exact, then **FR** results in redundant constraints that we remove before solving for the low-rank solution. While if the data is contaminated with noise, **FR** yields an overdetermined semidefinite least squares problem. We *flip* this problem to minimize the nuclear norm using a Pareto frontier approach. Instead of removing constraints from the overdetermined problem, we exploit the notion of *sketch matrix* to reduce the size of the overdetermined problem. The sketch matrix approach is studied in e.g., [18].

The problem of **LRMC** has many applications to real applications in data science, model reduction, collaborative filtering (the well known Netflix problem) sensor network localization, pattern recognition and various other machine learning scenarios, e.g., [21, 22]. See also the recent work in [1, 19, 23] and the references therein. Of particular interest is the case where the data is contaminated with noise. This falls into the area of *compressed sensing* or *compressive sampling*. An extensive collection of papers, books, codes is available at the Compressive Sensing Resources, <http://dsp.rice.edu/cs>.

The convex relaxation of minimizing the rank using the nuclear norm, the sum of the singular values, is studied in e.g., [9, 10, 19]. The solutions can be found directly by subgradient methods or by using **SDP** with interior point methods or low-rank methods, again see [19]. Many other methods have been developed, e.g., [16]. The two main approaches for rank minimization, convex relaxations and spectral methods, are discussed in [3, 14] along with a new algebraic combinatorial approach. A related analysis from a different viewpoint using rigidity in graphs is provided in [20].

## 1.1 Outline

We continue in Section 2 with the basic notions for **LRMC** using the nuclear norm and with the graph framework that we employ. Then in Section 3 we include preliminaries on cone facial structure and the details on how to exploit **FR**, for the **SDP** model to minimize the nuclear norm problem. The main result for the reduction is in Lemma 3.1.

The results for the noiseless case are given in Section 4. This includes an outline of the basic approach in Algorithm 3.1 and empirical results from randomly generated problems. The noisy case follows in Section 5 with empirical results and a comparison with results in [23]. Concluding remarks are given in Section 6.

## 2 Background on LRMC, NNM, SDP

We now consider our problem within the known framework on relaxing the low-rank matrix completion problem using the nuclear norm minimization and then using **SDP** to solve the relaxation. For the standard results we follow and include much of the known development in the literature

99 e.g., [9, 10, 19]. In this section we also include several useful tools and a graph theoretic framework  
 100 that allows us to exploit **FR** at the optimum.

## 101 2.1 Models

Suppose that we are given a (random) low rank  $m \times n$  real matrix  $Z \in \mathbb{R}^{m \times n}$  where a subset of entries are *sampled*. The **LRMC** can be modeled as follows:

$$(LRMC) \quad \begin{aligned} & \min \quad \text{rank}(M) \\ & \text{s.t.} \quad \mathcal{P}_{\hat{E}}(M) = z, \end{aligned} \quad (2.1) \text{?basicsetting?}$$

102 where  $\hat{E}$  is the set of indices containing the known (*sampled*) entries of  $Z$ ,  $\mathcal{P}_{\hat{E}}(\cdot) : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^{|\hat{E}|}$   
 103 is the projection onto the corresponding entries in  $\hat{E}$ , and  $z = \mathcal{P}_{\hat{E}}(Z)$  is the vector of known  
 104 entries formed from  $Z$ . However, the rank function is not a convex function and the **LRMC** is  
 105 computationally intractable, e.g., [13].

To set up the problem as a convex optimization problem, we can relax the rank minimization using *nuclear norm minimization*, **NNM**:

$$(NNM) \quad \begin{aligned} & \min \quad \|M\|_* \\ & \text{s.t.} \quad \mathcal{P}_{\hat{E}}(M) = z, \end{aligned} \quad (2.2) \text{?basicnuclear?}$$

where the nuclear norm  $\|\cdot\|_*$  is the sum of the singular values, i.e.,  $\|M\|_* = \sum_i \sigma_i(M)$ . The general primal-dual pair of problems for the **NNM** problem is

$$\begin{aligned} \min_M \quad & \|M\|_* & \max_y \quad & \langle z, y \rangle \\ \text{s.t.} \quad & \mathcal{A}(M) = z, & \text{s.t.} \quad & \|\mathcal{A}^*(y)\| \leq 1, \end{aligned} \quad (2.3) \text{?eq:pdpairnucln}$$

106 where  $\mathcal{A} : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^t$  is a linear mapping,  $z \in \mathbb{R}^t$ ,  $\mathcal{A}^*$  is the *adjoint of*  $\mathcal{A}$ , and  $\|\cdot\|$  is the operator  
 107 norm of a matrix, i.e., the largest singular value. The matrix norms  $\|\cdot\|_*$  and  $\|\cdot\|$  are a dual pair  
 108 of matrix norms akin to the vector  $\ell_1, \ell_\infty$  norms on the vector of singular values. Without loss of  
 109 generality, we further assume that  $\mathcal{A}$  is *surjective*. In general, the linear equality constraint is an  
 110 underdetermined linear system. In our case, we restrict to the case that  $\mathcal{A} = \mathcal{P}_{\hat{E}}$ .<sup>1</sup>

111 **Proposition 2.1.** *Suppose that, in the primal-dual pair (2.3), there exists  $\hat{M}$  with  $\mathcal{A}(\hat{M}) = z$ .  
 112 Then the pair of programs in (2.3) are a convex primal-dual pair and they satisfy both primal and  
 113 dual strong duality, i.e., the optimal values are equal and both values are attained.*

114 *Proof.* This is shown in [19, Prop. 2.1]. That primal and dual strong duality holds can be seen from  
 115 the fact that the generalized Slater condition trivially holds for both programs using  $M = \hat{M}, y = 0$ ,  
 116 respectively.  $\square$

117 **Corollary 2.1.** *The optimal sets for the primal-dual pair in (2.3) are nonempty, convex, compact  
 118 sets.*

119 *Proof.* This follows since both problems are regular, i.e., since  $\mathcal{A}$  is surjective, we conclude that  
 120 the primal satisfies the *Mangasarian-Fromovitz constraint qualification*; while  $y = 0$  shows that the  
 121 dual satisfies strict feasibility. It is well known that this constraint qualification is equivalent to the  
 122 dual problem having a nonempty, convex, compact optimal set, e.g., [11].  $\square$

---

<sup>1</sup>Note that the linear mapping  $\mathcal{A} = \mathcal{P}_{\hat{E}}$  corresponding to sampling is surjective as we can consider  $\mathcal{A}(M)_{ij \in \hat{E}} = \text{trace}(E_{ij}M)$ , where  $E_{ij}$  is the  $ij$ -unit matrix.

123 The following proposition shows that the nuclear norm minimization problem is **SDP** representable,  
 124 i.e., we can embed the problem into an **SDP** and solve it efficiently. Here  $Y \succeq 0$  denotes the Löwner  
 125 partial order that  $Y$  is symmetric and positive semidefinite, denoted  $Y \in \mathcal{S}_+^{m+n}$ . We let  $\succ 0, \mathcal{S}_{++}^n$   
 126 denote *positive definite*.

**Proposition 2.2.** *The optimal primal-dual solution set in (2.3) is the same as that in the **SDP** primal-dual pair:*

$$\begin{aligned} \min \quad & \frac{1}{2} \text{trace}(W_1 + W_2) \\ \text{s.t.} \quad & Y = \begin{bmatrix} W_1 & M \\ M^T & W_2 \end{bmatrix} \succeq 0 \\ & \mathcal{A}(M) = z \end{aligned} \qquad \begin{aligned} \max_y \quad & \langle z, y \rangle \\ \text{s.t.} \quad & \begin{bmatrix} I_m & \mathcal{A}^*(y) \\ \mathcal{A}^*(y)^T & I_n \end{bmatrix} \succeq 0. \end{aligned} \quad (2.4) \text{?eq:pdpairnucln}$$

127

□

This means that after ignoring the  $\frac{1}{2}$  we can further transform the **NNM** problem as:

$$\begin{aligned} \min \quad & \|Y\|_* = \text{trace}(Y) \\ \text{s.t.} \quad & \mathcal{P}_{\bar{E}}(Y) = z \\ & Y \succeq 0, \end{aligned} \quad (2.5) \text{?sdpnuclear?}$$

128 where  $\bar{E}$  is the set of indices in  $Y$  that correspond to  $\hat{E}$ , the known entries of the upper right block  
 129 of  $\begin{bmatrix} 0 & Z \\ Z^T & 0 \end{bmatrix} \in \mathcal{S}^{m+n}$ .

When the data is contaminated with noise, we reformulate the strict equality constraint by allowing the observed entries in the output matrix to be perturbed within a tolerance  $\delta$  for the norm, where  $\delta$  is normally a known noise level of the data, i.e.,

$$\begin{aligned} \min \quad & \|Y\|_* = \text{trace}(Y) \\ \text{s.t.} \quad & \|\mathcal{P}_{\bar{E}}(Y) - z\| \leq \delta \\ & Y \succeq 0. \end{aligned} \quad (2.6) \text{?sdpnuclearinex}$$

130 We emphasize that there is no constraint on the diagonal blocks of  $Y$  in (2.4) or in (2.5).  
 131 Therefore, we can always obtain a positive definite feasible solution in this exact case by setting  
 132 the diagonal elements of  $Y$  to be large enough. Therefore strict feasibility, the *Slater constraint*  
 133 *qualification*, always holds.

## 134 2.2 Graph Representation of the Problem

Our sampling yields elements  $z = \mathcal{P}_{\hat{E}}(Z)$ . With the matrix  $Z$  and the sampled elements we can associate a bipartite graph  $G_Z = (U_m, V_n, \hat{E})$ , where

$$U_m = \{1, \dots, m\}, \quad V_n = \{1, \dots, n\}.$$

135 Our algorithm exploits finding complete bipartite subgraphs, *bicliques*, in  $G_Z$ . We now relate this  
 136 approach to finding cliques by using the larger symmetric matrix  $Y$  in (2.4). This allows us to  
 137 exploit **FR** and apply the clique algorithms from [4, 15]. However, we keep the biclique notation  
 138 as much as possible.

Therefore, for our needs we associate  $Z$  with the *undirected graph*,  $G = (V, E)$ , with node set  $V = \{1, \dots, m, m+1, \dots, m+n\}$  and edge set  $E$  that satisfies

$$\{\{ij \in V \times V : i < j \leq m\} \cup \{ij \in V \times V : m+1 \leq i < j \leq m+n\}\} \subseteq E \subseteq \{ij \in V \times V : i < j\}.$$

Note that as above,  $\bar{E}$  is the set of edges excluding the trivial ones, that is,

$$\bar{E} = E \setminus \left\{ \{ij \in V \times V : i \leq j \leq m\} \cup \{ij \in V \times V : m+1 \leq i \leq j \leq m+n\} \right\}.$$

Recall that a *biclique*  $\alpha$  in the graph  $G_Z$  is a complete bipartite subgraph in  $G_Z$  with corresponding complete submatrix  $z[\alpha]$ . This corresponds to a nontrivial<sup>2</sup> *clique* in the graph  $G$ , a complete subgraph in  $G$ . The cliques of interest are  $C = \{i_1, \dots, i_k\}$  with cardinalities

$$|C \cap \{1, \dots, m\}| = p \neq 0, \quad |C \cap \{m+1, \dots, m+n\}| = q \neq 0. \quad (2.7) \text{ ?eq:cardspq?}$$

The submatrix  $z[\alpha]$  of  $Z$  for the corresponding biclique from the clique  $C$  is

$$z[\alpha] \equiv X \equiv \{Z_{i(j-m)} : ij \in C\}, \quad \text{sampled } p \times q \text{ rectangular submatrix.} \quad (2.8) \text{ ?eq:Xspecif?}$$

139 These non-trivial cliques in  $G$  that correspond to bicliques of  $G_Z$  are at the center of our algorithm.

**Example 2.1** (biclique for  $X$ ). *Let the  $m \times n$  data matrix of rank  $r$  with  $m = 7, n = 6, r = 2$  be*

$$Z = \begin{bmatrix} -5 & 15 & 10 & -20 & -21 & -6 \\ 4 & 0 & 4 & 4 & 6 & 6 \\ -3 & -35 & -38 & 32 & 27 & -8 \\ 5 & -5 & 0 & 10 & 12 & 7 \\ 0 & -30 & -30 & 30 & 27 & -3 \\ 3 & -5 & -2 & 8 & 9 & 4 \\ 5 & 5 & 10 & 0 & 3 & 8 \end{bmatrix}.$$

After sampling we have unknown entries denoted by NA and known entries in

$$\begin{bmatrix} -5 & NA & 10 & -20 & NA & -6 \\ 4 & 0 & 4 & 4 & 6 & 6 \\ -3 & NA & NA & 32 & 27 & NA \\ 5 & NA & 0 & 10 & 12 & NA \\ NA & -30 & NA & NA & 27 & NA \\ 3 & -5 & -2 & 8 & NA & 4 \\ 5 & 5 & NA & 0 & 3 & NA \end{bmatrix}.$$

140 Then  $z = \mathcal{P}_{\bar{E}}(Z)$  denotes a vector representation of the known entries.  $\bar{E}$  denotes the corresponding  
141 indices for  $\bar{E}$  when  $Z$  is considered in the big matrix  $Y$  and  $E$  is formed from  $\bar{E}$  by adding on the  
142 indices corresponding to the diagonal blocks.

Suppose that our algorithm found a biclique  $\alpha$  with indices

$$\bar{U}_m = \{6, 1, 2\}, \quad \bar{V}_n = \{1, 4, 3, 6\}.$$

---

<sup>2</sup>For  $G$  we have the additional trivial cliques of size  $k$ ,  $C = \{i_1, \dots, i_k\} \subset \{1, \dots, m\}$  and  $C = \{j_1, \dots, j_k\} \subset \{m+1, \dots, m+n\}$ , that are not of interest to our algorithm.



$\phi$  is an *exposing vector*. Let  $S$  be a subset of the convex cone  $K$ , then  $\text{face}(S)$  is the smallest face of  $K$  containing  $S$ . It is known that: a face of a face is a face; an intersection of faces is a face; and essential for our algorithm is the following for finding an intersection of exposed faces  $F_i \trianglelefteq K, i = 1, \dots, k$ , see [4],

$$F_i = K \cap \phi_i^\perp, \forall i \implies \bigcap_{i=1}^k F_i = \left( \sum_{i=1}^k \phi_i \right)^\perp \cap K.$$

If  $K = \mathcal{S}_+^n$ , then the facial structure is well understood. Faces are characterized by the ranges or nullspaces of the matrices in the face. Let  $X \in \mathcal{S}_+^n$  be rank  $r$  and

$$X = [P \ Q] \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} [P \ Q]^T$$

be the (orthogonal) spectral decomposition with  $D \in \mathcal{S}_{++}^r$ . Then the smallest face containing  $X$  is

$$\text{face}(X) = P\mathcal{S}_+^r P^T = \mathcal{S}_+^n \cap (QQ^T)^\perp.$$

The matrix  $QQ^T$  is an *exposing vector* for  $\text{face}(X)$ . Moreover, the relative interior satisfies

$$\text{relint}(\text{face}(X)) = P\mathcal{S}_{++}^r P^T = \text{relint}(\text{face}(\hat{X})), \quad \forall \hat{X} \in \text{relint}(\text{face}(X)),$$

150 i.e. the face and the exposing vectors are characterized by the eigenspace of any  $\hat{X}$  in the relative  
151 interior of the face.

152 For our application we use the following view of facial reduction and exposed faces.

**Theorem 3.1** ([5, Theorem 4.1]). *Consider a linear transformation  $\mathcal{M}: \mathcal{S}^n \rightarrow \mathbb{R}^m$  and a nonempty feasible set*

$$\mathcal{F} := \{X \in \mathcal{S}_+^n : \mathcal{M}(X) = b\},$$

for some point  $b \in \mathbb{R}^m$ . Then a vector  $v$  exposes a proper face of  $\mathcal{M}(\mathcal{S}_+^n)$  containing  $b$  if, and only if,  $v$  satisfies the auxiliary system

$$0 \neq \mathcal{M}^* v \in \mathcal{S}_+^n \quad \text{and} \quad \langle v, b \rangle = 0.$$

153 Let  $N$  denote the smallest face of  $\mathcal{M}(\mathcal{S}_+^n)$  containing  $b$ . Then the following are true.

154 1. We always have  $\mathcal{S}_+^n \cap \mathcal{M}^{-1}N = \text{face}(\mathcal{F})$ , the smallest face containing  $\mathcal{F}$ .

2. For any vector  $v \in \mathbb{R}^m$  the following equivalence holds:

$$v \text{ exposes } N \iff \mathcal{M}^* v \text{ exposes } \text{face}(\mathcal{F}). \tag{3.1} \text{?eq:exposnonuni}$$

155 □

156 The result in (3.1) details the facial reduction process for the matrix completion problem using  
157 exposing vectors. More precisely, if  $B \succeq 0$  is a principal submatrix of the data and  $\text{trace } VB =$   
158  $0, V \succeq 0$ , then  $V$  provides an exposing vector for the image of the coordinate map. We can then  
159 complete  $V$  with zeros to get  $Y \in \mathcal{S}_+^n$  an exposing vector for  $\mathcal{F}$ . Define the *triangular number*,  
160  $t(n) = n(n+1)/2$ , and the projection  $\text{vec}: \mathcal{S}^n \rightarrow \mathbb{R}^{t(n)}$  that vectorizes the upper-triangular part of  
161 a symmetric matrix columnwise. We let  $\text{Mat}: \mathbb{R}^{t(n)} \rightarrow \mathcal{S}^n$  denote the inverse mapping.



**Corollary 3.1.** Suppose that  $1 < k < n$  and  $\mathcal{M}$  in Theorem 3.1 is the coordinate projection onto the leading principal submatrix of order  $k, m = t(k)$ . Let  $B \in \mathcal{S}_+^k, b = \text{vec}(B) \in \mathbb{R}^{t(k)}$ , i.e., for  $X \in \mathcal{S}^n$ , we have

$$\mathcal{M}(X)_{ij} = b_{ij}, \quad \forall 1 \leq i \leq j \leq k.$$

Let

$$V \in \mathcal{S}_+^k, \text{trace}(VB) = 0, v = \text{vec } V.$$

Then  $Y = \mathcal{M}^*v$  is an exposing vector for the feasible set  $\mathcal{F}$ , i.e.,

$$\text{trace}(Y(\mathcal{F})) = 0.$$

162 *Proof.* The proof follows immediately from Theorem 3.1 as  $v$  exposes  $N$  and  $Y = \mathcal{M}^*v$  is an  
163 exposing vector for  $\text{face}(\mathcal{F})$ .  $\square$

### 164 3.2 Structure at Optimum

165 The results in Section 2 can now be used to prove the following special structure at the optimum.  
166 This structure is essential in our **FR** scheme.

**Corollary 3.2.** Let  $M^*$  be optimal for the primal in (2.4) with  $\text{rank}(M^*) = r_M$ . Let  $M^* = UDV^T$  be the compact SVD,  $D \in \mathcal{S}_{++}^{r_M}$ . Define

$$W_1 = UDU^T, \quad W_2 = VDV^T, \tag{3.2} \text{?eq:Winorm?}$$

and

$$Y = \begin{bmatrix} W_1 & M^* \\ (M^*)^T & W_2 \end{bmatrix} = \begin{bmatrix} U \\ V \end{bmatrix} D \begin{bmatrix} U \\ V \end{bmatrix}^T. \tag{3.3} \text{?eq:YDUV?}$$

Then we have  $Y \succeq 0$  and optimal in (2.4) with  $\text{rank}(Y) =: r_Y = r_M$  and

$$\|M^*\|_* = \frac{1}{2} \text{trace}(Y) = \text{trace}(D).$$

167 *Proof.* The matrices  $U, V$  have orthonormal columns. Therefore  $\text{trace}(Y) = 2 \text{trace}(D) = 2\|M\|_*$ .  
168  $\square$

Now suppose that there is a biclique  $\alpha$  of  $G_Z$  and a corresponding *sampled submatrix*,  $z[\alpha] \equiv X \in \mathbb{R}^{p \times q}$ , of  $Z \in \mathbb{R}^{m \times n}$ , with  $\text{rank}(X) = r_X$ . Without loss of generality, after row and column permutations if needed, we can assume that

$$Z = \begin{bmatrix} Z_1 & Z_2 \\ X & Z_3 \end{bmatrix}. \tag{3.4} \text{?eq:ZX?}$$

Let the **SVD** be

$$X = [U_1 \quad U_X] \begin{bmatrix} \Sigma & 0 \\ 0 & 0 \end{bmatrix} [V_1 \quad V_X]^T, \quad \Sigma \in \mathcal{S}_{++}^{r_X}; \tag{3.5} \text{?eq:svdX?}$$

and we have a full rank factorization  $X = \bar{P}\bar{Q}^T$  obtained using the compact **SVD**

$$X = \bar{P}\bar{Q}^T = U_1\Sigma V_1^T, \quad \bar{P} = U_1\Sigma^{1/2}, \quad \bar{Q} = V_1\Sigma^{1/2}.$$

We see below that such a desirable  $X$  (after a permutation if needed), that corresponds to a biclique  $\alpha \in G_Z$ ,  $z[\alpha] \equiv X$ , and at least one nontrivial exposing vector, is characterized by

$$C_X = \{m - p + 1, \dots, m, m + 1, \dots, m + q\}, \quad r \leq \min\{p, q\} < \max\{p, q\}. \quad (3.6) \text{ ?eq:suffbig?}$$

169 Here we use the *target rank*,  $r$ . We can exploit the information using these bicliques to obtain  
 170 exposing vectors of the *optimal face*,  $F^*$ , i.e., the smallest face of  $\mathcal{S}_+^{m+n}$  that contains the set of  
 171 optimal solutions of (2.4).

By abuse of notation, we can rewrite the optimal  $Y$  as in (3.3) in Corollary 3.2 and get

$$0 \preceq Y = \begin{bmatrix} U \\ P \\ Q \\ V \end{bmatrix} D \begin{bmatrix} U \\ P \\ Q \\ V \end{bmatrix}^T = \left[ \begin{array}{c|c|c|c} UDU^T & UDP^T & UDQ^T & UDV^T \\ \hline PDU^T & PDP^T & PDQ^T & PDV^T \\ \hline QDU^T & QDP^T & QDQ^T & QDV^T \\ \hline VDU^T & VDP^T & VDQ^T & VDV^T \end{array} \right]. \quad (3.7) \text{ ?eq:Ypartit?}$$

We see that  $X = PDQ^T = \bar{P}\bar{Q}^T$ . Since we assume that  $X$  satisfies (3.6) and so is *big enough*, we conclude that generically  $r_X = r_Y = r$ , see Lemma 3.2 below, and that the ranges satisfy

$$\begin{aligned} \text{Range}(X) &= \text{Range}(P) = \text{Range}(\bar{P}) = \text{Range}(U_1), \\ \text{Range}(X^T) &= \text{Range}(Q) = \text{Range}(\bar{Q}) = \text{Range}(V_1). \end{aligned} \quad (3.8) \text{ ?eq:PQbar?}$$

172 This is the key for facial reduction as we can use an *exposing vector* formed as  $U_X U_X^T$  as well as  
 173  $V_X V_X^T$ .

**Lemma 3.1** (basic **FR**). *Let  $Y$  be an optimal solution of the primal problem in (2.4) with  $\text{rank}(Y) = r = r_Z$ , i.e., the **NNM** heuristic yields an optimal  $Y$  that successful solves the **LRMC** problem. Let  $X \in \mathbb{R}^{p \times q}$  be a sampled submatrix of  $Z$  be as in (3.4) (after a permutation if needed) with  $\text{rank}(X) = r_X = r_Z \leq \min\{p, q\} < \max\{p, q\}$ , and with **SVD** as in (3.5). We now add appropriate blocks of zeros to the block exposing vectors  $U_X U_X^T, V_X V_X^T$  and get*

$$W_X = \left[ \begin{array}{c|c|c|c} 0 & 0 & 0 & 0 \\ \hline 0 & U_X U_X^T & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right] + \left[ \begin{array}{c|c|c|c} 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & V_X V_X^T & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right] = \left[ \begin{array}{c|c|c|c} 0 & 0 & 0 & 0 \\ \hline 0 & U_X U_X^T & 0 & 0 \\ \hline 0 & 0 & V_X V_X^T & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right].$$

Then all three matrices are exposing vectors for the optimal face, i.e., for  $W_X$  we have  $0 \neq W_X \succeq 0, W_X Y = 0$ . Moreover, if  $T$  is a full column rank matrix with the columns forming a basis for  $\text{Null}(W_X)$ , the nullspace of  $W_X$ , then a facial reduction step for the optimal face,  $F^*$ , the minimal face containing the optimal set, yields

$$F^* \preceq T \mathcal{S}_+^{(n+m)-(p+q-2r)} T^T.$$

174 *Proof.* That  $U_X U_X^T, V_X V_X^T$  provide exposing vectors is by construction based on the argument for  
 175 (3.8). The result follows from the fact that the sum of exposing vectors is an exposing vector.  
 176 Moreover, the block diagonal structure of the exposing vectors guarantees that the ranks add up  
 177 to get the size of the smaller face containing  $F^*$ .  $\square$

**Example 3.1** (pair of exposing vectors). We now present a matrix  $Y \in \mathcal{S}_+^{11}$  with  $\text{rank}(Y) = 2$ . Here  $(m, n) = (6, 5)$ .

$$Y = \begin{bmatrix} 0.0059877 & 0.10551 & -0.011994 & -0.036276 & -0.073807 & -0.049863 & -0.049795 & -0.02602 & 0.01314 & 0.022035 & -0.012187 \\ 0.10551 & 2.1638 & 0.035252 & -0.6439 & -1.5417 & -0.77074 & -1.9215 & -0.13496 & -0.23004 & 0.13318 & 0.239 \\ -0.011994 & 0.035252 & 0.22366 & 0.068878 & -0.04733 & 0.18725 & -0.74543 & 0.31405 & -0.39999 & -0.25065 & 0.39174 \\ -0.036276 & -0.6439 & 0.068878 & 0.21984 & 0.45085 & 0.30043 & 0.31772 & 0.15267 & -0.072518 & -0.12958 & 0.066865 \\ -0.073807 & -1.5417 & -0.04733 & 0.45085 & 1.1006 & 0.52923 & 1.4401 & 0.064661 & 0.20335 & -0.069711 & -0.2089 \\ -0.049863 & -0.77074 & 0.18725 & 0.30043 & 0.52923 & 0.45348 & 0.044817 & 0.33131 & -0.27295 & -0.27387 & 0.26224 \\ -0.049795 & -1.9215 & -0.74543 & 0.31772 & 1.4401 & 0.044817 & 3.9923 & -0.89251 & 1.4727 & 0.69104 & -1.4538 \\ -0.02602 & -0.13496 & 0.31405 & 0.15267 & 0.064661 & 0.33131 & -0.89251 & 0.45673 & -0.54736 & -0.3667 & 0.53491 \\ 0.01314 & -0.23004 & -0.39999 & -0.072518 & 0.20335 & -0.27295 & 1.4727 & -0.54736 & 0.72824 & 0.43489 & -0.71429 \\ 0.022035 & 0.13318 & -0.25065 & -0.12958 & -0.069711 & -0.27387 & 0.69104 & -0.3667 & 0.43489 & 0.29471 & -0.42483 \\ -0.012187 & 0.239 & 0.39174 & 0.066865 & -0.2089 & 0.26224 & -1.4538 & 0.53491 & -0.71429 & -0.42483 & 0.7007 \end{bmatrix}$$

We sample the elements in rows 4, 5, 6 and columns 7, 8, 9, 10 to obtain the  $(p = 3) \times (q = 4)$  matrix  $X$ . We let  $U_X, V_X$ , denote orthonormal bases for the nullspaces of  $X, X^T$ , respectively, i.e.,

$$XU_X = 0, \quad X^T V_X = 0.$$

Then the two exposing vectors are  $U_X U_X^T$  and  $V_X V_X^T$ , filled in with zeros. After adding them together, we get

$$W = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.81985 & -0.17015 & -0.34459 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -0.17015 & 0.035313 & 0.071516 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -0.34459 & 0.071516 & 0.14483 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0.023237 & -0.058066 & -0.12587 & 0.059006 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -0.058066 & 0.57988 & 0.34589 & 0.34729 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -0.12587 & 0.34589 & 0.68409 & -0.28395 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0.059006 & 0.34729 & -0.28395 & 0.71279 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

We see that

$$\|WY\| = 7.67638e-16,$$

178 thus verifying to 15 decimals that the sum of the two exposing vectors is indeed an exposing vector  
179 for  $\text{face}(Y)$ .

180 We emphasize that here we knew the two principal diagonal blocks of  $Y$  that corresponded to the  
181 clique  $C = \{4, 5, 6, 7, 8, 9, 10\}$ . But in general we do not and only know the sampled  $X$ . However,  
182 generically (Lemma 3.2, below), we get the exposing vectors correctly as done here. Moreover, here  
183 we only had a single sampled  $X$  and could permute it to an easy position to illustrate the exposing  
184 vector. In general, we will have many of these that are identified by the indices determining the  
185 corresponding clique. We then add them up to get a final exposing vector which is used for the  
186 **FR** step.

### 187 3.3 Bicliques, Weights and Final Exposing Vector

188 Given a partial matrix  $Z \in \mathbb{R}^{m \times n}$ , we need to find nontrivial bicliques  $\alpha$  and corresponding sampled  
189 submatrices  $z[\alpha] = X$  according to the properties in (2.7) and (2.8). Intuitively, we may want to  
190 find bicliques with size as large as possible so that we can expose  $Y$  immediately. However, we do  
191 not want to spend a great deal of time finding large bicliques. Instead we find it is more efficient  
192 to find many medium-size bicliques, satisfying the size-rank condition  $r \leq \min\{p, q\} < \max\{p, q\}$ .  
193 This rank condition guarantees that at least one of the two exposing vectors found from the biclique  
194 is not zero. We can then add the exposing vectors obtained from the equivalent cliques for these  
195 bicliques to finally expose a small face containing the optimal  $Y$ . This is equivalent to dealing with  
196 a small number of large bicliques. This consideration also comes from the expense of the singular

197 value decomposition for the sampled submatrix  $z[\alpha] = X$  for  $U_X, V_X$  in (3.5) when the biclique is  
 198 large.

199 The following lemma shows that, generically, we can restrict the search to bicliques correspond-  
 200 ing to a sampled submatrix  $X \in \mathbb{R}^{p \times q}$  that satisfies the rank condition  $r \leq \min\{p, q\} < \max\{p, q\}$   
 201 without losing rank magnitude.

202 **Lemma 3.2** (generic rank property). *Let  $r$  be a positive integer and  $Z_1 \in \mathbb{R}^{m \times r}$  and  $Z_2 \in \mathbb{R}^{n \times r}$   
 203 be continuous random variables with i.i.d. elements. Set  $Z = Z_1 Z_2^T$  and let  $X \in \mathbb{R}^{p \times q}$  be any  
 204 submatrix of  $Z$  with  $\min\{p, q\} \geq r$ . Then  $\text{rank}(X) = r$  with probability 1.*

205 *Proof.* Without loss of generality, we can assume that  $X$  is the top left corner of  $Z$ . Therefore,  
 206  $X = \bar{Z}_1 \bar{Z}_2$  for appropriate (top part) submatrices  $\bar{Z}_i$  of  $Z_i$ ,  $i = 1, 2$ . By the rank condition, we  
 207 have that  $X = \bar{Z}_1 \bar{Z}_2^T$  is a full rank factorization of  $X$  generically.  $\square$

208 **Remark 3.1.** *In our numerical tests we generate our matrices  $Z = Z_1 Z_2^T$  as done in the above  
 209 Lemma 3.2. Therefore, it clear generically that submatrices  $X$  with the specified size restriction  
 210 have  $\text{rank}(X) = \text{rank}(Z)$  generically. It is not clear if the converse is true, i.e., whether a given  
 211 random matrix  $Z$  with  $\text{rank}(Z) = r$  and full rank factorization  $Z = Z_1 Z_2^T$  implies that  $Z_1, Z_2$  have  
 212 random elements.*<sup>4</sup>

213 With the existence of noise (e.g., Gaussian), we know that generically the  $X$  found can only  
 214 have a higher rank but not a lower rank than  $r$ . In this case, since we assume that we know the  
 215 target rank of  $X$ , we can adjust the exposing vector so that it will not over-expose the completion  
 216 matrix. If the target rank is not known, then it can be estimated during the algorithm up to a  
 217 given tolerance, i.e., for a give sampled  $p \times q$  submatrix  $X$  we estimate the rank. If the estimated  
 218 rank  $r < \min\{p, q\}$ , then by our (generic) Lemma 3.2, we can assume that we have found our target  
 219 rank for  $Z$ . If this is not the case, then we need to look for bicliques of larger size. As soon as we  
 220 find  $r = \text{rank}(X) < \min\{p, q\}$  then we have found our estimated target rank  $r$ .

221 After finding a biclique  $\alpha$  corresponding to a sampled submatrix  $X$  and its full rank factorization  
 222  $X = \bar{P} \bar{Q}^T$ , we then construct *biclique weights*  $u_X^P$  and  $u_X^Q$  to measure how *noisy* the corresponding  
 223 exposing vectors are. We essentially use the *Eckart-Young distance* [7] to the nearest matrix of  
 224 rank  $r$  and include the size of the submatrix. If the problem is *noiseless* and we know the target  
 225 rank for  $Z$ , then these distances for the submatrices are 0.

**Definition 3.1** (biclique noise). *Suppose that  $X \in \mathbb{R}^{p \times q}$ , with singular values  $\sigma_1 \geq \dots \geq \sigma_{\min\{p, q\}}$ ,  
 is a given sampled submatrix corresponding to a biclique of the graph of the partial matrix  $Z$ . Let  
 $r$  be the target rank. Define the biclique noise*

$$u_X^P := \frac{\sum_{i=r+1}^{\min\{p, q\}} \sigma_i^2}{0.5p(p-1)}, \quad u_X^Q := \frac{\sum_{i=r+1}^{\min\{p, q\}} \sigma_i^2}{0.5q(q-1)}.$$

**Definition 3.2** (biclique weights). *Let  $\Theta$  be the set of all bicliques. For each biclique  $X \in \Theta$  of  
 the partial matrix  $Z$ , let  $p, q, u_X^P, u_X^Q$  be defined as in Definition 3.1. Let*

$$S = \sum_{X \in \Theta} (u_X^P + u_X^Q).$$

---

<sup>4</sup>The authors thank Dmitriy Drusvyatskiy for the simplification of our original proof of Lemma 3.2. Further discussions are given in [6].

Define the biclique weight

$$w_X^P = 1 - \frac{u_X^P}{S}, \quad w_X^Q = 1 - \frac{u_X^Q}{S}.$$

Using Lemma 3.1, we now present Algorithm 3.1, page 13, to find an exposing vector  $Y_{expo}$  for the optimal face, i.e., we get the block diagonal

$$0 \neq \left[ \begin{array}{c|c} \sum_{X \in \Theta} w_X^P U_X U_X^T & 0 \\ \hline 0 & \sum_{X \in \Theta} w_X^Q V_X V_X^T \end{array} \right] = Y_{expo} \succeq 0, \quad Y_{expo} Y^* = 0, \quad \forall \text{ optimal } Y^*.$$

Note that if

$$Y_{expo} = [U \quad V] \begin{bmatrix} \Lambda & 0 \\ 0 & 0 \end{bmatrix} [U \quad V]^T,$$

is the (orthogonal) spectral decomposition of  $Y_{expo}$ , with  $\Lambda \in \mathfrak{S}_{++}^{r_e}$ , then the optimal face satisfies

$$F^* \trianglelefteq V S_+^{m+n-r_e} V^T, \quad V = \begin{bmatrix} V_P & 0 \\ 0 & V_Q \end{bmatrix}.$$

Thus this **FR** process reduces the size of the problem.

---

**Algorithm 3.1** (finding the final exposing vector)

---

- 1: **INPUT:** partial matrix  $Z \in \mathcal{M}^{m \times n}$ , target rank  $r$ ;
- 2: **OUTPUT:** final *blocked exposing vector*  $Y_{expo}$  that exposes the optimal face for (2.5)
- 3: **PREPROCESSING:**  
find a set of bicliques  $\Theta$  of size within the given range {minsize, maxsize} with  $r < \text{minsize}$ ;
- 4: **for** each biclique  $\alpha \in \Theta$  and corresponding  $z[\alpha] = X$  **do**
- 5:    $[U_X, V_X] \leftarrow$  from **SVD** of  $X$  in 3.5
- 6:    $W_X^P \leftarrow U_X U_X^T$ ;  
    calculate biclique noise  $u_X^P$ ;
- 7:    $W_X^Q \leftarrow V_X V_X^T$ ;  
    calculate biclique noise  $u_X^Q$ ;
- 8: **end for**
- 9: calculate all the biclique weights  $w_X^i, i = P, Q, \alpha \in \Theta$ , from biclique noise;
- 10: sum over bicliques the weighted blocked matrices filled in with appropriate zeros.

$$0 \neq Y_{expo} \leftarrow \left[ \begin{array}{c|c} \sum_{X \in \Theta} w_X^P W_X^P & 0 \\ \hline 0 & \sum_{X \in \Theta} w_X^Q W_X^Q \end{array} \right].$$

11: **return**  $Y_{expo}$

---

226

227 **Remark 3.2.** We do not need to look for large bicliques in Algorithm 3.1 since we can take ad-  
 228 vantage of the fact that adding exposing vectors results in an exposing vector. Moreover, finding a  
 229 biclique is equivalent to finding a clique in  $G$ . Therefore, we use the algorithms for finding cliques  
 230 given in [15] and [4, Algorithm 2].

231 **4 Noiseless Case**

In the noiseless case, the biclique noise is 0 and the weights are all 1 and so ignored. The **FR** step finds the *blocked* exposing vector  $Y_{expo}$  and the *blocked* basis for  $\text{Null}(Y_{expo})^5$  given by the columns of

$$V = \begin{bmatrix} V_P & 0 \\ 0 & V_Q \end{bmatrix}, \quad V_P^T V_P = I_{r_p}, \quad V_Q^T V_Q = I_{r_q},$$

thus defining the dimensions  $r_p + r_q = r_v$ . Therefore an original feasible  $Y$  can be expressed as

$$Y = V R V^T = \begin{bmatrix} V_P R_p V_P^T & V_P R_{pq} V_Q^T \\ V_Q R_{pq}^T V_P^T & V_Q R_q V_Q^T \end{bmatrix} \quad (4.1) \text{?eq:VRVT?}$$

where the blocked

$$R = \begin{bmatrix} R_p & R_{pq} \\ R_{pq}^T & R_q \end{bmatrix} \in \mathcal{S}^{r_v}, r_v < m + n.$$

232 This means the problems (2.4) and (2.5) are in general reduced to the much smaller dimension  
 233  $\mathbb{R}^{r_p \times r_q}$ . And if we find enough bicliques we expect a reduction to  $r_p = r_q = r, r_v = 2r$ , twice the  
 234 target rank. If this is the case then we have exact recovery that can be obtained by a simple least  
 235 squares solution. Otherwise, we have to rely on the **NNM** heuristic.

The reduced model for  $Y$  after **FR** with **NNM** is

$$\begin{aligned} \min \quad & \text{trace}(R) && (= \text{trace}(V R V^T)) \\ \text{s.t.} \quad & \mathcal{P}_{\tilde{E}}(V_P R_{pq} V_Q^T) = z && (= \mathcal{P}_{\tilde{E}}(Z)) \\ & R = \begin{bmatrix} R_p & R_{pq} \\ R_{pq}^T & R_q \end{bmatrix} \succeq 0. \end{aligned} \quad (4.2) \text{?frnuclearnoise}$$

The **FR** typically results in low values for  $r_p, r_q$  and in the exact data case *many* of the linear equality constraints become redundant, i.e., we generally end up with an overdetermined linear system. We use the compact QR decomposition<sup>6</sup> to identify which constraints to choose that result in a linearly independent set with a relatively low condition number. Thus we have eliminated a portion of the sampling and we get the linear system

$$\mathcal{M}(R_{pq}) := \mathcal{P}_{\tilde{E}}(V_P R_{pq} V_Q^T) = \tilde{z}, \text{ for some } \tilde{E} \subseteq \hat{E}, \quad (4.3) \text{?eq:smallsyst?}$$

236 and  $\tilde{z}$  is the vector of corresponding elements in  $z$ .

1. We first solve the simple semidefinite constrained least squares problem

$$\min_{R \in \mathcal{S}_+^{r_v}} \|\mathcal{P}_{\tilde{E}}(V_P R_{pq} V_Q^T) - \tilde{z}\|.$$

237 If the optimal  $R$  has attained the target rank, then the exactness of the data implies that  
 238 necessarily the optimal value is zero; and we are done. (In fact, the **SDP** constraint is  
 239 redundant here as  $R$  can always be completed using an **SVD** decomposition of  $R_{pq}$ .)

<sup>5</sup>The MATLAB command *null* was used to find an orthonormal basis for the nullspace. However, this requires an SVD decomposition and fails for huge problems. In that case, we used the Lanczos approach with *eigs*.

<sup>6</sup>The MATLAB economical version function  $[\sim, R, E] = qr(\Phi, 0)$  finds the list of constraints for a well conditioned representation, where  $\Phi$  denotes the matrix of constraints.

240 2. If  $R$  does not have the target rank in Item 1 above, then we solve (4.2) for our minimum  
 241 nuclear norm solution. We note that the linear transformation  $\mathcal{M}$  in (4.3) is not one-one.  
 242 Therefore, we often need to add a small regularizing term to the objective, i.e., we use  
 243  $\min \text{trace}(R) + \gamma \|R\|_F$  with small  $\gamma > 0$ .

#### 244 4.1 Numerics Noiseless Case

245 We now present experiments with the algorithm on random noiseless instances. Averages (computer  
 246 times, rank, residuals) on **twenty** random instances are included in the tables<sup>7</sup>.

247 The tests were run with MATLAB version R2016a, and Windows 8, on a Dell Optiplex 9030,  
 248 Intel(R) Core(TM) i7-4790 CPU @ 3.60GHz and 16 GB RAM. <sup>8</sup> The times we present are the  
 249 wall-clock times in seconds. For the semidefinite constrained least squares problems we used the  
 250 MATLAB addon CVX [12] for simplicity. This means our computer times could be improved if we  
 251 replaced CVX with a recent **SDP** solver.

We generate the instances as done in the recent work [8]. The target matrices are obtained from  $Z = Z_L Z_R^T$ , where  $Z_L \in \mathbb{R}^{m \times r}$  and  $Z_R \in \mathbb{R}^{r \times n}$ . Each entry of the two matrices  $Z_L$  and  $Z_R$  is generated independently from a standard normal distribution  $N(0, 1)$ . We then generate a sparse  $m \times r$  matrix to obtain the random indices that are sampled. We evaluate our results using the same measurement as in [8], which we call “Residual” in our tables. It is calculated as:

$$\text{Residual} = \frac{\|\hat{Z} - Z\|_F}{\|Z\|_F},$$

252 where  $Z$  is the target matrix,  $\hat{Z}$  is the output matrix that we find, and  $\|\cdot\|_F$  is the Frobenius norm.

253  
 254 We observe that we far outperform the results in [8] both in accuracy and in time; and we  
 255 solve much larger problems. We are not as competitive for the low density problems as our method  
 256 requires a sufficient number of cliques in  $G$  (bicliques in  $G_Z$ ). We could combine our preprocessing  
 257 approach using the bicliques before the method in [8] is applied.

258 In Tables 1 to 7 we present the results with noiseless data with target ranks ranging from  $r = 2$   
 259 to  $r = 6$ . We see that we get efficient *high* accuracy recovery in *every* instance. The accuracy is  
 260 significantly higher than what one can expect from an **SDP** interior point solver. The computer  
 261 time is almost entirely spent on finding the matrix representation and on its QR factorization that  
 262 is used as a heuristic for finding a correct subset of well-conditioned linear constraints. However,  
 263 we do not use any refinement steps for these tests. For higher rank and sparse problems we end  
 264 up with a larger **FR** problem and a large matrix representation. This can be handled using  
 265 the sketch matrix and refinement described in the noisy case. For the lower density problems, we  
 266 remove the rows and columns of the original data matrix corresponding to zero diagonal elements  
 267 of the final exposing matrix. These rows and columns have no sampled entries in them and so it  
 268 does not make sense to include them in the algorithm. We include the percentage of the number of  
 269 elements of the original data matrix that are recovered and the corresponding percentage residual.  
 270 Since the accuracy is high for this recovered submatrix, it can then be used with further sampling  
 271 to recover the complete original matrix.

<sup>7</sup>The density  $p$  in the tables are reported as “mean( $p$ )” because the real density obtained is usually not the same as the one set for generating the problem. We report the mean of the real densities over the five instances.

<sup>8</sup>The Tables 4 with rank 6 and 5 with rank 8 were done using a MacBookPro12,1, Intel Core i5, 2.7 GHz with two cores and 8 GB RAM. The version of MATLAB was the same R2016a.

272 These problems involved relatively low target ranks  $r = 2$  to  $r = 8$ . Larger ranks mean that we  
 273 need to find larger bicliques/cliques, e.g.,  $r = 20$  means that the cliques need to be of size bigger  
 274 than 40. This means that the values for  $r_p, r_q$  can be large and we need to solve a large **SDP** least  
 275 squares problem. We include a purify step to do this in the noisy case discussed below.

276 Note that the largest problems in the last of the noiseless tables, 6 and 7, have, respectively,  
 277 48,000,000 and 50,000,000 data entries in  $Z$  with approximately 35,000,000 unknown values that  
 278 were recovered successfully with **extremely** high accuracy. The target rank was recovered in every  
 279 instance. We used the MATLAB command *null* in Table 6 to find the nullspaces to derive  $V$  in  
 280 (4.1). This is based on an **SVD** decomposition of a full matrix and is expensive. We used MATLAB  
 281 *eigs* rather than *null* in Table 7 which resulted in lower computer times but lower accuracy. We  
 282 could not use *null* in the noisy case as this results in essentially full rank each time due to the noise.  
 283 We changed to a sparse QR decomposition which estimates the rank, has the lowest computer times  
 284 while still maintaining high accuracy.

Table 1: noiseless:  $r = 2$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications			$r_v$	Rcvrd (%Z)	Time (s)	Rank	Residual (%Z)
$m$	$n$	mean( $p$ )					
2100	4000	0.33	4.00	100.00	46.35	2.0	1.4298e-13
2100	4000	0.26	4.00	100.00	44.69	2.0	4.3546e-14
2100	4000	0.22	4.00	100.00	43.43	2.0	9.8758e-14
2100	4000	0.18	4.00	100.00	42.66	2.0	1.4409e-13
2100	4000	0.14	4.00	99.78	42.16	2.0	8.9667e-14

Table 2: noiseless:  $r = 3$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications			$r_v$	Rcvrd (%Z)	Time (s)	Rank	Residual (%Z)
$m$	$n$	mean( $p$ )					
2100	4000	0.33	6.00	100.00	50.46	3.0	8.6855e-13
2100	4000	0.26	6.00	100.00	49.88	3.0	1.0738e-12
2100	4000	0.22	6.00	100.00	48.56	3.0	1.1436e-12
2100	4000	0.18	6.00	99.81	47.90	3.0	2.5695e-12
2100	4000	0.14	6.20	95.15	46.69	3.0	8.5525e-12

Table 3: noiseless:  $r = 5$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications			$r_v$	Rcvrd (%Z)	Time (s)	Rank	Residual (%Z)
$m$	$n$	mean( $p$ )					
2100	4000	0.45	10.00	100.00	52.48	5.0	2.2232e-10
2100	4000	0.42	10.00	100.00	53.16	5.0	2.3748e-11
2100	4000	0.39	10.00	100.00	52.45	5.0	1.5950e-10
2100	4000	0.36	10.00	99.99	49.78	5.0	4.5280e-11
2100	4000	0.33	10.00	99.79	47.60	5.0	2.5057e-10



Table 4: noiseless:  $r = 6$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications			$r_v$	Rcvrd (%Z)	Time (s)	Rank	Residual (%Z)
$m$	$n$	mean( $p$ )					
2100	4000	0.48	12.00	100.00	84.83	6.0	4.4311e-10
2100	4000	0.45	12.00	99.98	78.81	6.0	7.2856e-10
2100	4000	0.42	12.00	99.78	76.11	6.0	1.3813e-11
2100	4000	0.39	12.00	98.46	73.48	6.0	2.8688e-10
2100	4000	0.36	13.65	92.08	74.52	6.0	5.6545e-08

Table 5: noiseless:  $r = 8$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications			$r_v$	Rcvrd (%Z)	Time (s)	Rank	Residual (%Z)
$m$	$n$	mean( $p$ )					
1000	3000	0.53	16.10	96.39	37.29	8.0	1.1072e-10
1000	3000	0.50	17.65	88.99	36.50	8.0	4.6569e-10
1000	3000	0.48	32.15	71.66	72.14	8.5	2.0413e-07

Table 6: noiseless:  $r = 3$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications			$r_v$	Rcvrd (%Z)	Time (s)	Rank	Residual (%Z)
$m$	$n$	mean( $p$ )					
700	2000	0.33	6.00	100.00	5.58	3.0	2.6857e-13
1000	5000	0.33	6.00	100.00	58.31	3.0	3.0256e-12
1400	9000	0.33	6.00	100.00	296.91	3.0	1.4185e-12
1900	14000	0.33	6.00	100.00	1043.46	3.0	1.9995e-12
3000	16000	0.33	6.00	100.00	1758.76	3.0	2.5250e-12

Table 7: noiseless:  $r = 4$ ; 100% recovered; nullspace with eigs; mean 5 instances.

Specifications			Time (s)	Rank	Residual (%Z)
$m$	$n$	mean( $p$ )			
700	2000	0.36	12.80	4.0	1.5217e-12
1000	5000	0.36	49.66	4.0	1.0910e-12
1400	9000	0.36	131.53	4.0	6.0304e-13
1900	14000	0.36	291.22	4.0	3.4847e-11
2500	20000	0.36	798.70	4.0	7.2256e-08

## 285 5 Noisy Case

286 This case is similar to the noiseless case but with the addition of a refinement step. (The refinement  
287 step can also be used for the noiseless case when the **FR** problem dimension  $r_v$  is too large.)  
288 We include the rank and residual outputs for both before refinement and the total of both after  
289 refinement. We see that in most cases when the graph is sufficiently dense, refinement is *not*  
290 needed, and near perfect completion (recovery) is obtained relative to the noise. In particular, the  
291 low target rank was attained most times.

We generate the data as in the noiseless case and then perturb the known entries by additive noise, i.e.,

$$Z_{ij} \leftarrow Z_{ij} + \sigma \xi_t \|Z\|_\infty, \quad \forall ij \in \bar{E},$$

where  $\xi_t \sim N(0, 1)$  and  $\sigma$  is a noise factor that can be changed. The computer and software were similar as in the noiseless case. The tests were run on MATLAB version R2016a as above, but on a Dell Optiplex 9030, with Windows 8, Intel(R) Core(TM) i7-4790 CPU @ 3.60GHz and 16 GB RAM.

As above we proceed to first complete **FR** in order to reduce the dimension of  $Y$ , i.e., the dimension of  $R$ ,  $r_v$ , is dramatically smaller. In the low density and/or high rank cases it is difficult to find enough cliques and in this case the final exposing vector  $Y_{expo}$  contains many zero rows. This essentially means that we have not sampled rows and/or columns of  $Z$ . In these cases we have ignored the rows and columns that used no sampled entries.

After **FR** we first solve the simple semidefinite constrained least squares problem

$$\delta_0 = \min_{R \in \mathcal{S}_+^{r_v}} \|\mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - z\|, \quad z = \mathcal{P}_{\hat{E}}(Z).$$

However, unlike in the noiseless case, we cannot remove redundant constraints, even though there may be many. This problem is now highly overdetermined and may also be ill-posed in that the constraint transformation may not be one-one. We use the notion of *sketch matrix* to reduce the size of the system, e.g., [18]. The matrix  $A$  is a random matrix of appropriate size with a relatively small number of rows in order to dramatically decrease the size of the constrained least squares problem

$$\delta_0 = \min_{R \in \mathcal{S}_+^{r_v}} \|A (\mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - z)\|.$$

As noted in [18], this leads to surprisingly good results. If  $s$  is the dimension of  $R$ , then we use a random sketch matrix of size  $2t(s) \times |\hat{E}|$ , where  $t(\cdot)$  is the number of variables on and above the diagonal of a symmetric matrix, i.e., the triangular number

$$t(s) = \frac{s(s+1)}{2}.$$

If the optimal  $R$  has the correct target rank, then we are done.

## 5.1 Refinement Step with Dual Multiplier

If the result from the constrained least squares problem does not have the target rank, we now use this  $\delta_0$  as a best target value for our parametric approach as done in [4]. Our **NNM** problem can be stated as:

$$\begin{aligned} \min \quad & \text{trace}(R) \\ \text{s.t.} \quad & \left\| A \left( \mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - z \right) \right\| \leq \delta_0 \\ & R \succeq 0. \end{aligned} \tag{5.1} \text{?frsdpnuclear?}$$

To attempt to find a lower rank solution, we use the approach in [4] and *flip* this problem:

$$\begin{aligned} \varphi(\tau) := \min \quad & \left\| A \left( \mathcal{P}_{\hat{E}}(V_P R_{pq} V_Q^T) - z \right) \right\| + \gamma \|R\|_F \\ \text{s.t.} \quad & \text{trace}(R) \leq \tau \\ & R \succeq 0. \end{aligned} \tag{5.2} \text{?eq:flipNoisyfi}$$

303 As in the noiseless case, the least squares problem may be underdetermined. We add a regularizing  
304 term  $+\gamma\|R\|_F$  to the objective with  $\gamma > 0$  small. The starting value of  $\tau$  is obtained from the  
305 unconstrained least squares problem, and from which we can reduce the value of the trace of  $R$   
306 to reduce the nuclear norm and so heuristically reduce the rank. We refer to this process as the  
307 refinement step.

308 This process requires a tradeoff between low-rank and low-error. Specifically, the trace con-  
309 straint may not be tight at the starting value of  $\tau$ , which means we can lower the trace of  $R$   
310 without sacrificing accuracy, however, if the trace is pushed lower than necessary, the error starts  
311 to get larger. To detect the balance point between low-rank and low-error, we exploit the role as  
312 sensitivity coefficient for the *dual multiplier* of the inequality constraint. The value of the dual  
313 variable indicates the rate of increase of the objective function. When the the dual multiplier be-  
314 comes positive then we know that decreasing  $\tau$  further will increase the residual value. We have  
315 used the value of .01 to indicate that we should stop decreasing  $\tau$ .

## 316 5.2 Numerics Noisy Case

317 The noisy case results with increasing ranks 2, 3, 4 and various sizes and densities follow in Tables  
318 8, 9, 10. With the densities we use the recovery is essentially 100%. We consider problems with  
319 relatively high density to ensure that we can find enough cliques. We have not included tests with  
320 higher rank as those are done in the noiseless case and are similar here.

Table 8: noisy:  $r = 2$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications				Rcvd (%Z)	Time (s)		Rank		Residual (%Z)	
$m$	$n$	% noise	$p$		initial	refine	initial	refine	initial	refine
1100	3000	0.50	0.33	100.00	33.72	48.53	2.00	2.00	8.53e-03	8.53e-03
1100	3000	1.00	0.33	100.00	33.67	49.09	2.00	2.00	2.70e-02	2.70e-02
1100	3000	2.00	0.33	100.00	34.13	48.84	2.00	2.00	9.75e-02	9.75e-02
1100	3000	3.00	0.33	100.00	36.34	92.73	5.00	5.00	5.48e-01	1.40e-01
1100	3000	4.00	0.33	100.00	51.45	186.28	11.00	8.00	1.25e+00	1.28e-01

Table 9: noisy:  $r = 3$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications				Rcvd (%Z)	Time (s)		Rank		Residual (%Z)	
$m$	$n$	% noise	$p$		initial	refine	initial	refine	initial	refine
700	1000	1.00	0.33	99.99	2.58	16.54	3.35	3.35	1.29e+00	1.07e+00
800	2000	1.00	0.33	100.00	10.72	29.59	3.75	3.75	1.15e+00	1.07e+00
900	4000	1.00	0.33	100.00	61.92	94.40	3.25	3.20	1.47e+00	1.07e+00
1000	8000	1.00	0.33	100.00	404.26	672.60	8.70	6.45	3.94e+00	7.11e-01
1100	16000	1.00	0.33	100.00	3553.81	4230.73	9.00	6.65	4.00e+00	6.66e-01

## 321 6 Conclusion

322 In this paper we have shown that we can apply facial reduction through the exposing vector  
323 approach used in [4] in combination with the nuclear norm heuristic to efficiently find low-rank

Table 10: noisy:  $r = 4$ ;  $m \times n$  size; density  $p$ ; mean 20 instances.

Specifications				Rcvd (%Z)	Time (s)		Rank		Residual (%Z)	
$m$	$n$	% noise	$p$		initial	refine	initial	refine	initial	refine
1100	3000	0.00	0.36	100.00	30.27	42.44	4.00	4.00	9.04e-13	9.04e-13
1200	3500	1.00	0.33	100.00	52.48	198.22	8.20	6.70	6.45e+00	1.08e+00
1300	4000	2.00	0.32	100.00	81.09	388.68	11.80	7.85	1.88e+01	1.28e+00
1400	4500	3.00	0.31	100.00	117.40	573.87	12.00	7.40	2.51e+01	1.45e+00
1500	5000	4.00	0.31	100.00	142.86	699.06	12.00	6.90	2.42e+01	1.61e+00

324 matrix completions. This exploits the degenerate structure of the optimal solution set even though  
 325 the nuclear norm heuristic problem itself satisfies strict feasibility.

326 Specifically, whenever enough complete bipartite subgraphs are available for the graph of the  
 327 matrix of the problem, we are able to find a proper face with a *significantly reduced dimension* that  
 328 contains the optimal solution set. We then solve this smaller minimum trace problem by *flipping*  
 329 the problem and using a refinement with a parametric point approach. If we cannot find enough  
 330 bicliques, the matrix can still be partially completed. Having an insufficient number of bicliques  
 331 is indicative of not having enough initial data to recover the unknown elements for our algorithm.  
 332 This is particularly true for large  $r$  where larger bicliques are needed. Throughout we see that the  
 333 facial reduction both regularizes the problem and reduces the size and often allows for a solution  
 334 without any refinement.

335 Our *preliminary* numerical results are promising as they efficiently and accurately recover large  
 336 scale problems. The numerical tests are ongoing with improvements in using biclique algorithms  
 337 rather than clique algorithms thus exploiting the block structure of the cliques; and with solving  
 338 the lower dimensional flipped problems. In our paper we have started our tests with knowing the  
 339 target rank  $r$ . In forthcoming tests we plan on using estimating the target rank using sampled  
 340 submatrices.

341 In addition, theoretical results on *exact recovery* are discussed in many papers, e.g., [2, 3, 19].  
 342 They use the so-called *restricted isometry property*, **RIP**, for vectors extended to the matrix case.  
 343 However, the **RIP** condition is difficult to verify. It appears from our work above that exact recovery  
 344 guarantees can be guaranteed from rigidity questions in the graph of  $Z$ , i.e., in the number and  
 345 density of the bicliques. Moreover, there are interesting questions on how to extend these results  
 346 from the simple matrix completion to general solutions of linear equations,  $\mathcal{A}(Z) = b$ , where  $\mathcal{A}$  is  
 347 some linear transformation.

## 348 Acknowledgment

349 The authors would like to thank Nathan Krislock for his help with parts of the MATLAB coding;  
 350 and we would also like to thank two anonymous referees for carefully reading the paper and helping  
 351 us improve the details and the presentation.

## Index

- 352  $C_X = \{m - p + 1, \dots, m, m + 1, \dots, m + q\}$ , 10  
 353  $F \trianglelefteq K$ , face, 7  
 354  $F^*$ , optimal face, 10  
 355  $G = (V, E)$ , undirected graph, 6  
 356  $G_Z = (U_m, V_n, \hat{E})$ , 5  
 357  $K^*$ , dual cone, 7  
 358  $P$ -biclique weight,  $w_X^P$ , 13  
 359  $R_p$ , 14  
 360  $R_q$ , 14  
 361  $R_{pq}$ , 14  
 362  $X \in \mathbb{R}^{p \times q}$ , sampled submatrix, 9  
 363  $Y \succeq 0$ , positive semidefinite, 5  
 364  $Y_{expo}$ , 13  
 365  $\text{Mat}$ , 8  
 366  $\mathcal{S}^n$ , space of real symmetric matrices, 5  
 367  $\mathcal{S}_+^n$ , positive semidefinite matrices, 5  
 368  $\mathcal{S}_{++}^n$ , positive definite matrices, 5  
 369  $\bar{E}$ , indices in  $Y$  of sampled entries of  $Z$ , 5  
 370  $\hat{E}$ , indices of sampled entries of  $Z$ , 4  
 371  $\text{vec}$ , 8  
 372  $\|\cdot\|_*$ , nuclear norm, 4  
 373  $\|\cdot\|_F$ , Frobenius norm, 15  
 374  $r$ , target rank, 10  
 375  $r_X$ , 9  
 376  $r_Y$ , 9  
 377  $r_p$ , 14  
 378  $r_q$ , 14  
 379  $t(k) = k(k + 1)/2$ , triangular number, 8  
 380  $u_X^P$ , biclique noise, 12  
 381  $u_X^P$ , biclique weight, 12  
 382  $w_X^P$ ,  $P$ -biclique weight, 13  
 383  $z[\alpha]$ , complete submatrix, 6  
 384 **FR**, facial reduction, 3  
 385 **LRMC**, low-rank matrix completion problem,  
 386 3, 4  
 387 **NNM**, nuclear norm minimization, 4  
 388 **RIP**, restricted isometry property, 20  
 389 **SDP**, semidefinite programming, 3  
 390 adjoint of  $\mathcal{A}$ , 4  
 391 biclique, 5, 6  
 392 biclique noise,  $u_X^P$ , 12  
 393 biclique weight,  $u_X^P$ , 12  
 394 clique, 6  
 395 complete submatrix,  $z[\alpha]$ , 6  
 396 conjugate face,  $F^*$ , 7  
 397 dual cone,  $K^*$ , 7  
 398 dual multiplier, 19  
 399 Eckart-Young distance, 12  
 400 exposed face, 7  
 401 exposing vector, 3, 8, 10  
 402 face,  $F \trianglelefteq K$ , 7  
 403 x **FR**, 3  
 404 Frobenius norm,  $\|\cdot\|_F$ , 15  
 405 indices in  $Y$  of sampled entries of  $Z$ ,  $\bar{E}$ , 5  
 406 indices of sampled entries of  $Z$ ,  $\hat{E}$ , 4  
 407 low-rank matrix completion problem, **LRMC**, 4  
 408 x **LRMC**, 3  
 409 Mangasarian-Fromovitz constraint qualification,  
 410 4  
 411 x **NNM**, 4  
 412 nuclear norm,  $\|\cdot\|_*$ , 4  
 413 optimal face,  $F^*$ , 10  
 414 orthogonal complement, 7  
 415 partial matrix, 3  
 416 positive definite matrices,  $\mathcal{S}_{++}^n$ , 5  
 417 positive semidefinite matrices,  $\mathcal{S}_+^n$ , 5  
 418 x **RIP**, 20  
 419 sampled submatrix,  $z[\alpha] \equiv X \in \mathbb{R}^{p \times q}$ , 9  
 420 x **SDP**, 3  
 421 sketch matrix, 3, 18  
 422 Slater constraint qualification, 5  
 423 space of real symmetric matrices,  $\mathcal{S}^n$ , 5  
 424 target rank,  $r$ , 10  
 425 triangular number,  $t(n) = n(n + 1)/2$ , 8  
 426 undirected graph,  $G = (V, E)$ , 6

## References

- MR3436616** [1] J.D. Blanchard, J. Tanner, and K. Wei. CGIHT: conjugate gradient iterative hard thresholding for compressed sensing and matrix completion. *Inf. Inference*, 4(4):289–327, 2015. 3  
429
- MR2565240** [2] E.J. Candès and B. Recht. Exact matrix completion via convex optimization. *Found. Comput. Math.*, 9(6):717–772, 2009. 20  
431
- MR2723472** [3] E.J. Candès and T. Tao. The power of convex relaxation: near-optimal matrix completion. *IEEE Trans. Inform. Theory*, 56(5):2053–2080, 2010. 3, 20  
433
- ChDrWor14** [4] D. Drusvyatskiy, N. Krislock, Y-L. Cheung Voronin, and H. Wolkowicz. Noisy sensor network localization: robust facial reduction and the Pareto frontier. *SIAM Journal on Optimization*, pages 1–30, 2017. arXiv:1410.6852, 20 pages, to appear. 3, 5, 7, 8, 13, 18, 19  
435  
436
- DrPaWor14** [5] D. Drusvyatskiy, G. Pataki, and H. Wolkowicz. Coordinate shadows of semidefinite and Euclidean distance matrices. *SIAM J. Optim.*, 25(2):1160–1178, 2015. 8  
438
- DrusWolk16** [6] D. Drusvyatskiy and H. Wolkowicz. The many faces of degeneracy in conic optimization. Technical report, University of Waterloo, Waterloo, Ontario, 2017. submitted, 97 pages. 12  
440
- EckartYoung36** [7] C. Eckart and G. Young. The approximation of one matrix by another of lower rank. *Psychometrica*, 1(3):211–218, 1936. 12  
442
- angLiuTohZhou15** [8] E.X. Fang, H. Liu, K.-C. Toh, and W.-X. Zhou. Max-norm optimization for robust matrix recovery. Technical report, Department of Operations Research and Financial Engineering, Princeton University, Princeton, 2015. 15  
444  
445
- Fazel02** [9] M. Fazel. *Matrix Rank Minimization with Applications*. PhD thesis, Stanford University, Stanford, CA, 2001. 3, 4  
447
- elhindiBoyd01** [10] M. Fazel, H. Hindi, and S.P. Boyd. A rank minimization heuristic with application to minimum order system approximation. In *Proceedings American Control Conference*, pages 4734–4739, 2001. 3, 4  
449  
450
- MR0489903** [11] J. Gauvin. A necessary and sufficient regularity condition to have bounded multipliers in nonconvex programming. *Math. Programming*, 12(1):136–138, 1977. 4  
452
- cvxunpub1** [12] M. Grant and S. Boyd. CVX: Matlab software for disciplined convex programming, version 1.21. <http://cvxr.com/cvx>, April 2011. 15  
454
- kaRaghWeigz14** [13] M. Hardt, R. Meka, P. Raghavendra, and B. Weitz. Computational limits for matrix completion. In *JMLR Proceedings*, pages 1–23. JMLR, 2014. 4  
456
- MR3417586** [14] F.J. Király, L. Theran, and R. Tomioka. The algebraic combinatorial approach for low-rank matrix completion. *J. Mach. Learn. Res.*, 16:1391–1436, 2015. 3  
458
- kriswolk15** [15] N. Krislock and H. Wolkowicz. Explicit sensor network localization using semidefinite representations and facial reductions. *SIAM Journal on Optimization*, 20(5):2679–2708, 2010. 5, 7, 13  
460  
461

- [JainDhillon4509](#) [16] R. Meka, P. Jain, and I.S. Dhillon. Guaranteed rank minimization via singular value projection. Technical Report Advanced Studies in Pure Mathematics, Department of Computer Science, University of Texas at Austin, Austin, 2009. 3  
463  
464
- [PatakiSVW4599](#) [17] G. Pataki. Geometry of Semidefinite Programming. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook OF Semidefinite Programming: Theory, Algorithms, and Applications*. Kluwer Academic Publishers, Boston, MA, 2000. 7  
466  
467
- [Wainwright4515](#) [18] M. Pilanci and M.J. Wainwright. Randomized sketches of convex programs with sharp guarantees. *IEEE Trans. Infor. Theory*, 61(9):5096–5115, 2015. 3, 18  
469
- [ParriloFazel](#) [19] B. Recht, M. Fazel, and P. Parrilo. Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization. *SIAM Rev.*, 52(3):471–501, 2010. 3, 4, 20  
471
- [MR2595541](#) [20] A. Singer and M. Cucuringu. Uniqueness of low-rank matrix completion by rigidity theory. *SIAM J. Matrix Anal. Appl.*, 31(4):1621–1641, 2009/10. 3  
473
- [MR2717223](#) [21] N. Srebro. *Learning with matrix factorizations*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—Massachusetts Institute of Technology. 3  
475
- [MR2203286](#) [22] N. Srebro and A. Shraibman. Rank, trace-norm and max-norm. In *Learning theory*, volume 3559 of *Lecture Notes in Comput. Sci.*, pages 545–560. Springer, Berlin, 2005. 3  
477
- [MR3440480](#) [23] J. Tanner and K. Wei. Low rank matrix completion by alternating steepest descent methods. *Appl. Comput. Harmon. Anal.*, 40(2):417–429, 2016. 3  
479
- [talkdimacs46](#) [24] H. Wolkowicz. Tutorial: Facial reduction in cone optimization with applications to matrix completions, at: Dimacs workshop on distance geometry: Theory and applications, 2016. Based on survey paper: The many faces of degeneracy in conic optimization, (with D. Drusvyatskiy).  
481  
482  
483 1