

C&O 367
Assignment 1

Due on Tuesday, Jan. 11, 2005 Instructor H. Wolkowicz

Notes:

1. Please explain your answers and proofs carefully. A yes or no does not constitute a valid answer; nor does a numerical value with no explanation constitute a valid answer.
-

1. GEOMETRY

For a vector $x = (x_1, \dots, x_n)^T \in \mathfrak{R}^n$, we define the (*Euclidean*) *norm* and (*Euclidean*) *inner product*

$$\|x\| := (x_1^2 + \dots + x_n^2)^{1/2},$$

$$\langle x, y \rangle := x \cdot y := x_1 y_1 + \dots + x_n y_n.$$

Theorem 1 (*Cauchy-Schwarz*) $|x \cdot y| \leq \|x\| \|y\|$, with equality if and only if $x = \lambda y$, for some λ .

EXERCISES

- (a) (10) Explain the geometrical significance for the vectors x and y of:
 - i.

$$x \cdot y = 0.$$

- ii.

$$x \cdot y > 0.$$

- (b) (10) Prove that the function $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$ defined by $f(x) = a \cdot x$ is continuous, where a is a given vector.

2. CALCULUS

For a function $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$, the *gradient* is denoted

$$\nabla g := \begin{pmatrix} \frac{\partial g}{\partial x_1} \\ \frac{\partial g}{\partial x_2} \\ \dots \\ \frac{\partial g}{\partial x_n} \end{pmatrix}$$

EXERCISES

- (a) (5) If $g(x) = \|x\|$, calculate $\nabla g(x)$.
- (b) (10) Suppose $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$, $a, b \in \mathfrak{R}^n$, and $f(t) := g(a + tb)$. Calculate $f'(t)$.

Suppose $f : \mathfrak{R} \rightarrow \mathfrak{R}$ is infinitely differentiable at $x = a$. The *Taylor Series* of f about a is:

$$f(a) + f'(a)(x - a) + \frac{1}{2!}f''(a)(x - a)^2 + \frac{1}{3!}f'''(a)(x - a)^3 + \dots$$

EXERCISE

Write down the Taylor series of:

- (a) (5)
- $$f(x) = x^3, \text{ about } x = 1.$$
- (b) (5)
- $$f(x) = \log(1 + x), \text{ about } x = 0.$$

3. TOPOLOGY

The *open ball* $B(x; r) := \{y \in \mathfrak{R}^n : \|x - y\| < r\}$. Suppose that D is a subset of \mathfrak{R}^n .

Interior: $x \in \text{int } D$ if there exists $r > 0$ with $B(x; r) \subset D$.

Closure: $x \in \text{cl } D$ if there exists a sequence $x^k \in D$ with $x^k \rightarrow x$.

Boundary: $x \in \partial D$ if $x \in \text{cl } D \setminus \text{int } D$.

D is *open* if $D = \text{int } D$. D is *closed* if $D = \text{cl } D$.

EXERCISES

- (a) (15) For each of the following sets, find the interior, the closure, and the boundary. Then determine which of the sets are open, closed, neither, or both.

i.

$$\{(x_1, x_2) : x_1 \geq 0, x_2 \geq 0\}.$$

ii.

$$\{(x_1, x_2) : x_1 > 0, x_2 > 0\}.$$

iii.

$$\{(x_1, x_2) : x_1 > 0, x_2 \geq 0\}.$$

iv.

$$\mathfrak{R}^n$$

v.

$$\{(x_1, x_2) : x_1^2 + x_2^2 < 0\}.$$

vi.

$$\emptyset.$$

- (b) i. (10) Prove that D is closed if and only if the complement D^c is open.
ii. (10) Prove that $x \in \partial D$ if and only if for any $r > 0$ there exists a $y \in B(x; r) \cap D$ and a $z \in B(x; r) \cap D^c$.

4. MATRICES

EXERCISES

(a) (10) Let

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 2 & 1 \end{pmatrix}$$

- i. Calculate the determinant of A .
- ii. Calculate the rank of A .
- iii. What is the rank of A^T .

(b) (10) Let

$$B = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$$

Calculate the eigenvalues and eigenvectors of B .

(c) (10) Let

$$C = \begin{pmatrix} 3 & 1 & 1 & 4 \\ \rho & 4 & 10 & 1 \\ 1 & 7 & 17 & 3 \\ 2 & 2 & 4 & 3 \end{pmatrix}$$

Using elementary transformations (elementary row and column operations), find the value of ρ that minimizes the rank of C .