Bibliography

[1] D. Vanden Abeele and H. Deconinck. Development of a Godunov-type solver for 2D ideal MHD problems. Technical Report 13, von Karman Institute, 1995.

[2] A. Akhiezer, G. Liubarskii, and R. Polovin. The stability of shock waves in magnetohydrodynamics. *Sov. Phys. JETP*, 35:507–511, 1959.

[3] A. I. Akhiezer, I. A. Akhiezer, R. V. Polovin, A. G. Sitenko, and K. N. Stepanov. *Plasma electrodynamics*, volume 1. Pergamon Press, Oxford, 1975.

[4] J. Edward Anderson. *Magnetohydrodynamic shock waves.* PhD thesis, M. I. T., 1963.

[5] D. S. Balsara and D. S. Spicer. A staggered mesh algorithm using high order Godunov fluxes to ensure solenoidal magnetic fields in magnetohydrodynamic simulations. *J. Comput. Phys.*, 149:270–292, 1999.

[6] A. A. Barmin, A. G. Kulikovskiy, and N. V. Pogorelov. Shock-capturing approach and nonevolutionary solutions in magnetohydrodynamics. *J. Comput. Phys.*, 126:77–90, 1996.

[7] Timothy J. Barth. Numerical methods for gasdynamic systems on unstructured meshes. In D. Kröner, M. Ohlberger, and C. Rohde, editors, *An introduction to recent developments in theory and numerics for conservation laws*, Lecture notes in Computational Science and Engineering, pages 195–284. Springer-Verlag, Berlin, 1998.

[8] D. Biskamp. *Nonlinear Magnetohydrodynamics.* Cambridge University Press, Cambridge, 1993.

[9] B. Bottin, M. Carbonaro, S. Zemsch, and G. Degrez. Aerothermodynamic design of an inductively-coupled plasma wind tunnel. *AIAA Paper*, 97-2498, 1997.

[10] J. U. Brackbill and D. C. Barnes. The effect of nonzero $\nabla \cdot \vec{B}$ on the numerical solution of the magnetohydrodynamic equations. *J. Comput. Phys.*, 35:426–430, 1980.

[11] M. Brio and C. C. Wu. Characteristic fields for the equations of magnetohydrodynamics. *Contemporary mathematics*, 60:19–23, 1987.

[12] M. Brio and C. C. Wu. An upwind differencing scheme for the equations of ideal magnetohydrodynamics. *J. Comput. Phys.*, 75:400–422, 1988.

[13] R. Broglia, M. Manna, H. Deconinck, and G. Degrez. Development and validation of an axi-symmetric Navier-Stokes solver for hypersonic flows. Technical Report 188, von Karman Institute, 1995.

[14] J. T. Burkepile and O. C. St. Cyr. A revised and expanded catalogue of mass ejections observed by the Solar Maximum Mission coronagraph. Technical Report NCAR/TN-369+STR, National Center for Atmospheric Research, Boulder, Colorado, USA, 1993.

[15] L. Burlaga, R. Fitzenreiter, R. Lepping, K. Ogilvie, A. Szabo, A. Lazarus, J. Steinberg, G. Gloeckler, R. Howard, D. Michels, C. Farrugia, R. P. Lin, and D. E. Larson. A magnetic cloud containing prominence material: January 1997. *J. Geophys. Res.*, 103(A1):277–285, 1998.

[16] S. Cable and Y. Lin. Three-dimensional MHD simulations of interplanetary rotational discontinuities impacting the Earth's bow shock and magnetosheath. *J. Geophys. Res.*, 103(A12):29,551–29,567, 1998.

[17] S. Cable and R. S. Steinolfson. Three-dimensional MHD simulations of the interaction between Venus and the solar wind. *J. Geophys. Res.*, 100(A11):21,645–21,658, 1995.

[18] J. K. Chao. Intermediate shocks: observations. *Adv. Space Res.*, 15(8/9):521–530, 1995.

[19] J. Contopoulos. General axisymmetric magnetohydrodynamic flows: theory and solutions. *ApJ*, 460:185–198, 1996.

[20] R. Courant and K. O. Friedrichs. *Supersonic flow and shock waves.* Interscience publishers, New York, 1948.

[21] R. Courant and D. Hilbert. *Methods of mathematical physics*, volume 2. Interscience publishers, New York, 1962.

[22] A. Csik, H. Deconinck, and S. Poedts. Monotone residual distribution schemes for the ideal 2D magnetohydrodynamic equations on unstructured grids. *AIAA*, paper:99–3325, 1999.

[23] W. Dai and P. R. Woodward. An approximate Riemann solver for ideal magnetohydrodynamics. *J. Comput. Phys.*, 111:354–372, 1994.

[24] W. Dai and P. R. Woodward. Extension of the piecewise parabolic method (PPM) to multidimensional ideal magnetohydrodynamics. *J. Comput. Phys.*, 115:485, 1994.

[25] W. Dai and P. R. Woodward. Interactions between magnetohydrodynamical shocks and denser clouds. *Astrophys. J.*, 436:776–783, 1994.

[26] W. Dai and P. R. Woodward. On the divergence-free condition and conservation laws in numerical simulations for supersonic magnetohydrodynamic flows. *Astrophys. J.*, 494:317–335, 1998.

[27] D.Burgess. Collisionless shocks. In M. G. Kivelson and C. T. Russell, editors, *Introduction to space physics*, pages 129–163. Cambridge University Press, Cambridge, 1995.

[28] C. R. DeVore. Flux-corrected transport techniques for multidimensional compressible magnetohydrodynamics. *J. Comput. Phys.*, 92:142, 1991.

[29] M. A. Dopita, I. Evans, and R. D. Schwartz. Herbig-haro objects 46 and 47 - evidence for bipolar ejection from a young star. *Astrophys. J. Lett.*, 263:L73–L77, 1982.

[30] M. Van Dyke. *An Album of Fluid Motion*. Parabolic Press, Stanford, 1982.

[31] G. E. Brueckner et al. The large angle spectroscopic coronagraph (LASCO). *Solar Phys.*, 162:357–402, 1995.

[32] G. M. Simnett et al. LASCO observations of disconnected magnetic structures out to beyond 28 solar radii during coronal mass ejections. *Solar Phys.*, 175(2), 1997.

[33] C. R. Evans and J. F. Hawley. Simulation of magnetohydrodynamic flows: a constrained transport method. *Astrophys. J.*, 332:659–677, 1988.

[34] S. A. E. G. Falle. Some notes on magnetohydrodynamics. 1994.

[35] S. A. E. G. Falle and S. S. Komissarov. On the inadmissibility of non-evolutionary shocks. *Submitted to J. Plasma Phys.*, 1999.

[36] S. A. E. G. Falle, S. S. Komissarov, and P. Joarder. A multidimensional upwind scheme for magnetohydrodynamics. *Mon. Not. R. Astron. Soc.*, 297:265–277, 1998.

[37] W. C. Feldman, D. N. Baker, S. J. Bame, J. Birn, J. T. Gosling, , Jr. E. W. Hones, and S. J. Schwartz. Slow-mode shocks: a semipermanent feature of the distant geomagnetic tail. *J. Geophys. Res.*, 90(A1):233–240, 1985.

[38] W. C. Feldman, S. J. Schwartz, S. J. Bame, D. N. Baker, J. Birn, J. T. Gosling, Jr. E. W. Hones, J. A. Slavin, E. J. Smith, and R. D. Zwickl. Evidence for slow-mode shocks in the deep geomagnetic tail. *Geophys. Res. Lett.*, 11(6):599–602, 1984.

[39] T. G. Forbes and E. R. Priest. A comparison of analytical and numerical models for steadily driven magnetic reconnection. *Rev. Geophysics*, 25(8):1583–1607, 1987.

[40] H. Freistuehler. Instability of vanishing viscosity approximation to hyperbolic systems of conservation laws with rotational invariance. *J. Differential Equations*, 87(2):205–226, 1990.

[41] H. Freistuehler. Linear degeneracy and shock waves. *Math. Z.*, 207:583–596, 1991.

[42] H. Freistuehler. Rotational degeneracy of hyperbolic systems of conservation laws. *Arch. Rational Mech. Anal.*, 113:39–64, 1991.

[43] H. Freistuehler. Some remarks on the structure of intermediate magnetohydrodynamic shocks. *J. Geophys. Res.*, 96(A3):3825–3827, 1991.

[44] H. Freistuehler. Dynamical stability and vanishing viscosity: a case-study of a non-strictly hyperbolic system. *Comm. Pure. Appl. Math.*, 45:561–582, 1992.

[45] H. Freistuehler. Hyperbolic systems of conservation laws with rotationally equivariant flux function. *Mat. Aplic. Comp.*, 11(2):167–188, 1992.

[46] H. Freistuehler. Contributions to the mathematical theory of magnetohydrodynamic shock waves. In X.-X. Ding and T.-P. Liu, editors, *Nonlinear evolutionary partial differential equations*, volume 3 of *AMS/IP Studies in Advanced Mathematics*, pages 175–187. AMS/IP, 1997.

[47] H. Freistuehler. Small amplitude intermediate magnetohydrodynamic shock waves. *Phys. Scripta*, T74:26–29, 1998.

[48] H. Freistuehler and E. B. Pitman. A numerical study of a rotationally degenerate hyperbolic system. Part 1. The Riemann problem. *J. Comput. Phys.*, 100:306–321, 1992.

[49] H. Freistuehler and P. Szmolyan. Existence and bifurcation of viscous profiles for all intermediate magnetohydrodynamic shock waves. *SIAM J. Math. Anal.*, 26(1):112–128, 1995.

[50] S. E. Gibson and B. C. Low. A time-dependent three-dimensional hydromagnetic model of the coronal mass ejection. *ApJ*, 1998.

[51] S. K. Godunov. Symmetrical form of the magnetohydrodynamic equations. *report of the Computer center of the Siberian branch of the USSR Academy of Sciences*, pages 26–34, 1972.

[52] S. K. Godunov. In C. Carasso, P.-A. Raviart, and D. Serre, editors, *Nonlinear hyperbolic problems: Proceedings of an advanced research workshop*, volume 1270 of *Lecture Notes in Mathematics*, Springer-Verlag, Berlin, 1987.

[53] J. P. Goedbloed and A. Lifshitz. Stationary symmetric magnetohydrodynamic flows. *Phys. Plasmas*, 4(10):3544–3564, 1997.

[54] T. I. Gombosi. *Physics of the space environment.* Atmospheric and space science series. Cambridge University Press, Cambridge, 1998.

[55] T. I. Gombosi, K. G. Powell, and D. L. De Zeeuw. Axisymmetric modeling of cometary mass loading on an adaptively refined grid: MHD results. *J. Geophys. Res.*, 99(A11):21,525–21,539, 1994.

[56] T. I. Gombosi, D. L. De Zeeuw, R. L. Haberli, and K. G. Powell. Three-dimensional multiscale MHD model of cometary plasma environments. *J. Geophys. Res.*, 101:15233–15253, 1996.

[57] J. A. Gonzalez-Esparza, M. Neugebauer, E. J. Smith, and J. L. Phillips. Radial evolution of ejecta characteristics and transient shocks - ulysses in-ecliptic observations. *J. Geophys. Res.*, 103:4767, 1998.

[58] N. Gopalswamy, Y. Hanaoka, T. Kosugi, R. P. Lepping, J. T. Steinberg, S. Plunkett, R. A. Howard, B. J. Thompson, J. Gurman, G. Ho, N. Nitta, and H. S. Hudson. On the relationship between coronal mass ejections and magnetic clouds. *Geophys. Res. Lett.*, 25:2485–2488, 1998.

[59] H. Grad. Reducible problems in magneto-fluid dynamic steady flows. *Revs. Modern Phys.*, 32(4):830–847, 1960.

[60] W. Gropp, E. Lusk, and A. Skjellum. *Using MPI.* Scientific and Engineering Computation. MIT Press, Cambridge, Massachusetts, 1994.

[61] E. Hameiri. The equilibrium and stability of rotating plasmas. *Phys. Fluids*, 26(1):230–237, 1983.

[62] M. Heinemann and S. Olbert. Axisymmetric ideal MHD stellar wind flow. *J. Geophys. Res.*, 83:2457–2460, 1978.

[63] R. D. Henderson. Adaptive spectral element methods for turbulence and transition. In H. Deconinck and T. J. Barth, editors, *High order discretization methods in computational fluid dynamics*, von Karman Institute Lecture Series. 1998.

[64] T. E. Holzer. Interaction between the solar wind and the interstellar medium. *Annu. Rev. Astron. Astrophys.*, 27:199–234, 1989.

[65] G. Hornig. *Zur kovarianten Formulierung der magnetischen Rekonnexion.* PhD thesis, Ruhr-Universität Bochum, 1997.

[66] R. A. Howard, Jr. N. R. Sheely, M. J. Koomen, and D. J. Michels. Coronal mass ejections: 1979-1981. *J. Geophys. Res.*, 90:8173–8191, 1985.

[67] A. J. Hundhausen. The solar wind. In M. G. Kivelson and C. T. Russell, editors, *Introduction to space physics*, pages 91–128. Cambridge University Press, Cambridge, 1995.

[68] A. J. Hundhausen. Coronal mass ejections. In Keith T. Strong, Julia L. R. Saba, Bernhard M. Haisch, and Joan T. Schmelz, editors, *The many faces of the sun*, pages 143–200. Springer-Verlag, New York, 1998.

[69] A. J. Hundhausen, J. T. Burkepile, and O. C. St. Cyr. Speeds of coronal mass ejections: SMM observations from 1980 and 1984-1989. *J. Geophys. Res.*, 99(A4):6543–6552, 1994.

[70] A. Jeffrey. *Magnetohydrodynamics.* Oliver & Boyd, Edinburgh, 1966.

[71] A. Jeffrey and T. Taniuti. *Non-linear wave propagation.* Academic Press, New York, 1964.

[72] H. Ji, M. Yamada, and S. Hsu R. Kulsrud. Experimental test of the Sweet-Parker model of magnetic reconnection. *Phys. Rev. Lett.*, 80(15):3256–3259, 1998.

[73] D. E. Jones, E. J. Smith, J. A. Slavin, B. T. Tsurutani, G. L. Siscoe, and D. A. Mendis. The bow wave of comet Giacobini-Zinner: ICE magnetic field observations. *Geophys. Res. Lett.*, 13(3):243–246, 1986.

[74] K. Kabin, T. I. Gombosi, D. L. De Zeeuw, K. G. Powell, and P. L. Israelevich. Interaction of the Saturnian magnetosphere with Titan: results of a three-dimensional MHD simulation. *To appear in J. Geophys. Res.*, 1999.

[75] M. L. Kaiser, M. J. Reiner, N. Gopalswamy, R. A. Howard, O. C. St. Cyr, B. J. Thompson, and J.-L. Bougeret. Type II radio emissions in the frequency range from 1–14 Mhz associated with the April 7, 1997 solar event. *Geophys. Res. Lett.*, 25(14):2501–2504, 1998.

[76] H. Karimabadi. Physics of intermediate shocks: a review. *Adv. Space Res.*, 15(8/9):507–520, 1995.

[77] H. Karimabadi, D. Krauss-Varban, and N. Omidi. Kinetic structure of intermediate shocks: implications for the magnetopause. *J. Geophys. Res.*, 100(A7):11,957–11,979, 1995.

[78] H. Karimabadi and N. Omidi. Hybrid simulations of intermediate shocks: coplanar and noncoplanar solutions. *Geophys. Res. Lett.*, 19(17):1723–1726, 1992.

[79] Y. Kato and T. Taniuti. Hydromagnetic plane steady flow in compressible ionized gases. *Prog. Theor. Phys.*, 21(4):606–612, 1959.

[80] C. F. Kennel, R. D. Blandford, and P. Coppi. MHD intermediate shock discontinuities. Part 1. Rankine-Hugoniot conditions. *J. Plasma Physics*, 42:299–319, 1989.

[81] C. F. Kennel, R. D. Blandford, and C. C. Wu. Structure and evolution of small-amplitude intermediate shock waves. *Phys. Fluids B*, 2(2):253–269, 1990.

[82] R. Keppens and J. P. Goedbloed. Numerical simulations of stellar winds: polytropic models. *Astron. Astrophys.*, 343:251–260, 1998.

[83] R. Keppens, G. Toth, M. A. Botchev, and A. Van Der Ploeg. Implicit and semi-implicit schemes: algorithms. *Int. J. Numer. Meth. Fluids*, 30:335–352, 1999.

[84] M. G. Kivelson and C. T. Russell. *Introduction to space physics.* Cambridge University Press, Cambridge, 1995.

[85] A. G. Kulikovskiy and G. A. Lyubimov. *Magnetohydrodynamics.* Addison-Wesley, Reading, MA, 1965.

[86] R. M. Kulsrud. Magnetic reconnection in a magnetohydrodynamic plasma. *Phys. Plasmas*, 5(5):1599–1606, 1998.

[87] L. D. Landau and E. M. Lifshitz. *Electrodynamics of continuous media.* Pergamon Press, Oxford, 1984.

[88] P. D. Lax. Hyperbolic systems of conservation laws II. *Comm. Pure Appl. Math.*, 10:537–566, 1957.

[89] L. C. Lee, L. Huang, and J. K. Chao. On the stability of rotational discontinuities and intermediate shocks. *J. Geophys. Res.*, 94(A7):8813–8825, 1989.

[90] R. J. Leveque. *Numerical methods for conservation laws.* Lectures in Mathematics ETH Zurich. Birkhauser-Verlag, Basel, 1992.

[91] R. J. Leveque. Nonlinear conservation laws and finite volume methods for astrophysical fluid flow. In O. Steiner and A. Gautschy, editors, *Computational methods for astrophysical fluid flow.* Springer-Verlag, Berlin, 1998.

[92] M. A. Liberman and A. L. Velikovich. *Physics of shock waves in gases and plasmas*, volume 19 of *Springer Series in Electrophysics.* Springer-Verlag, Berlin, 1985.

[93] A. Lifshitz and J. P. Goedbloed. Transonic magnetohydrodynamic flows. *J. Plasma Physics*, 58(1):61–99, 1997.

[94] Y. Lin and L. C. Lee. Study of the dayside reconnection layer in resistive MHD and hybrid models. *J. Geophys. Res.*, 98(A3):3919–3934, 1993.

[95] Y. Lin and L. C. Lee. Reconnection layer at the flank magnetopause in the presence of shear flow. *Geophys. Res. Lett.*, 21(10):855–858, 1994.

[96] Y. Lin, L. C. Lee, and C. F. Kennel. The role of intermediate shocks in magnetic reconnection. *Geophys. Res. Lett.*, 19(3):229–232, 1992.

[97] Y. Lin, D. W. Swift, and L. C. Lee. Simulation of pressure pulses in the bow shock and magnetosheath driven by variations in interplanetary magnetic field direction. *J. Geophys. Res.*, 101(A12):27,251–27,269, 1996.

[98] Y. Lin and H. Xie. Formation of reconnection layer at the dayside magnetopause. *Geophys. Res. Lett.*, 24(24):3145–3148, 1997.

[99] T. J. Linde. *A three-dimensional adaptive multifluid MHD model of the heliosphere.* PhD thesis, University of Michigan, 1998.

[100] Timur J. Linde, Tamas I. Gombosi, Philip L. Roe, Kenneth G. Powell, and Darren L. De Zeeuw. Heliosphere in the magnetized local interstellar medium: Results of a three-dimensional MHD simulation. *J. Geophys. Res.*, 103(A2):1889–1904, 1998.

[101] H.-L. Liu and M. E. Hagan. Local heating/cooling of the mesosphere due to gravity wave and tidal coupling. *Geophys. Res. Lett.*, 25:2941, 1998.

[102] T.-P. Liu. On the viscosity criterion for hyperbolic conservation laws. In M. Shearer, editor, *Viscous profiles and numerical methods for shock waves*, Siam Proceedings Series, pages 105–114. SIAM, Philadelphia, 1991.

[103] T.-P. Liu and Z. Xin. Overcompressive shock waves. In B. L. Keyfitz and M. Shearer, editors, *Nonlinear evolution equations that change type*, The IMA volumes in mathematics and its applications, pages 139–145. Springer-Verlag, Berlin, 1990.

[104] B. C. Low. Solar activity and the corona. *Solar Phys.*, 167:217–265, 1996.

[105] G. S. S. Ludford. The structure of a hydromagnetic shock in steady plasma motion. *J. Fluid Mech.*, 5:67–80, 1959.

[106] M. Manna. *A three dimensional high resolution compressible flow solver.* PhD thesis, Universite Catholique de Louvain, 1992.

[107] S. A. Markovskii. Nonevolutionary discontinuous magnetohydrodynamic flows in a dissipative medium. *Phys. Plasmas*, 5(7):2596–2604, 1998.

[108] E. Morano, R. Sanders, and M.-C. Druguet. Multidimensional dissipation for upwind schemes: stability and applications to gas dynamics. *Preprint*, 1997.

[109] R. S. Myong and P. L. Roe. Shock waves and rarefaction waves in magnetohydrodynamics. Part 1. A model system. *J. Plasma Physics*, 58:485–519, 1997.

[110] R. S. Myong and P. L. Roe. Shock waves and rarefaction waves in magnetohydrodynamics. Part 2. The MHD system. *J. Plasma Physics*, 58:521–552, 1997.

[111] R. S. Myong and P. L. Roe. On Godunov-type schemes for magnetohydrodynamics. 1. A model system. *J. Comput. Phys.*, 147:545–567, 1998.

[112] N. Omidi, H. Karimabadi, and D. Krauss-Varban. Hybrid simulation of the curved dayside magnetopause during southward IMF. *Geophys. Res. Lett.*, 25(17):3273–3276, 1998.

[113] I. Percival and D. Richards. *Introduction to dynamics.* Cambridge University Press, Cambridge, 1982.

[114] S. M. Petrinec and C. T. Russell. Hydrodynamic and MHD equations across the bow shock and along the surfaces of planetary obstacles. *Space Sci. Rev.*, 79:757–791, 1997.

[115] J. L. Phillips, S. J. Bame, M. F. Thomsen, B. E. Goldstein, and E. J. Smith. Ulysses plasma observations in the Jovian magnetosheath. *J. Geophys. Res.*, 98(A12):21,189–21,202, 1993.

[116] S. Poedts and E. Schwartz. Computation of the ideal-MHD continuous spectrum in axisymmetric plasmas. *Journal of Computational Physics*, 105:165–168, 1993.

[117] N. V. Pogorelov and T. Matsuda. Influence of the interstellar magnetic field direction on the shape of the global heliopause. *J. Geophys. Res.*, 103(A1):237–245, 1998.

[118] K. G. Powell, P. L. Roe, R. S. Myong, T. I. Gombosi, and D. L. De Zeeuw. An upwind scheme for magnetohydrodynamics. *AIAA Paper*, 95-1704-CP, 1995.

[119] E.R. Priest. *Solar Magnetohydrodynamics.* Geophysics and Astrophysics Monographs. D. Reidel Publishing Company, Dordrecht, 1982.

[120] K. B. Quest. Theory and simulation of collisionless parallel shocks. *J. Geophys. Res.*, 93(A9):9649–9680, 1988.

[121] James J. Quirk. A contribution to the great Riemann solver debate. *Int. J. Numer. Methods Fluids*, 18:555–574, 1994.

[122] J. Raeder, J. Berchem, and M. Ashour-Abdalla. The Geospace Environment Modeling grand challenge: results from a Global Geospace Circulation Model. *J. Geophys. Res.*, 103:14,787, 1998.

[123] H. U. Rahman, G. Yur, G. Wong, and R. S. White. Laboratory simulation of the large-scale Birkeland current system in the polar region with northward interplanetary magnetic field. *J. Geophys. Res.*, 94(A6):6873–6878, 1989.

[124] R. Ratkiewicz, A. Barnes, G. A. Molvik, J. R. Spreiter, S. S. Stahara, M. Vinokur, and S. Venkateswaran. Effect of varying strength and orientation of local interstellar magnetic field on configuration of exterior heliosphere: 3D MHD simulations. *Astron. Astrophys.*, 335:363–369, 1998.

[125] M. J. Reiner, M. L. Kaiser, J. Fainberg, and R. G. Stone. A new method for studying remote type ii radio emissions from coronal mass ejection-driven shocks. *J. Geophys. Res.*, 103:29651, 1998.

[126] B. Reipurth and S. Heathcote. The jet and energy source of HH 46/47. *Astron. Astrophys.*, 246:511–534, 1991.

[127] P. L. Roe and D. S. Balsara. Notes on the eigensystem of magnetohydrodynamics. *SIAM J. Appl. Math.*, 56:57–67, 1996.

[128] D. Ryu and T. W. Jones. Numerical magnetohydrodynamics in astrophysics: algorithm and tests for one-dimensional flow. *Astrophys. J.*, 442:228–258, 1995.

[129] D. Ryu, T. W. Jones, and A. Frank. Numerical magnetohydrodynamics in astrophysics: algorithm and tests for multidimensional flow. *Astrophys. J.*, 452:785–796, 1995.

[130] J. Safrankova, Z. Nemecek, L. Prech, G. Zastenker, K. I. Paularena, N. Nikolaeva, M. Nozdrachev, A. Skalsky, and T. Mukai. The January 10–11, 1997 magnetic cloud: multipoint measurements. *Geophys. Res. Lett.*, 25(14):2549–2552, 1998.

[131] T. Sakurai. Magnetohydrodynamic solar/stellar wind models. *Comput. Phys. Rep.*, 12:247, 1990.

[132] T. Sato, T. Hayashi, K. Watanabe, R. Horiuchi, M. Tanaka, N. Sawairi, and K. Kusano. Role of compressibility on driven magnetic reconnection. *Phys. Fluids B*, 4(2):450–457, 1992.

[133] M. Scholer. Asymmetric time-dependent and stationary magnetic reconnection at the dayside magnetopause. *J. Geophys. Res.*, 94(A11):15,099–15,111, 1989.

[134] M. Scholer and R.-F. Lottermoser. On the kinetic structure of the magnetotail reconnection layer. *Geophys. Res. Lett.*, 25(17):3281–3284, 1998.

[135] C. Schwab. hp-fem for fluid flow simulation. In H. Deconinck and T. J. Barth, editors, *High order discretization methods in computational fluid dynamics*, von Karman Institute Lecture Series. 1998.

[136] Jr. Sheeley, N. R., R. A. Howard, D. J. Michels, M. J. Koomen, R. Schwenn, K. H. Muehlhaeuser, and H. Rosenbauer. Coronal mass ejections and interplanetary shocks. *J. Geophys. Res.*, 90:163–175, 1985.

[137] J. Smoller and B. Temple. Shock-wave solutions in closed form and the Oppenheimer–Snyder limit in general relativity. *SIAM Journal on Applied Mathematics*, 58(1):15–33, 1998.

[138] P. Song and C. T. Russell. What do we really know about the magnetosheath. *Adv. Space Res.*, 20(4/5):747–765, 1997.

[139] P. Song, C. T. Russell, T. I. Gombosi, J. R. Spreiter, S. S. Stahara, and X. X. Zhang. On the processes in the terrestrial magnetosheath, 1: scheme development. *To appear in J. Geophys. Res.*, 1999.

[140] P. Song, C. T. Russell, J. T. Gosling, M. F. Thomsen, and R. C. Elphic. Observations of the density profile in the magnetosheath near the stagnation streamline. *Geophys. Res. Lett.*, 17(11):2035–2038, 1990.

[141] P. Song, C. T. Russell, and M. F. Thomsen. Slow mode transition in the frontside magnetosheath. *J. Geophys. Res.*, 97(A6):8295–8305, 1992.

[142] P. Song, C. T. Russell, X. X. Zhang, S. S. Stahara, J. R. Spreiter, and T. I. Gombosi. On the processes in the terrestrial magnetosheath, 2: case study. *To appear in J. Geophys. Res.*, 1999.

[143] J. R. Spreiter and S. S. Stahara. The location of planetary bow shocks: a critical overview of theory and observations. *Adv. Space Res.*, 15(8/9):433–449, 1995.

[144] R. S. Steinolfson and S. Cable. Venus bow shocks at unusually large distances from the planet. *Geophys. Res. Lett.*, 20(8):755–758, 1993.

[145] R. S. Steinolfson and A. J. Hundhausen. Concave-outward slow shocks in coronal mass ejections. *J. Geophys. Res.*, 95:15,251, 1990.

[146] R. S. Steinolfson and A. J. Hundhausen. Coronal mass ejection shock fronts containing the two types of intermediate shocks. *J. Geophys. Res.*, 95(A12):20,693–20,699, 1990.

[147] R. S. Steinolfson and A. J. Hundhausen. MHD intermediate shocks in coronal mass ejections. *J. Geophys. Res.*, 95(A5):6389–6401, 1990.

[148] H. De Sterck, A. Csik, D. Vanden Abeele, S. Poedts, and H. Deconinck. Stationary two-dimensional magnetohydrodynamic flows with shocks: characteristic analysis and grid convergence study. *J. Comput. Phys.*, 2000. in press.

[149] H. De Sterck, H. Deconinck, S. Poedts, and D. Roose. A bow shock flow containing (almost) all types of ('exotic') MHD discontinuities. In *Proceedings of the Seventh International Conference on Hyperbolic Problems*, pages 195–204, Birkhauser-Verlag, Basel, 1998.

[150] H. De Sterck, B. C. Low, and S. Poedts. Complex magnetohydrodynamic bow shock topology in field-aligned low-β flow around a perfectly conducting cylinder. *Phys. Plasmas*, 5(11):4015–4027, 1998.

[151] H. De Sterck, B. C. Low, and S. Poedts. Characteristic analysis of a complex two-dimensional magnetohydrodynamic bow shock flow with steady compound shocks. *Phys. Plasmas*, 6(3):954–969, 1999.

[152] H. De Sterck and S. Poedts. Field-aligned magnetohydrodynamic bow shock flows in the switch-on regime. Parameter study of the flow around a cylinder and results for the axi-symmetrical flow over a sphere. *Astron. Astrophys.*, 343:641–649, 1999.

[153] H. De Sterck and S. Poedts. Stationary slow shocks in the magnetosheath for solar wind conditions with $\beta < 2/\gamma$: Three-dimensional MHD simulations. *J. Geophys. Res.*, 104(A10):22,401, 1999.

[154] H. De Sterck and S. Poedts. Slow shocks and space weather: 3d MHD simulations of the earth's bow shock and magnetosheath. *Submitted to J. Atmos. Solar-Terr. Phys.*, July 1999.

[155] H. De Sterck, S. Poedts, and J. P. Goedbloed. Dynamics of hot filaments in a tokamak plasma. *J. Plasma Physics*, 59/2:277–302, 1998.

[156] E.C. Stone, A.M. Frandsen, R.A. Mewaldt, E.R. Christian, D. Margolies, J.F. Ormes, and F. Snow. The advanced composition explorer. *Space Science Reviews*, 86:1, 1998.

[157] J. M. Stone, J. F. Hawley, C. R. Evans, and M. L. Norman. A test suite for magnetohydrodynamical simulations. *Astrophys. J.*, 388:415–437, 1992.

[158] J. M. Stone and M. L. Norman. ZEUS-2D: A radiation magnetohydrodynamics code for astrophysical flows in two space dimensions.

ii. the magnetohydrodynamic algorithms and tests. *Astrophys. J. Suppl.*, 80:791–818, 1992.

[159] T. Tanaka. Configurations of the solar wind flow and magnetic field around the planets with no magnetic field: calculation by a new MHD simulation scheme. *J. Geophys. Res.*, 98(A10):17,251–17,262, 1993.

[160] P. A. Thompson. *Compressible-fluid dynamics.* McGraw-Hill, New York, 1972.

[161] G. Toth, R. Keppens, and M. A. Botchev. Implicit and semi-implicit schemes in the Versatile Advection Code: numerical tests. *Astron. Astrophys.*, 332:1159–1170, 1998.

[162] G. Toth and D. Odstrcil. Comparison of some Flux Corrected Transport and Total Variation Diminishing numerical schemes for hydrodynamic and magnetohydrodynamic problems. *J. Comput. Phys.*, 128:82, 1996.

[163] K. Tsinganos, C. Sauty, G. Surlantzis, E. Trussoni, and J. Contopoulos. On the relation of limiting characteristics to critical surfaces in magnetohydrodynamic flows. *Mon. Not. R. Astron. Soc.*, 283:811–820, 1996.

[164] E. Turkel. Accuracy of schemes with non-uniform meshes for compressible fluid flows. Technical Report 85-43, ICASE Langley, VA, 1985.

[165] Y. Uchida, Y. Todo, R. Rosner, and K. Shibata. A magnetohydrodynamic model for Herwig-Haro objects: magnetically guided shocked flows associated with optical jets from young stellar objects. *Publ. Astron. Soc. Japan*, 44:227–243, 1992.

[166] S. Umemura, K. Iki, K. Shibata, and Y. Sofue. MHD shock waves through a galactic nuclear disk with a vertical magnetic field. *Publ. Astron. Soc. Japan*, 40:25–46, 1988.

[167] E. van der Weide. *Compressible flow simulation on unstructured grids using multi-dimensional upwind schemes.* PhD thesis, Technische Universiteit Delft, 1998.

[168] R. J. Walker and C. T. Russell. Solar-wind interactions with magnetized planets. In M. G. Kivelson and C. T. Russell, editors, *Introduction to space physics*, pages 164–182. Cambridge University Press, Cambridge, 1995.

[169] G. B. Whitham. *Linear and nonlinear waves.* Wiley-Interscience. John Wiley, New York, 1974.

[170] C. C. Wu. On MHD intermediate shocks. *Geophys. Res. Lett.*, 14(6):668–671, 1987.

[171] C. C. Wu. The MHD intermediate shock interaction with an intermediate wave: are intermediate shocks physical? *J. Geophys. Res.*, 93(A2):987–990, 1988.

[172] C. C. Wu. Formation, structure and stability of MHD intermediate shocks. *J. Geophys. Res.*, 95:8149, 1990.

[173] C. C. Wu. New theory of MHD shock waves. In M. Shearer, editor, *Viscous profiles and numerical methods for shock waves*, Siam Proceedings Series, pages 209–236. SIAM, Philadelphia, 1991.

[174] C. C. Wu. MHD flow past an obstacle: large-scale flow in the magnetosheath. *Geophys. Res. Lett.*, 19(2):87–90, 1992.

[175] C. C. Wu. Magnetohydrodynamic Riemann problem and the structure of the magnetic reconnection layer. *J. Geophys. Res.*, 100(A04):5579–5598, 1995.

[176] C. C. Wu and T. Hada. Formation of intermediate shocks in both two-fluid and hybrid models. *J. Geophys. Res.*, 96(A3):3769–3778, 1991.

[177] C. C. Wu and C. F. Kennel. Structural relations for time-dependent intermediate shocks. *Geophys. Res. Lett.*, 19(20):2087–2090, 1992.

[178] C. C. Wu and C. F. Kennel. Structure and evolution of time-dependent intermediate shocks. *Phys. Rev. Lett.*, 68(1):56–59, 1992.

[179] M. Yamada, H. Ji, S. Hsu, T. Carter, R. Kulsrud, N. Bretz, F. Jobes, Y. Ono, and F. Perkins. Study of driven magnetic reconnection in a laboratory plasma. *Phys. Plasmas*, 4(5):1936–1944, 1997.

[180] M. Yan and L. C. Lee. Generation of slow-mode waves in front of the dayside magnetopause. *Geophys. Res. Lett.*, 21(7):629–632, 1994.

[181] M. Yokosawa. Formation of shock wave in magnetohydrodynamical accretion onto a black hole. *Publ. Astron. Soc. Japan*, 46:73–86, 1994.

[182] T. Yokoyama and K. Shibata. Numerical simulation of solar coronal X-ray jets based on the magnetic reconnection model. *Publ. Astron. Soc. Japan*, 48:353–376, 1996.

[183] A. L. Zachary, A. Malagoli, and P. Colella. A higher-order method for multidimensional ideal magnetohydrodynamics. *SIAM J. Sci. Comput.*, 15(2):263–284, 1994.

[184] K Zumbrun and D. Serre. Viscous and inviscid stability of multi-dimensional planar shock fronts. *Preprint*, 1999.