

## A CORRIGENDUM TO “D-GROUPS AND THE DME”

REMI JAQUI AND RAHIM MOOSA

The purpose of this note, which is not intended for publication, is to fill a small gap that appears in the proof of Proposition 3.5 of [1]. At the bottom of page 364 of the published version there is a mistake: when addressing the case of  $n = 1$ ; that is, the case when  $a$  and  $\log \delta a := \frac{\delta a}{a}$  are algebraically dependent over  $k$ ; it is claimed that something forces the polynomial  $P_0(x)$  to be trivial. In fact,  $P_0(x) = d(1 - x)$  for any  $d \in k$  is possible, and should have been dealt with.

Instead of following the details in [1] we will give a direct and more conceptual proof of Proposition 3.5 of [1] in this special case:

**Proposition.** *Suppose  $k$  is a field of characteristic zero,  $R$  is a commutative affine Hopf  $k$ -algebra, and  $\delta$  is a  $k$ -linear derivation on  $R$  that is an  $a$ -coderivation for some group-like  $a \in R$ . If  $a$  and  $\log \delta a$  are algebraically dependent over  $k$  then  $\log \delta a = d(1 - a)$  for some  $d \in k$ .*

*In particular, the conclusion of [1, Proposition 3.5] holds with  $c := -\frac{d^2}{2}$ .*

*Proof.* There is an affine algebraic group  $G$ , a nontrivial character  $a : G \rightarrow \mathbb{G}_m$ , and an  $a$ -twisted  $D$ -group structure  $s : G \rightarrow TG$ , all over  $k$ , such that  $R = k[G]$  and  $\delta$  is the derivation on  $k[G]$  induced by  $s$ .

Consider the map  $\pi := (a, \log \delta a) : G \rightarrow \mathbb{G}_m \times \mathbb{G}_a$  that appears in [1] where it is shown, by an easy computation at the beginning of the proof of Proposition 3.8, that, since  $a$  is a character and  $s$  is  $a$ -twisted,  $\pi$  is a morphism of algebraic groups. Since  $\{a, \log \delta a\}$  is algebraically dependent over  $k$ , and  $a \neq 1$ , the (connected) algebraic subgroup  $H := \pi(G) \leq \mathbb{G}_m \times \mathbb{G}_a$  must be 1-dimensional. Hence the coordinate projection  $\pi_1 : H \rightarrow \mathbb{G}_m$  is surjective with finite kernel. It follows that  $\pi_1$  factors as  $\rho_n \phi$ , for some  $n > 0$ , where  $\phi : H \rightarrow \mathbb{G}_m$  is an isomorphism of algebraic groups and  $\rho_n : \mathbb{G}_m \rightarrow \mathbb{G}_m$  is given by  $\rho_n(x) = x^n$ . Let  $\mathbb{G}_m \times_n \mathbb{G}_a$  be the semidirect product where  $\mathbb{G}_m$  acts on  $\mathbb{G}_a$  by  $(x, y) \mapsto x^n y$ .

We first claim that  $F : H \rightarrow \mathbb{G}_m \times_n \mathbb{G}_a$  given by  $(x, y) \mapsto (\phi(x, y), y)$ , is a group homomorphism. Indeed,

$$\begin{aligned}
 F((x, y)(x', y')) &= F(xx', y + xy') \\
 &= (\phi(xx', y + xy'), y + xy') \\
 &= (\phi(x, y)\phi(x', y'), y + xy') \quad \text{as } \phi \text{ is a group homomorphism} \\
 &= (\phi(x, y)\phi(x', y'), y + \phi(x, y)^n y') \quad \text{as } \phi(x, y)^n = x \\
 &= (\phi(x, y), y)(\phi(x', y'), y') \\
 &= F(x, y)F(x', y')
 \end{aligned}$$

as desired.

Next, we claim that  $\chi := \pi_2 \phi^{-1} : \mathbb{G}_m \rightarrow \mathbb{G}_a$  is a  $\rho_n$ -twisted additive character. That is, that  $\chi(xx') = \chi(x) + x^n \chi(x')$ . Indeed, note that the image  $H' = F(H)$  is

the graph of  $\chi$ . So for  $x, x' \in \mathbb{G}_m$  we have  $(xx', \chi(xx')) \in H'$ . But as  $F$  is a group homomorphism,  $H'$  is a subgroup of  $\mathbb{G}_m \times_n \mathbb{G}_a$ , and so

$$(x, \chi(x))(x', \chi(x')) = (xx', \chi(x) + x^n \chi(x')) \in H',$$

as well. It follows that  $\chi(xx') = \chi(x) + x^n \chi(x')$ , as desired.

Choose  $b \in k$  such that  $b^n \neq 1$ . We have, for  $x \in \mathbb{G}_m$ ,

$$\begin{aligned} \chi(xb) &= \chi(x) + x^n \chi(b) & \text{and} \\ \chi(bx) &= \chi(b) + b^n \chi(x) \end{aligned}$$

so that  $\chi(x)(1 - b^n) = \chi(b)(1 - x^n)$ . Letting  $d := \frac{\chi(b)}{1 - b^n}$  we get  $\chi(x) = d(1 - x^n)$ .

So, for all  $g \in G$ , we have

$$\begin{aligned} \log \delta a(g) &= \pi_2 \pi(g) \\ &= \chi \phi \pi(g) & \text{as } \chi = \pi_2 \phi^{-1} \\ &= d(1 - \phi \pi(g)^n) \\ &= d(1 - \pi_1 \pi(g)) & \text{as } \pi_1 = \rho_n \phi \\ &= d(1 - a(g)). \end{aligned}$$

That is,  $\log \delta a = d(1 - a)$ , as desired.

For the ‘‘in particular’’ clause, a direct computation shows that for  $c := -\frac{d^2}{2}$  we have the identity  $a\delta^2 a = \frac{3}{2}(\delta a)^2 + c(a^2 - a^4)$  as claimed in Proposition 3.5 of [1].  $\square$

It may be worth pointing out that what  $\log \delta a = d(1 - a)$  says geometrically is that we have the short exact sequence

$$1 \longrightarrow (\ker(a), u) \longrightarrow (G, s) \xrightarrow{a} (\mathbb{G}_m, t_a) \longrightarrow 1$$

where  $u := s|_{\ker(a)}$  makes  $(\ker(a), u)$  a  $D$ -group,  $t_a(x) := d(x - x^2)$  makes  $(\mathbb{G}_m, t_a)$  an id-twisted  $D$ -group, and the morphisms are algebraic group homomorphisms that are also morphisms of  $D$ -varieties.

## REFERENCES

- [1] Jason Bell, Omar Le3n S3nchez, and Rahim Moosa.  $D$ -groups and the Dixmier-Moeglin equivalence. *Algebra Number Theory*, 12(2):343–378, 2018.