

SYDE Advanced Math 2, Solutions to Assignment 5

1: (a) Solve the system $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ and draw the direction field and some solution curves.

Solution: Let $A = \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix}$. Then $\det(A - rI) = \det \begin{pmatrix} 1-r & 2 \\ 1 & -r \end{pmatrix} = r^2 - r - 2 = (r+1)(r-2)$.


When $r = -1$ we have $A - rI = \begin{pmatrix} 2 & 2 \\ 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$ and so an eigenvector is $\mathbf{u} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$.

When $r = 2$ we have $A - rI = \begin{pmatrix} -1 & 2 \\ 1 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & -2 \\ 0 & 0 \end{pmatrix}$ so an eigenvector is $\mathbf{v} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$.

The solution to the system is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^{-t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + be^{2t} \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$

To sketch the direction field, note that the isoclines are given by $m = \frac{y'}{x'} = \frac{x}{x+2y}$, that is $mx + 2my = x$, or equivalently $y = \frac{1-m}{2m}x$. This is the line through $(0,0)$ with slope $\frac{1-m}{2m}$. Some isoclines are shown in yellow, the direction field is shown in green, and some solution curves are shown in blue.



(b) Solve the system $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ and draw the direction field and some solution curves.

Solution: Let $A = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix}$. Then $\det(A - rI) = \det \begin{pmatrix} 1-r & 2 \\ -2 & 1-r \end{pmatrix} = r^2 - 2r + 5 = (r-1)^2 + 4$.

The eigenvalues are $r = 1 \pm 2i$.

When $r = 1 + 2i$ we have $A - rI = \begin{pmatrix} -2i & 2 \\ -2 & -2i \end{pmatrix} \sim \begin{pmatrix} -i & 1 \\ 0 & 0 \end{pmatrix}$ so an eigenvector is $\mathbf{u} = \begin{pmatrix} 1 \\ i \end{pmatrix}$.

A complex solution is

$$\begin{pmatrix} z \\ w \end{pmatrix} = e^{(1+2i)t} \begin{pmatrix} 1 \\ i \end{pmatrix} = e^t (\cos 2t + i \sin 2t) \begin{pmatrix} 1 \\ i \end{pmatrix} = e^t \begin{pmatrix} \cos 2t + i \sin 2t \\ -\sin 2t + i \cos 2t \end{pmatrix}.$$


Two independent real solutions are obtained from the real and imaginary parts of the above complex solution.

The general solution is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^t \begin{pmatrix} \cos 2t \\ -\sin 2t \end{pmatrix} + be^t \begin{pmatrix} \sin 2t \\ \cos 2t \end{pmatrix}.$$

The isoclines are given by $m = \frac{y'}{x'} = \frac{-2x+y}{x+2y}$, that is $mx + 2my = -2x + y$, or equivalently $y = \frac{2+m}{1-2m}x$.

This is the line through $(0,0)$ of slope $\frac{2+m}{1-2m}$. Some isoclines are shown in yellow, the slope field is shown in green, and some solution curves are shown in blue.



2: (a) Find the solution to the system $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ with $\begin{pmatrix} x(0) \\ y(0) \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$.

Solution: Let $A = \begin{pmatrix} 1 & -2 \\ 2 & -3 \end{pmatrix}$. Then $\det(A - rI) = \det \begin{pmatrix} 1-r & -2 \\ 2 & -3-r \end{pmatrix} = r^2 + 2r + 1 = (r+1)^2$.

When $r = -1$ we have $A - rI = \begin{pmatrix} 2 & -2 \\ 2 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$ so an eigenvector is $\mathbf{u} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$.

Note also that when $\mathbf{v} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ we have $(A - rI)\mathbf{v} = \begin{pmatrix} 2 & -2 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} = 2\mathbf{u}$.

Thus the general solution to the system is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^{-t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + be^{-t} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + bte^{-t} \begin{pmatrix} 2 \\ 2 \end{pmatrix}.$$

To get $x(0) = 2$ we need $2 = a + b$ and to get $y(0) = 1$ we need $1 = a$, so we must take $a = 1$ and $b = 1$, and we obtain the solution

$$\begin{pmatrix} x \\ y \end{pmatrix} = e^{-t} \begin{pmatrix} 2 \\ 1 \end{pmatrix} + te^{-t} \begin{pmatrix} 2 \\ 2 \end{pmatrix}$$

(b) Find the solution to the system $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ 4 & 5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ with $\begin{pmatrix} x(0) \\ y(0) \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

Solution: Let $A = \begin{pmatrix} 1 & -2 \\ 4 & 5 \end{pmatrix}$. Then $\det(A - rI) = \det \begin{pmatrix} 1-r & -2 \\ 4 & 5-r \end{pmatrix} = r^2 - 6r + 13 = (r-3)^2 + 4$,

The eigenvalues are $r = 3 \pm 2i$.

When $r = 3 + 2i$ we have $A - rI = \begin{pmatrix} -2-2i & -2 \\ 4 & 2-2i \end{pmatrix} \sim \begin{pmatrix} 2 & 1-i \\ 0 & 0 \end{pmatrix}$ so an eigenvector is $\mathbf{u} = \begin{pmatrix} 1-i \\ -2 \end{pmatrix}$.

A complex solution is

$$\begin{pmatrix} z \\ w \end{pmatrix} = e^{(3+2i)t} \begin{pmatrix} 1-i \\ -2 \end{pmatrix} = e^{3t} (\cos 2t + i \sin 2t) \begin{pmatrix} 1-i \\ -2 \end{pmatrix} = e^{3t} \begin{pmatrix} (\cos 2t + \sin 2t) + i(\sin 2t - \cos 2t) \\ -2 \cos 2t - 2i \sin 2t \end{pmatrix}.$$

Two independent real solutions are given by the real and imaginary parts of this complex solution, so the general solution to the system is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^{3t} \begin{pmatrix} \sin 2t + \cos 2t \\ -2 \cos 2t \end{pmatrix} + be^{3t} \begin{pmatrix} \sin 2t - \cos 2t \\ -2 \sin 2t \end{pmatrix}.$$

To get $x(0) = 1$ we need $1 = a - b$ and to get $y(0) = 2$ we need $2 = -2a$, so we must have $a = -1$ and $b = -2$, and so the solution is

$$\begin{pmatrix} x \\ y \end{pmatrix} = -e^{3t} \begin{pmatrix} \sin 2t + \cos 2t \\ -2 \cos 2t \end{pmatrix} - 2e^{3t} \begin{pmatrix} \sin 2t - \cos 2t \\ -2 \sin 2t \end{pmatrix} = e^{3t} \begin{pmatrix} \cos 2t - 3 \sin 2t \\ 4 \sin 2t + 2 \cos 2t \end{pmatrix}.$$

3: (a) Solve the system $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 6t+1 \\ e^t \end{pmatrix}$.

Solution: First we solve the associated homogeneous system. Let $A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$.

Then $\det(A - rI) = \det \begin{pmatrix} 1-r & 2 \\ -1 & 4-r \end{pmatrix} = r^2 - 5r + 6 = (r-2)(r-3)$ so the eigenvalues are $r = 2, 3$.

When $r = 2$ we have $A - rI = \begin{pmatrix} -1 & 2 \\ -1 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & -2 \\ 0 & 0 \end{pmatrix}$ so an eigenvector is $\mathbf{u} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$.

When $r = 3$ we have $A - rI = \begin{pmatrix} -2 & 2 \\ -1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix}$ so an eigenvector is $\mathbf{v} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$.

Thus the general solution to the homogeneous system is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^{2t} \begin{pmatrix} 2 \\ 1 \end{pmatrix} + be^{3t} \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

We shall find a particular solution to the given non-homogeneous system using variation of parameters.

We try $\begin{pmatrix} x_p \\ y_p \end{pmatrix} = X \begin{pmatrix} r \\ s \end{pmatrix}$ where $X = \begin{pmatrix} 2e^{2t} & e^{3t} \\ e^{2t} & e^{3t} \end{pmatrix}$. Putting this in the non-homogeneous system gives

$$\begin{pmatrix} r' \\ s' \end{pmatrix} = \begin{pmatrix} 2e^{2t} & e^{3t} \\ e^{2t} & e^{3t} \end{pmatrix}^{-1} \begin{pmatrix} 6t+1 \\ e^t \end{pmatrix} = e^{-5t} \begin{pmatrix} e^{3t} & -e^{3t} \\ -e^{2t} & 2e^{2t} \end{pmatrix} \begin{pmatrix} 6t+1 \\ e^t \end{pmatrix}$$

$$= e^{-5t} \begin{pmatrix} (6t+1)e^{3t} - e^{4t} \\ -(6t+1)e^{2t} + 2e^{3t} \end{pmatrix} = \begin{pmatrix} (6t+1)e^{-2t} - e^{-t} \\ -(6t+1)e^{-3t} + 2e^{-2t} \end{pmatrix}.$$

We integrate (using integration by parts) to obtain

$$r = \int (6t+1)e^{-2t} - e^{-t} dt = -\frac{1}{2}(6t+1)e^{-2t} - \frac{3}{2}e^{-2t} + e^{-t} = -(3t+2)e^{-2t} + e^{-t}, \text{ and}$$

$$s = \int -(6t+1)e^{-3t} + 2e^{-2t} dt = \frac{1}{3}(6t+1)e^{-3t} + \frac{2}{3}e^{-3t} - e^{-t} = (2t+1)e^{-3t} - e^{-2t}.$$

Thus we obtain the particular solution

$$\begin{pmatrix} x_p \\ y_p \end{pmatrix} = \begin{pmatrix} 2e^{2t} & e^{3t} \\ e^{2t} & e^{3t} \end{pmatrix} \begin{pmatrix} -(3t+2)e^{-2t} + e^{-t} \\ (2t+1)e^{-3t} - e^{-2t} \end{pmatrix} = \begin{pmatrix} -2(3t+2) + 2e^t + (2t+1) - e^t \\ -(3t+2) + e^t + (2t+1) - e^t \end{pmatrix} = \begin{pmatrix} -4t-3+e^t \\ -t-1 \end{pmatrix}.$$

The general solution to the given system is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^{2t} \begin{pmatrix} 2 \\ 1 \end{pmatrix} + be^{3t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} -4t-3+e^t \\ -t-1 \end{pmatrix}.$$

(b) Solve the system $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 3 & -4 \\ 5 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1 \\ 5e^{3t} - 4 \end{pmatrix}$.

Solution: First we solve the associated homogeneous system. Let $A = \begin{pmatrix} 3 & -4 \\ 5 & -1 \end{pmatrix}$.

Then $\det(A - rI) = \det \begin{pmatrix} 3-r & -4 \\ 5 & -1-r \end{pmatrix} = r^2 - 2r + 17 = (r-1)^2 + 16$, so the eigenvalues are $r = 1 \pm 4i$.

When $r = 1+4i$ we have $A - rI = \begin{pmatrix} 2-4i & -4 \\ 5 & -2-4i \end{pmatrix} \sim \begin{pmatrix} 1-2i & -2 \\ 0 & 0 \end{pmatrix}$ so an eigenvector is $\mathbf{u} = \begin{pmatrix} 2 \\ 1-2i \end{pmatrix}$.
A complex solution is

$$\begin{pmatrix} z \\ w \end{pmatrix} = e^{(1+4i)t} \begin{pmatrix} 2 \\ 1-2i \end{pmatrix} = e^t (\cos 4t + i \sin 4t) \begin{pmatrix} 2 \\ 1-2i \end{pmatrix} = e^t \begin{pmatrix} 2 \cos 4t + i \sin 4t \\ (2 \sin 4t + \cos 4t) + i(\sin 4t - 2 \cos 4t) \end{pmatrix}$$

so the general solution to the associated homogeneous system is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^t \begin{pmatrix} 2 \cos 4t \\ 2 \sin 4t + \cos 4t \end{pmatrix} + be^t \begin{pmatrix} 2 \sin 4t \\ \sin 4t - 2 \cos 4t \end{pmatrix}.$$

We shall find a particular solution to the non-homogeneous system using the method of undetermined coefficients. We try $\begin{pmatrix} x_p \\ y_p \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix} + e^{3t} \begin{pmatrix} r \\ s \end{pmatrix}$. Put this into the system of DEs. The left side is

$$LS = \begin{pmatrix} x_p' \\ y_p' \end{pmatrix} = \begin{pmatrix} 3re^{3t} \\ 3se^{3t} \end{pmatrix}$$

and the right side is

$$\begin{aligned} RS &= \begin{pmatrix} 3 & -4 \\ 5 & -1 \end{pmatrix} \begin{pmatrix} x_p \\ y_p \end{pmatrix} + \begin{pmatrix} 1 \\ 5e^{3t} - 4 \end{pmatrix} = \begin{pmatrix} 3 & -4 \\ 5 & -1 \end{pmatrix} \begin{pmatrix} c + re^{3t} \\ d + se^{3t} \end{pmatrix} + \begin{pmatrix} 1 \\ 5e^{3t} - 4 \end{pmatrix} \\ &= \begin{pmatrix} 3c + 3re^{3t} - 4d - 4se^{3t} + 1 \\ 5c + 5re^{3t} - d - se^{3t} + 5e^{3t} - 4 \end{pmatrix}. \end{aligned}$$

In order to have $LS = RS$ we need $3re^{3t} = 3c + 3re^{3t} - 4d - 4se^{3t} + 1 = 0$, that is $(3c - 4d + 1) + (-4s)e^{3t} = 0$, and $3se^{3t} = 5c + 5re^{3t} - d - se^{3t} + 5e^{3t} - 4$, that is $(5c - d - 4) + (5r - 4s + 5)e^{3t} = 0$. Since 1 and e^{3t} are linearly independent, the coefficients must all vanish, so we have $3c - 4d + 1 = 0$, $5c - d - 4 = 0$, $-4s = 0$ and $5r - 4s + 5 = 0$. The first two of these equations give $c = 1$ and $d = 1$ and the second two give $r = -1$ and $s = 0$. Thus we obtain the particular solution

$$\begin{pmatrix} x_p \\ y_p \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} + e^{3t} \begin{pmatrix} -1 \\ 0 \end{pmatrix}.$$

The general solution to the given system is

$$\begin{pmatrix} x \\ y \end{pmatrix} = ae^t \begin{pmatrix} 2 \cos 4t \\ 2 \sin 4t + \cos 4t \end{pmatrix} + be^t \begin{pmatrix} 2 \sin 4t \\ \sin 4t - 2 \cos 4t \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} + e^{3t} \begin{pmatrix} -1 \\ 0 \end{pmatrix}.$$

4: Find the solution to the system $\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 2 & -2 & 1 \\ 1 & -1 & 1 \\ 2 & -4 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ with $\begin{pmatrix} x(0) \\ y(0) \\ z(0) \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$.

Solution: Let $A = \begin{pmatrix} 2 & -2 & 1 \\ 1 & -1 & 1 \\ 2 & -4 & 3 \end{pmatrix}$. Then

$$\begin{aligned} \det(A - rI) &= \det \begin{pmatrix} 2-r & -2 & 1 \\ 1 & -1-r & 1 \\ 2 & -4 & 3-r \end{pmatrix} \\ &= -(r-2)(r-3)(r+1) - 4 - 4 - 4(r-2) - 2(r-3) + 2(r+1) \\ &= -(r-2)(r-3)(r+1) - 4r + 8 = -(r-2)((r-3)(r+1) + 4) \\ &= -(r-2)(r^2 - 2r + 1) = -(r-2)(r-1)^2. \end{aligned}$$

When $r = 2$ we have

$$A - rI = \begin{pmatrix} 0 & -2 & 1 \\ 1 & -3 & 1 \\ 2 & -4 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & -3 & 1 \\ 0 & -2 & 1 \\ 0 & 2 & -1 \end{pmatrix} \sim \begin{pmatrix} 1 & -3 & 1 \\ 0 & 1 & -\frac{1}{2} \\ 0 & 2 & -1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & -\frac{1}{2} \\ 0 & 0 & 0 \end{pmatrix}$$

so an eigenvector is $\mathbf{u} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$. When $r = 1$ we have

$$A - rI = \begin{pmatrix} 1 & -2 & 1 \\ 1 & -2 & 1 \\ 2 & -4 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & -2 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

so we have the two independent eigenvectors $\mathbf{v} = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$ and $\mathbf{w} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$. Thus the general solution to the system is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = ae^{2t} \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} + be^t \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + ce^t \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}.$$

To get $x(0) = 1$ we need $a + b + 2c = 1$, to get $y(0) = 2$ we need $a + 0b + c = 2$, and to get $z(0) = 1$ we need $2a - b + 0c = 1$. We solve these three equations:

$$\left(\begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 1 & 0 & 1 & 2 \\ 2 & -1 & 0 & 1 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 0 & 1 & 1 & -1 \\ 0 & 3 & 4 & 1 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & 1 & 4 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -5 \\ 0 & 0 & 1 & 4 \end{array} \right).$$

Thus $a = -2$, $b = -5$ and $c = 4$, so the solution is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = -2e^{2t} \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} - 5e^t \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + 4e^t \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} = -2e^{2t} \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} + e^t \begin{pmatrix} 3 \\ 4 \\ 5 \end{pmatrix}.$$

5: Solve the system $\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 1 & 1 & -2 \\ -2 & -2 & 2 \\ 3 & 2 & -3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} + e^{-2t} \begin{pmatrix} -1 \\ 2 \\ 2 \end{pmatrix}$.

Solution: Let $A = \begin{pmatrix} 1 & 1 & -2 \\ -2 & -2 & 2 \\ 3 & 2 & -3 \end{pmatrix}$. Then

$$\begin{aligned} \det(A - rI) &= \det \begin{pmatrix} 1-r & 1 & -2 \\ -2 & -2-r & 2 \\ 3 & 2 & -3-r \end{pmatrix} \\ &= -(r-1)(r+2)(r+3) + 6 + 8 + 4(r-1) - 2(r+3) - 6(r+2) \\ &= -(r-1)(r+2)(r+3) - 4r - 8 = -(r+2)((r-1)(r+3) + 4) \\ &= -(r+2)(r^2 + 2r + 1) = -(r+2)(r+1)^2. \end{aligned}$$

When $r = -2$ we have

$$A - rI = \begin{pmatrix} 3 & 1 & -2 \\ -2 & 0 & 2 \\ 3 & 2 & -1 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 & 0 \\ -2 & 0 & 2 \\ 3 & 2 & -1 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

so an eigenvector is $\mathbf{u} = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$. When $r = -1$ we have

$$A - rI = \begin{pmatrix} 2 & 1 & -2 \\ -2 & -1 & 2 \\ 3 & 2 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 & 0 \\ -2 & -1 & 2 \\ 3 & 2 & -2 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 1 & 2 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

so one eigenvector is $\mathbf{v} = \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$. We look for a vector \mathbf{w} such that $(A - rI)\mathbf{w} = \mathbf{v}$:

$$\left(\begin{array}{ccc|c} 2 & 1 & -2 & 2 \\ -2 & -1 & 2 & -2 \\ 3 & 2 & -2 & 1 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 0 & -1 \\ -2 & -1 & 2 & -2 \\ 3 & 2 & -2 & 1 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 0 & -1 \\ 0 & 1 & 2 & -4 \\ 0 & 1 & 2 & -4 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & -2 & 3 \\ 0 & 1 & 2 & -4 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

and we obtain $\mathbf{w} = \begin{pmatrix} 3 \\ -4 \\ 0 \end{pmatrix}$. Thus the general solution to the associated homogeneous system is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = ae^{-2t} \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} + be^{-t} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} + ce^{-t} \begin{pmatrix} 3 \\ -4 \\ 0 \end{pmatrix} + cte^{-t} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}.$$

To find a particular solution to the given non-homogeneous system, we use the method of undetermined coefficients. We try $\mathbf{x} = e^{-2t}\mathbf{p} + te^{-2t}\mathbf{q}$, where $\mathbf{x} = (x_p, y_p, z_p)^T$ and \mathbf{p} and \mathbf{q} are vectors in \mathbb{R}^3 . We put \mathbf{x} into the non-homogeneous system of DEs. The left side is

$$LS = \mathbf{x}' = -2e^{-2t}\mathbf{p} + e^{-2t}\mathbf{q} - 2te^{-2t}\mathbf{q} = e^{-2t}(-2\mathbf{p} + \mathbf{q}) + te^{-2t}(-2\mathbf{q})$$

and, writing $\mathbf{b} = (-1, 2, 2)^T$, the right side is

$$RS = A\mathbf{x} + e^{-2t}\mathbf{b} = A(e^{-2t}\mathbf{p} + te^{-2t}\mathbf{q}) + e^{-2t}\mathbf{b} = e^{-2t}(A\mathbf{p} + \mathbf{b}) + te^{-2t}A\mathbf{q}.$$

To get $LS = RS$ we need $-2\mathbf{p} + \mathbf{q} = A\mathbf{p} + \mathbf{b}$ and $-2\mathbf{q} = A\mathbf{q}$. Note that $-2\mathbf{q} = A\mathbf{q} \iff (A + 2I)\mathbf{q} = 0$ so \mathbf{q} must be an eigenvector of $r = -2$, and so we must have $\mathbf{q} = k\mathbf{u}$ for some $k \in \mathbb{R}$. Also, note that $-2\mathbf{p} + \mathbf{q} = A\mathbf{p} + \mathbf{b} \iff (A + 2I)\mathbf{p} = \mathbf{q} - \mathbf{b}$, so we solve this to find \mathbf{p} :

$$\begin{array}{c} \left(\begin{array}{ccc|c} 3 & 1 & -2 & k+1 \\ -2 & 0 & 2 & -k-2 \\ 3 & 2 & -1 & k-2 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 0 & -1 \\ -2 & 0 & 2 & -k-2 \\ 3 & 2 & -1 & k-2 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 0 & -1 \\ 0 & 2 & 2 & -k-4 \\ 0 & 1 & 1 & -k-1 \end{array} \right) \\ \sim \left(\begin{array}{ccc|c} 1 & 1 & 0 & -1 \\ 0 & 1 & 1 & -k-1 \\ 0 & 2 & 2 & -k-4 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & -1 & k \\ 0 & 1 & 1 & -k-1 \\ 0 & 0 & 0 & k-2 \end{array} \right) \end{array}$$

To get a solution, we must take $k = 2$ and we have $\mathbf{q} = k\mathbf{u} = \begin{pmatrix} 2 \\ -2 \\ 2 \end{pmatrix}$, and $\mathbf{p} = \begin{pmatrix} k \\ -k-1 \\ k-2 \end{pmatrix} = \begin{pmatrix} 2 \\ -3 \\ 0 \end{pmatrix}$.

Thus we obtain the particular solution

$$\begin{pmatrix} x_p \\ y_p \\ z_p \end{pmatrix} = e^{-2t} \begin{pmatrix} 2 \\ -3 \\ 0 \end{pmatrix} + te^{-2t} \begin{pmatrix} 2 \\ -2 \\ 2 \end{pmatrix}.$$

The general solution to the given (non-homogeneous) system is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = ae^{-2t} \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} + be^{-t} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} + ce^{-t} \begin{pmatrix} 3 \\ -4 \\ 0 \end{pmatrix} + cte^{-t} \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} + e^{-2t} \begin{pmatrix} 2 \\ -3 \\ 0 \end{pmatrix} + te^{-2t} \begin{pmatrix} 2 \\ -2 \\ 2 \end{pmatrix}.$$