


PARADOXES: Simpson's Paradox

1. Background Review – Lurking Variates, Confounding and Comparison Error [optional reading]

As background to answering Question(s) about an \mathbf{X} - \mathbf{Y} relationship between a focal variate \mathbf{X} and a response variate \mathbf{Y} , $\mathbf{Z}_1, \mathbf{Z}_2, \dots, \mathbf{Z}_k$ in the schema at the right are called **lurking variates**, a phrase that means lurking *explanatory* variates in that each \mathbf{Z} accounts, at least in part, for changes from element to element in the value of the response variate. The importance of lurking variates is that if the distributions of their values *differ* between groups of elements [like (sub)populations or samples] with different values of the focal variate, an Answer about the \mathbf{X} - \mathbf{Y} relationship may differ from the true state of affairs unless the differences in the values of relevant \mathbf{Z} s are taken into account.



The *same* statistical issue raised by lurking variates is involved, with different terminology, in **confounding**; the difference is that the behaviour of lurking variates (the entity responsible) is *why* confounding (the statistical issue) occurs.

An explanatory variate responsible for confounding is called a **confounder** or **confounding variate**; these two terms are synonyms for a lurking variate whose distribution of values (over a group of elements) differs for different values of the focal variate.

The following definitions summarize the foregoing discussion:

- * **Lurking variate:** a non-focal explanatory variate (\mathbf{Z}) whose differing distributions of values over groups of elements or units with different values of the focal variate, if taken into account, would meaningfully change an Answer about an \mathbf{X} - \mathbf{Y} relationship.
- * **Confounding:** differing distributions of values of one or more *non-focal* explanatory variate(s) among two (or more) groups of elements or units [like (sub)populations or samples] with different values of the focal variate.
 - **Confounder (confounding variate):** a non-focal explanatory variate involved in confounding.

'Confounding' and 'confounder' have the convenience of being one-word terminology rather than the multi-word phrases involving 'lurking variates' which convey the same ideas.

- * **Comparison error:** for an Answer about an \mathbf{X} - \mathbf{Y} relationship that is based on comparing attributes of groups of elements or units with different values of the focal variate(s), comparison error is the difference from the *intended* (or *true*) state of affairs arising from:
 - differing distributions of lurking variate values between (or among) the groups of elements or units OR – confounding.


The alternate wording of the last phrase accommodates the equivalent terminologies of lurking variates and confounding; in a particular context, we use the version of the definition appropriate to that context:

- 'lurking variates' can more readily accommodate phenomena like Simpson's Paradox discussed in this Highlight #51;
- 'confounding' is more common in the context of comparative Plans, as illustrated in Statistical Highlight #63, but the variety of usage of 'confounding' can be a source of difficulty (see Statistical Highlight #3).

Comparison error in experimental and observational Plans is discussed in Statistical Highlight #9.


The schema at the right below (from, for instance, page HL6.4 in Statistical Highlight #6) reminds us of several matters.

- Data-based investigating is concerned initially with four groups of elements or units – the *target* population, the *study* population, the *respondent* population and the *sample*.
- Associated with each of these groups are (one or more) *attributes* of interest.
- Answer(s) to Question(s) are usually given in terms of attributes, often their values.
- Our six categories of error, of which five are defined in terms of attributes – model error is the exception.
 - In the schema, the four arrows arising from comparison error point to *boxes* representing *groups* of elements or units (a population or a sample) rather than, as for the other five error categories, to *lines joining boxes*; the rightmost head of these four arrows is to be taken as pointing to *both* sample ellipses.
 - + Multiple arrows for comparison error are a consequence of its different manifestations in different Question contexts, as illustrated in Table HL60.2 on the lower half of page HL60.4 in Statistical Highlight #60.



In other discussion (e.g., in Statistical Highlights #57 and #63), the context for comparison error due to lurking variate(s)/confounding is comparative investigating of a *treatment* effect; the relevant *causal* structure (e.g., from the upper half of page HL59.1 in Statistical Highlight #59), is case (8), shown at the upper right, with focal variate \mathbf{X} , response variate \mathbf{Y} and lurking variate/confounder \mathbf{Z} . In this Highlight #51, as summarized in the structure (A)₂ at the lower right, we broaden the discussion in two ways:

- we have two (or three) 'focal' variates [not necessarily all of equal interest in the Question context];



- we are *unconcerned* with *causation* as the reason for the \mathbf{X}_1 - \mathbf{Y} and \mathbf{Z} - \mathbf{Y} associations, because the nature of the focal variates is such that we *cannot* set their levels and this precludes using such focal variate(s) to manipulate the value of \mathbf{Y} ;
 – this is why the lower structure at the right overleaf on page HL51.1 has *lines rather than arrows* between the variate symbols.

The phenomenon known as Simpson's Paradox can arise in a comparative investigation where the attributes are proportions – that is, the response variate \mathbf{Y} is *qualitative* [discrete (categorical)] in nature; the dramatic name ('Paradox') is a reflection of how the effect of lurking variate(s) can *reverse* the sign of a relationship. The data in twenty of the first twenty-one Tables HL51.1 to HL51.21 used in discussion of Simpson's Paradox in this Highlight #51 are hypothetical. The discussion is in six sections:

2. Illustrations of Simpson's Paradox
3. 'Simpson's Paradox' with a quantitative response variate.
4. Reasons for Simpson's Paradox – properties of proportions.
5. Reasons for Simpson's Paradox – population subgroups and weighted averages.
6. Reasons for Simpson's Paradox – probability distributions.
7. A Plan for an investigation to answer the Question of sex discrimination.


The discussion is framed in terms of *populations*, because there are no *inherent* sampling issues in Simpson's Paradox; when the groups being compared are *samples*, there is the additional statistical issue of managing sample error.

2. Illustrations of Simpson's Paradox

The data in Table HL51.1 below come from the discussion of Simpson's Paradox in Program 11 of *Against All Odds: Inside Statistics*; the context is possible sex discrimination in graduate admissions. Overall, the admission rate [or *proportion* (an *attribute*)] is *lower* for women (50% vs. 55% for men – see the bottom line of the Table) but, when the data are subdivided by school (Law and Business), the female admission rate is *higher* (by 5 percentage points) for *each* school. The (binary) response variate is school admission (Yes, No) and the lurking variate is women-to-men ratio among applicants; its effect is because:

- * the two schools had appreciably *different* admission rates: 80 and 75% for Law, 20 and 15% for Business;
- * *half* as many women as men (120 vs. 240) applied to Law but *equal* numbers of women and men (120) applied to Business.

SCHOOLWOMEN.....		MEN.....		
	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	240	180	75
Business	120	24	20	120	18	15
Both	240	120	51	360	198	55



The diagram to the right of Table HL51.1 shows its data in graphical form; Simpson's Paradox is the *positive* slope of the middle dashed line for the data for *both* schools changing to a *negative* slope in the upper and lower lines for the schools *individually*:

In this illustration, the variates in the lower structure (A_2) at the lower right overleaf on page HL51.1 are:

\mathbf{X}_1 is an applicant's sex (female, male), \mathbf{X}_2 is the school applied to (Law, Business),

[In Tables HL51.5 and HL51.6 on the facing page HL51.3, \mathbf{X}_3 is the level of study (Masters, Doctoral)],

\mathbf{Z} is the (lurking variate) women-to-men ratio among applicants (discussed further in Sections 3 and 5 on pages HL51.3 and HL51.4),


\mathbf{Y} is the response to an applicant (admitted, not admitted). [On page HL51.3, \mathbf{Y} is time for degree completion (minimum, longer).]

Unlike investigating a treatment effect when there is more than one focal variate (e.g., using a factorial treatment structure), the focal variate of primary interest in *this* Question context is \mathbf{X}_1 , an applicant's sex.


The limitation imposed by lurking variates on an Answer to a Question about an \mathbf{X} - \mathbf{Y} relationship is illustrated further by the data in Tables HL51.2 to HL51.4; as the diagrams to the right of the tables emphasize, it is also possible to have:

- * the *same* overall admission rate for women and men but a *higher* rate for women in the two schools individually (Table HL51.2);

SCHOOLWOMEN.....		MEN.....		
	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	168	126	75
Business	120	24	20	120	18	15
Both	240	120	50	288	144	50




SCHOOLWOMEN.....		MEN.....		
	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	240	192	80
Business	120	24	20	120	24	20
Both	240	120	51	360	216	60



- * a *lower* overall admission rate for women but the *same* rate for women and men in the two schools individually (Table HL51.3);
- * a *higher* rate overall *and* in the two schools individually for women (Table HL51.4).

The effect of lurking variates on an \mathbf{X} - \mathbf{Y} relationship at a *second* level of subdivision is illustrated in Tables HL51.5 and HL51.6 at the upper right of the facing page HL51.3; a context

SCHOOLWOMEN.....		MEN.....		
	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	120	90	75
Business	120	24	20	120	18	15
Both	240	120	51	240	108	45




(continued)

PARADOXES: Simpson's Paradox (continued 1)


for these data is the proportion of graduate students who complete their degree in the minimum time. In Table HL51.5, the proportion for women is *lower* overall, *higher* when subdivided by subject area (Law or Business) but again *lower* when subject area is subdivided by level (Masters or Doctoral). Similar effects are seen in Table HL51.6, except the proportions for women become *equal* when subdivided by subject area and *higher* when further subdivided by level.

Probabilistically, subdividing is *conditioning* so that Tables HL51.1 to HL51.6, in illustrating Simpson's Paradox, show the limitation on an Answer which involves comparing *conditional* probabilities for a response variate with *different* conditionings; that is, comparing probabilities for Y given X_1 and X_2 with Y given only X_1 (in Tables HL51.1 to HL51.4) or for Y given X_1 , X_2 and X_3 with Y given X_1 and X_2 or Y given only X_1 (in Tables HL51.5 and HL51.6) – see Section 6 overleaf on page HL51.4. Four other illustrations of Simpson's Paradox are given in Note 2 on pages HL51.5 and HL51.6 and three more illustrative tables (like Table HL51.9 overleaf on page HL51.4) are discussed on pages HL51.6 and HL51.7 in the Appendix of this Highlight #51.

SCHOOLWOMEN.....		MEN.....		
	Number of Students	COMPLETIONS Number	%	Number of Students	COMPLETIONS Number	%
Law: Masters	60	51	85	60	54	90
Doctoral	60	33	55	300	180	60
Bus.: Masters	60	27	45	20	10	50
Doctoral	60	9	15	100	20	20
Law	120	84	70	360	234	65
Business	120	36	30	120	30	25
Both	240	120	50	480	264	55




SCHOOLWOMEN.....		MEN.....		
	Number of Students	COMPLETIONS Number	%	Number of Students	COMPLETIONS Number	%
Law: Masters	60	54	90	240	204	85
Doctoral	60	42	70	80	52	65
Bus.: Masters	60	18	30	120	30	25
Doctoral	60	6	10	40	2	5
Law	120	96	80	320	256	80
Business	120	24	20	160	32	20
Both	240	120	51	480	288	60



3. 'Simpson's Paradox' with a Quantitative Response Variate

Simpson's Paradox is usually presented in the context of comparing *proportions* but the same phenomenon can occur with a *continuous* response variate. As illustrated by the data in Table HL51.7 and the diagram to its right, whose context is graduate studies admission averages, the average is *lower* overall for women than men (84% vs. 86%) but, when the data are subdivided by school, both averages are *higher* for women. The response variate here is an applicant's average, the attribute is the *average* of these averages (e.g., 90 and 88 for Law, 82 and 80 for Business) and the lurking variate is women-to-men ratio among applicants (1:3 for Law, 3:1 for Business). With 1:1 ratios, there is *no* 'paradox'.

SCHOOLWOMEN.....	MEN.....	
	Number of Applicants	Applicants' Average (%)	Number of Applicants	Applicants' Average (%)
Law	50	90	150	88
Business	150	82	50	80
Both	200	84	200	86




The illustration in Table HL51.7 shows that Simpson's Paradox is *not* solely a phenomenon which may arise when comparing *proportions*. Its origin lies in the *relative* 'natural' group sizes arising from the process of subdividing (or its inverse of combining) used to manage comparison error in observational Plans. Such a lurking variate (called the women-to-men ratio in the discussion of Table HL51.1 on page HL51.2 and Table HL51.7 above) is *different* in nature to Z in the upper causal structure of case (8) at the lower right of page HL51.1, which we think of as being able to *cause* an element to change the value of its response variate. Thus, we now recognize *two* ways a change in a lurking variate can affect attribute value(s):

- by *causing* elements' response variate (and, hence, their attribute) values to change, OR:
- by distorting attribute *calculation* when subdividing is used to manage comparison error in an observational Plan.

4. Reasons for Simpson's Paradox – properties of proportions

Quantities (like variate and attribute values) which are *single* numbers are relatively straight-forward to compare: 4 is greater than 2 is greater than -6, although the latter has a larger *magnitude* than the first two. However, when quantities (like proportions or fractions and the coordinates of points on a scatter diagram) involve *two* numbers, comparisons may raise complications. For example, in the diagram at the right, points A and C with *different* coordinates are the *same* distance from the origin and point B is *closer* to the origin than A and C despite its coordinates being *larger* than one of those of A and C. The (surprising) result for fractions (or proportions), exhibited in



(continued overleaf)

Simpson's Paradox, is that for eight (positive) integers a, b, \dots, h , it is possible to have (as in Table HL51.1 on page HL51.2):

$$\frac{a}{b} > \frac{c}{d} \text{ and } \frac{e}{f} > \frac{g}{h} \text{ but at the same time to have: } \frac{a+e}{b+f} < \frac{c+g}{d+h}, \quad \text{e.g., } \frac{32}{40} > \frac{90}{120} \text{ and } \frac{12}{60} > \frac{9}{60} \text{ but } \frac{44}{100} < \frac{99}{180}.$$

This property also applies to *more* than two pairs of fractions (as in Table HL51.15 on page HL51.5) and for other combinations of inequality and equality (as in Tables HL51.2, HL51.3, HL51.5 and HL51.6 on page HL51.2 and overleaf on page HL51.3)

When the fraction $\frac{90}{120}$ is instead $\frac{30}{40}$, we see at the right that there is *no* 'paradox' (as in Table HL51.4), reminding us that it is the group sizes (in the denominators) under subdividing that may engage the property of proportions which generates the 'paradox' – recall Section 3 on page HL51.3.

5. Reasons for Simpson's Paradox – population subgroups and weighted averages

The distorted calculation of the values of (population) attributes (like proportions and averages), which generates the 'paradox' illustrated in Sections 2 and 3, is an instance of *weighted* combinations of the corresponding attributes of population *subgroups*. As shown in Table HL51.8 at the right, the attribute values in the last line of each of Tables HL51.1 to HL51.4 are weighted combinations of the attributes in the two table lines above them; what produces the changes in attribute values *relative* to each other is a change in *weights*. Each weight is determined by the (natural) *size* of a population subgroup; this size is the *lurking* variate whose change is responsible for the change in (the sign of) the \mathbf{X} - \mathbf{Y} relationship. The same idea applies to *each* of the two levels of subdivision in Tables HL51.5 and HL51.6 and to the averages in Table HL51.7. When the weights are *equal* (as in Table HL51.4), there is *no* 'paradox'.

Table HL51.8: Weighted percentage

	Weights	
Table HL51.1:	$\frac{120}{240} \times 80 + \frac{120}{240} \times 20 = 50$	$\frac{1}{2} \quad \frac{1}{2}$
	$\frac{240}{360} \times 75 + \frac{120}{360} \times 15 = 55$	$\frac{2}{3} \quad \frac{1}{3}$
Table HL51.2:	$\frac{120}{240} \times 80 + \frac{120}{240} \times 20 = 50$	$\frac{1}{2} \quad \frac{1}{2}$
	$\frac{168}{288} \times 75 + \frac{120}{288} \times 15 = 50$	$\frac{7}{12} \quad \frac{5}{12}$
Table HL51.3:	$\frac{120}{240} \times 80 + \frac{120}{240} \times 20 = 50$	$\frac{1}{2} \quad \frac{1}{2}$
	$\frac{240}{360} \times 80 + \frac{120}{360} \times 20 = 60$	$\frac{2}{3} \quad \frac{1}{3}$
Table HL51.4:	$\frac{120}{240} \times 80 + \frac{120}{240} \times 20 = 50$	$\frac{1}{2} \quad \frac{1}{2}$
	$\frac{120}{240} \times 75 + \frac{120}{240} \times 15 = 45$	$\frac{1}{2} \quad \frac{1}{2}$

6. Reasons for Simpson's Paradox – probability distributions

Table HL51.1 on page HL51.2 provides data from which the probability function of a discrete trivariate distribution can be estimated. To obtain this model, we first extend Table HL51.1 as in Table HL51.9 below to include three extra columns for 'Both sexes.' We then define five events and use estimates for ten probabilities – the vertical line means 'given that' in the eight *conditional* probabilities and \cap denotes an *intersection* of events.

SCHOOLWOMEN.....		MEN.....		BOTH SEXES.....		
	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	240	180	75	360	276	76.6
Business	120	24	20	120	18	15	240	42	17.5
Both schools	240	120	50	360	198	55	600	318	53

The (joint) trivariate model is shown in Table HL51.10 at the right below; summing its probabilities for one variate, we obtain the three

Event A: Applicant is admitted (\mathbf{Y} =yes; the complement $\bar{\mathbf{A}}$ is \mathbf{Y} =no)

Event F: Applicant is female (\mathbf{X}_1 =female)

Event M: Applicant is male (\mathbf{X}_1 =male)

Event L: Applicant applies to Law (\mathbf{X}_2 =Law)

Event B: Applicant applies to Business (\mathbf{X}_2 =Business)

$$\Pr(\mathbf{F}) = 0.4$$

$$\Pr(\mathbf{M}) = 0.6$$

$$\Pr(\mathbf{A}|\mathbf{F}) = 0.5$$

$$\Pr(\mathbf{A}|\mathbf{M}) = 0.55$$

$$\Pr(\mathbf{A}|\mathbf{L}) = 0.76$$

$$\Pr(\mathbf{A}|\mathbf{B}) = 0.175$$

$$\Pr(\mathbf{A}|\mathbf{F} \cap \mathbf{L}) = 0.8$$

$$\Pr(\mathbf{A}|\mathbf{M} \cap \mathbf{L}) = 0.2$$

$$\Pr(\mathbf{A}|\mathbf{M} \cap \mathbf{B}) = 0.15$$

(marginal) *bivariate* models in Tables HL51.11 to HL51.13. The smaller **bold** annotations in Tables HL51.10 to HL51.12 show how eight of the nine percentages in Table HL51.9 arise; for example, the 80% of women admitted to Law is $\frac{0.16}{0.2}$.

Table HL51.10: Trivariate model for \mathbf{Y} , \mathbf{X}_1 and \mathbf{X}_2

F.....	M.....		
	L	B	L	B	
A	0.16	0.04	0.3	0.03	0.53
$\bar{\mathbf{A}}$	0.8	0.2	0.75	0.15	0.47
	0.2	0.2	0.4	0.2	

Table HL51.11: Bivariate model for \mathbf{Y} and \mathbf{X}_1

	F	M	
A	0.2	0.33	0.53
$\bar{\mathbf{A}}$	0.8	0.67	0.47
	0.4	0.6	

Table HL51.12: Bivariate model for \mathbf{Y} and \mathbf{X}_2

	L	B	
A	0.46	0.07	0.53
$\bar{\mathbf{A}}$	0.54	0.93	0.47
	0.6	0.4	

Table HL51.13: Bivariate model for \mathbf{X}_1 and \mathbf{X}_2

	L	B	
F	0.2	0.2	0.4
M	0.4	0.2	0.6
	0.6	0.4	

We see that Table HL51.1 on page HL51.2 involves *parts* of the two multivariate distributions in Tables HL51.10 and HL51.11; it is therefore *unsurprising* if comparisons among these parts, taken in isolation, yield seeming 'paradoxes'. It can be confusing that Table HL51.1 and those like it do not show *explicitly* percentages involving *complements* [like applicants 'not admitted' (event $\bar{\mathbf{A}}$)].

7. A Plan for an Investigation to Answer the Question of Sex Discrimination

Comparing proportions of women and men admitted among applicants to graduate studies (as in the context of Table HL51.1 on page HL51.2) is *not* an adequate Plan to answer the Question of possible sex discrimination, for two reasons:

- there is the possibility of Simpson's Paradox and no clear way to define the level of subdivision at which to make comparisons;
- applicants' *qualifications* are not taken into account.

Both matters are addressed by a Plan which involves taking *pairs* of applicants, one female and one male, with the *same* qualifications for admission and then comparing the proportions of women and men who are admitted across a number of such


PARADOXES: Simpson's Paradox (continued 2)

pairs that is adequate, in the investigation context, to manage all relevant categories of error.

* For comparison error, pairing manages the *group sizes* (and hence, the weights in the attribute calculations) in a way that precludes Simpson's Paradox; matching manages equality of qualifications for the groups of women and men being compared. However, as with *any* observational Plan (that gathers data from a population in its *natural* state), there is still the limitation on Answer(s) imposed by comparison error due to other (unrecognized) lurking variates.

* When investigating the much-discussed issue of comparable worth (whether women are paid the same as men for the same work), relevant explanatory variates to manage include qualifications, experience and hours worked per month or per year.

The schema at the right is a pictorial reminder of the lurking variate of group (population or sample) sizes when developing an observational Plan to answer a Question with a *causative* aspect, which (usually) involves comparing attribute values (calculated or obtained from a scatter diagram in the *Analysis* stage of the FDEAC cycle) for broad subpopulations (like women and men). By contrast, when answering a Question with a *descriptive* aspect (e.g., a Question about *both* sexes), differing attribute values at different levels of subdivision are more obvious and so lurking variate(s) are usually less troublesome. These matters are illustrated, using information from Table HL51.5 (near the top of page HL51.3), in Table HL51.14 at the right below.



NOTES: 1. Simpson's Paradox is so surprising, particularly when first encountered, that it is easy to lose sight of key *statistical* issues.

- The proportions *are* correctly calculated – Simpson's Paradox is *not* the result of mistakes in arithmetic.

- Simpson's Paradox is *not* confined to attributes that are proportions (as discussed in Section 3 on page HL51.3).

- Simpson's Paradox occurs when subdividing (or combining) data for categories and only in *some* circumstances.

Lessons for data-based investigating are:

- * recognize and manage the (surprising) property of proportions discussed in Section 4 on pages HL51.3 and HL51.4;
- * manage relevant non-focal explanatory variates – this includes sometimes being able to identify an appropriate level of subdivision at which to make comparisons (as in Table HL63.3 on page HL63.4 of Statistical Highlight #63).

There is then no 'paradox' for a *clear* Question investigated with an adequate Plan, suggesting that the name Simpson's *Paradox* can be misleading;

2. Four more illustrations of Simpson's Paradox are:

Table HL51.15: The context is the same as that of Table HL51.1 on page HL51.1 but there are now *six* programs (A, ..., F) instead of two schools (Law, Business).

Like Table HL51.1, there is a *lower* percentage of women admitted overall but a *higher* percentage for *each* of the six programs.

Table HL51.16: Baseball batting averages – the batter with the *lower* average for the whole season has a *higher* average in both *half* seasons. Recalling Section 7 and Note 1 above, it is of interest to develop a Plan to answer the Question of which batter to take if only one can be chosen.

Table HL51.17: Death rates (per 1,000 lives) in two regions of the U.S. for smokers and non-smokers.

[These data were gathered by a life insurance company which was issuing whole life policies countrywide on a non-medical issue basis; in 1986, 3,800 policies were issued to males aged 40-45. The company's files were kept in two locations – Nashville for policies issued east of the Mississippi and Los Angeles for policies issued west of the Mississippi. Nashville issued 2,000 policies and processed 13 deaths, Los Angeles issued 1,800 policies and processed 8 deaths.]

REFERENCE: Dolins, J.G.: Actuaries ... be careful! *The Actuary*, March, 1989, page 11.

Table HL51.14: (Based on Table HL51.5 data)		Women		Men		Both sexes	
Group		Group size	%	Group size	%	Group size	%
Individuals		1	--	1	--	1	--
Smaller subgroups	Law: Masters	60	85	60	90	120	88
	Doctoral	60	55	300	60	360	59
	Bus.: Masters	60	45	20	50	80	46
	Doctoral	60	15	100	20	160	18
Larger subgroups	Law	120	70	360	65	480	66
	Business	120	30	120	25	240	27
Population		240	50	480	55	720	53

PROGRAMWOMEN.....		MEN.....		
	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Archeology	108	89	82	825	512	62
Biology	25	17	68	560	353	63
Chemistry	593	219	37	325	114	35
Drama	375	131	35	417	138	33
English	393	106	27	191	48	25
French	341	27	8	373	22	6
All	1,825	589	32	2,691	1,187	44

Time PeriodBATTER #1.....		BATTER #2.....		
	Hits	At bats	Average	Hits	At bats	Average
First half	15	70	.214	25	130	.192
Second half	15	50	.300	80	280	.286
Whole season	30	120	.250	105	410	.256

LOCATIONSMOKERS.....			NON-SMOKERS		
	Deaths	Policies	Rate	Deaths	Policies	Rate
Nashville	6	900	6.67	7	1,100	6.36
Los Angeles	5	1,100	4.55	3	700	4.29
Either	11	2,000	5.50	10	1,800	5.56

(continued overleaf)

NOTES: 2. **Table HL51.18:** Effect of jury challenges on conviction rates in trials in the U.K.

[In early 1987, an article by Bernard Levin in *The Times* raised the question of whether jury challenges assist those who are guilty in avoiding conviction.

Mr. Levin concluded this was *not* the case, on the

basis of data showing a conviction rate of 53% in trials with no challenges, *lower* than the conviction rate of 60% in trials *with* challenges. However, this answer does *not* necessarily follow from these conviction rates; in the *hypothetical* data in Table HL51.18 (at the right above), the conviction rate for *guilty* defendants is substantially *higher* in trials with *no* challenges. Unfortunately, this counter-argument is speculative because the number of defendants *actually* guilty and innocent, and the rates of challenge and of conviction in both these groups, are not readily accessible. Nevertheless, an article in a major newspaper which uses flawed reasoning from data to answer a Question on a substantive issue is a serious matter.]

REFERENCE: Hill, I.D.: Rebutting the media. *The Royal Statistical Society NEWS & NOTES* 16(#1), September, 1989, page 4.

There is discussion and further illustrations of Simpson's Paradox in Wagner, C.H.: Simpson's Paradox in Real Life. *American Statistician* 36 (#1, February): 46-48 (1982).

DEFENDENT STATUS	...NO CHALLENGE...		CHALLENGE.....		
	Number of Trials	CONVICTIONS Number	%	Number of Trials	CONVICTIONS Number	%
Guilty	20	16	80	70	42	60
Innocent	10	0	0	0	0	0
Either	30	16	53	70	42	60

- ① Referring to the data in Table HL51.17 overleaf at the lower right of page HL51.5, suggest a plausible explanation for the *lower* death rates for both smokers and non-smokers whose files were kept in Los Angeles, compared with those kept in Nashville.
- ② Referring to the matters raised by the numbers in Table HL51.18 at the right above, outline how you would try to *reduce* the uncertainties which are present and so obtain an Answer with fewer limitations about the effect(s) of jury challenges on conviction rates for the guilty in the U.K.
 - What effect(s) of jury challenges on the conviction of *innocent* defendants is indicated by the numbers in Table HL51.18? Explain briefly.

8. Appendix: Simpson's Paradox and Interaction

For extending the discussion of Simpson's Paradox on the first six sides (pages HL51.1 to HL51.6) of this Highlight #51, for convenience in this Appendix (including labelling the three diagrams to the right of Tables HL51.19 to HL51.21 below and on the facing page HL51.7) we use the notation defined near the middle of page HL51.2:


X_1 is an applicant's sex (female, male), X_2 is the school applied to (Law, Business), X_3 is the level of study [Masters, Doctoral], Y is the response to an applicant (admitted, not admitted) or time for degree completion (minimum, longer), \bar{Y} (the *average* of Y) is the *percentage* of applicants admitted or who complete their degree in the minimum time.

The diagrams illustrating Simpson's Paradox to the right of Tables HL51.1 to HL51.6 (on pages HL51.2 and HL51.3) are reminiscent of a diagram showing interaction (e.g., in Note 6 on page HL68.2 in Statistical Highlight #68); however, there are *differences*:

- the Simpson's Paradox diagrams have an additional (dashed) line for the overall X_1 - \bar{Y} relationship;
- the instances of Simpson's Paradox in Tables HL51.1 to HL51.6 have only *parallel* (solid) lines for the X_1 - \bar{Y} relationships for different values of X_2 – that is, there is *no* interaction of X_1 and X_2 in their effects on Y .

This restriction is *removed* in (another) reworking of Table HL51.1 and its diagram in Table HL51.19 below, where there *is* interaction of X_1 and X_2 in their effects on Y because the two solid lines in the diagram to the right of the Table are *not* parallel.

SCHOOLWOMEN.....		MEN.....		BOTH SEXES.....		
	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	480	360	75	600	456	76
Business	120	36	30	120	12	10	240	48	20
Both schools	240	132	55	600	372	62			



Thus, interaction *may* be involved in Simpson's Paradox but is not *required* for it to occur.

Earlier discussion at the upper left of page HL51.3 and on page HL51.4 in Section 6, and in this Appendix, reminds us that Simpson's Paradox and interaction *both* involve (estimated) values of *conditional* probabilities for Y , BUT:

- Simpson's Paradox involves comparing these probabilities conditioned on two (or three) of the X s with probabilities conditioned on one *fewer* (one or two) X s; WHEREAS:
- interaction is absent or present depending on the values of probabilities with the *same* conditioning on the X s – these values determine whether the corresponding lines are or are not parallel.

NOTES: 3. Illustration of Simpson's Paradox from comparing *across* Tables HL51.1 to HL51.6 can overshadow comparisons *down* such tables. For example, in Table HL51.1 (reworked as Table HL51.9 on page HL51.4), the six **bold** percentages for X_2 (80 and 20, 75 and 15, 76.6 and 17.5) address a Question *different* from possible sex discrimination:

PARADOXES: Simpson's Paradox (continued 3)

NOTES: 3. ● How do the admission standards of the Law and Business schools compare?
(cont.)

The (hypothetical) data in Table HL51.9 on page HL51.4 (and Tables HL51.19 to HL51.21 on the facing page HL51.6 and below on this page) indicate an appreciably *higher* admission standard for Business than for Law, unless the abilities of the two applicant pools are remarkably different.

4. A second reworking of Table HL51.1 and its diagram is given below in Table HL51.20; as in Table HL51.19, there *is* interaction of X_1 and X_2 in their effects on Y but, for both schools combined, there is *no* sex difference in proportions, due to cancellation of effects in *opposite* directions for the schools individually.

Table HL51.20:

WOMEN.....		MEN.....		BOTH SEXES.....		
SCHOOL	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	180	153	85	300	249	83
Business	120	24	20	180	27	15	300	51	17
Both schools	240	120	50	360	180	50			





Table HL51.21 below and its diagram show, like Table HL51.20, *no* sex difference for both schools combined but this is now a consequence of the *individual schools* also showing this same behaviour – there is *no* interaction.

Table HL51.21:

WOMEN.....		MEN.....		BOTH SEXES.....		
SCHOOL	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%	Number of Applicants	ADMISSIONS Number	%
Law	120	96	80	80	64	80	200	160	80
Business	120	24	20	80	16	20	200	40	20
Both schools	240	120	50	160	80	50			



5. Across Tables HL51.1 to HL51.6 on pages HL51.2 and HL51.3 and Tables HL51.19 to HL51.21 on the facing page HL51.6 and above, different weights in the proportion calculations (like those in Table HL51.8 on page HL51.4) yield a noteworthy *variety* in the percentages for women compared to those for men. This is summarized in Table HL51.22 at the right below; three categories are distinguished.

- In four tables, there *is* an X_1 - \bar{Y} relationship, there is no interaction of X_1 and X_2 in their effects on Y and, in two of the tables, the X_1 - \bar{Y} relationship is *unexceptional* in light of the effect of subdivision by X_2 ; by contrast, in Table HL51.3 and Table HL51.6 between the first and second levels of subdivision by X_2 , the exceptional behaviour is the X_1 - \bar{Y} relationship *disappearing* when the data are subdivided by X_2 .

- Notation like (1,3) [or (1,2)] on Table HL51.5 (or Table HL51.6) in Table HL51.22 refers to the first and third (or first and second) levels of subdivision by X_2 .

- In four tables, there is *no* X_1 - \bar{Y} relationship but three of these are 'false negative' Answers – when the data are subdivided by X_2 , there *is* an X_1 - \bar{Y} relationship and so they are designated 'exceptional' in the fourth column.

- In Table HL51.20, interaction is the *reason* for the exceptional behaviour but interaction is absent in the other three tables.

- In five tables, there *is* (again) an X_1 - \bar{Y} relationship, interaction is absent or incidental, and each is a case of the exceptional behaviour known as Simpson's Paradox (SP).


Apart from understanding the properties of proportions and weighted averages and using an adequate Plan (discussed in Sections 4, 5 and 7 and Note 1 on pages HL51.3 to HL51.5), the summary in Table HL51.22 above reminds us that:

- ⊙ Simpson's Paradox is merely the *most* exceptional case (*change of direction* of an X_1 - \bar{Y} relationship) in a context (involving *discrete* variates) that can give rise to less exceptional or even *unexceptional* behaviour;
- ⊙ interaction is rarely the reason for the exceptional behaviour (only in Table HL51.20 above).

Table HL51.22: X_1 - \bar{Y} RELATIONSHIPS IN NINE TABLES
(SP in the fourth column denotes 'Simpson's Paradox')

Table	Relationship	Interaction	Exceptional behaviour
HL51.3	Yes	No	Yes
HL51.4	Yes	No	No
HL51.5 (1,3)	Yes	No	No
HL51.6 (1,2)	Yes	No	Yes
HL51.2	No	No	Yes
HL51.6 (2,3)	No	No	Yes
HL51.20	No	Essential	Yes
HL51.21	No	No	No
HL51.1	Yes	No	Yes: SP
HL51.5 (1,2)	Yes	No	Yes: SP
HL51.5 (2,3)	Yes	No	Yes: SP
HL51.6 (1,3)	Yes	No	Yes: SP
HL51.19	Yes	Incidental	Yes: SP

6. In meeting the obligation to deal with *relationships* in an introductory statistics course, the lengthy discussion (e.g., in Statistical Highlights #9, #10 and #57 to #70) shows the (unexpected) complexities, for only *three* variates, arising from issues of causation, confounding, interaction and Simpson's Paradox. The schema at the right reminds us there are common themes *and* differences among these four matters – see also Appendix 2 on pages HL3.3 and HL3.4 in Statistical Highlight #3.




NOTES: 7. As summarized on the left of the schema at the right, a relationship in *statistics* is often considered in terms of one or more of association, confounding, causation, interaction and Simpson's Paradox.

In *probability* (on the right of the schema), a relationship is considered in terms of *dependence*, which comes in great variety and is often difficult to mathematize; as a consequence, introductory courses emphasize *probabilistic independence*, as it applies to events, random variables and processes. Even the first two of these three involve an appreciable set of ideas and may be all a course has time to discuss.

Connection between statistical and probabilistic considerations of a relationship arises in the probability *models* statistics uses in the Analysis stage of the FDEAC cycle.

- Emphasis on probabilistic *independence* in introductory courses can obscure the fact that it is a mathematical *idealization*. In the real world, *dependence* is the norm – it may be that the behaviour of *every* particle in the universe depends on (*i.e.*, is affected by) *every other* particle, no matter how minute the degree of dependence.
 - This may be why lurking variates are usually so *numerous* when answering Questions with a causative aspect.

The schema at the lower right of page HL57.1 in Statistical Highlight #57, which provides a framework for our discussion of data-based investigating of statistical relationships, is shown again at the right; *unsurprisingly*, it is a more detailed version of the left-hand ('statistics') side of the schema above it.



8. The (equivalent) diagrams at the right show the effects of two (binary) focal variates X_1 and X_2 on (the average of) Y ; one focal variate is on the horizontal axis of a diagram, the other distinguishes the two lines by its level.

- The *nonparallel* lines show there is an X_1 - X_2 *interaction*.
- The left-hand diagram shows that X_1 and Y are *independent* when X_2 is Lo (the relevant line has *zero* slope) but *not* when X_2 is Hi.
- The right-hand diagram shows there is *no* independence of X_2 and Y – *neither* line has zero slope.

For these two diagrams, it is *wrong* to involve *conditional (probabilistic) independence* because it implies that we have undertaken the difficult task of formulating a probability model for this situation – see also Section 13 on page HL89.18 in Statistical Highlight #89.

Thus, different variate contexts can yield different assessments of (complicated) relationships.

9. Table HL51.7 on page HL51.3 illustrates, for *averages* of a *continuous* response variate, a phenomenon analogous to Simpson's Paradox. A similar 'paradox' occurs for an organization of (say) 10 employees who each work for the organization for (say) 30 years or until age 55, whichever comes first. In constant dollars (*i.e.*, with the effect of inflation removed), each employee is hired at a salary determined by their experience, taken as proportional to their age which is at least 25; each receives the *same* salary *increase* each year. At start-up, the organization hires people with a variety of levels of experience/age but a steady state is then maintained by hiring a new employee aged 25 to replace a person leaving the organization at age 55. This situation is portrayed graphically at the right above for the first 60 years of the organization's existence – each stepped line represents one employee's salary over time. Despite every employee's *increasing* salary, the organization's employee experience/age structure (a 'lurking variate') and, hence, the *average* salary for the organization, remain nearly *constant* over time.

Although the organization described is an idealization, it is reasonable model for a college or university with a constant faculty complement. The 'paradox' could result in a specious claim, at the time of salary negotiations, that faculty salaries are being unfairly constrained because the *average* salary is not increasing over time.

