

§8.2 - Integration by Parts (IBP)

Recall: If u and v are functions of x , then by the product rule,

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

Integrating both sides, we have

$$\int \frac{d}{dx}(uv) dx = \int u \frac{dv}{dx} dx + \int v \frac{du}{dx} dx$$
$$\implies uv = \int u dv + \int v du$$

Rearrange

\implies

$$\int u dv = uv - \int v du.$$

↑ Integration by parts (IBP) formula.

Remarks:

1. IBP allows us to trade one integral, $\int u dv$ for another (hopefully simpler) integral, $\int v du$.

2. IBP can help when integrating a product of functions. One function, u , will be differentiated while the other, dv , will be integrated.

Ex: $\int x \cos x \, dx$

Strategy: Pick u to be a function that gets simpler when differentiated. Everything else is dv .

Solution: Let's define u and dv as follows:

$$\begin{array}{ccc} u = x & & v = \sin x \\ \text{differentiate} \downarrow & & \uparrow \text{integrate} \\ du = 1 \cdot dx & & dv = \cos x \, dx \end{array}$$

Then

$$\int \frac{x}{u} \frac{\cos x \, dx}{dv} = uv - \int v \, du$$

$$= x \sin x - \int \sin x \, dx$$

$\boxed{= -\cos x + C}$

$$= \boxed{x \sin x + \cos x + C}$$

A better strategy for picking u is the LIATE method.

Logs
 Inverse trig
 Algebraic (powers of x)
 Trig
 Exponential

Let u be the first function on
 this list that appears in your
 integral. Let everything else
 be dv .

Ex: Evaluate $\int x^2 \ln x \, dx$

Solution: Using IBP and the LIATE method,

$$u = \ln x \quad v = x^3/3$$

$$du = \frac{1}{x} \, dx \quad dv = x^2 \, dx$$

$$\begin{aligned}\therefore \int x^2 \ln x \, dx &= uv - \int v \, du \\ &= (\ln x) \cdot \frac{x^3}{3} - \int \frac{x^3}{3} \cdot \frac{1}{x} \, dx\end{aligned}$$

$$= \frac{x^3}{3} \ln x - \frac{1}{3} \int x^2 dx$$

$$= \boxed{\frac{x^3}{3} \ln x - \frac{x^3}{9} + C}$$

Ex: Evaluate $\int x^2 e^x dx$

Solution: Let $u = x^2$ $v = e^x$
 $du = 2x dx$ $dv = e^x dx$

$$\begin{aligned} \int x^2 e^x dx &= x^2 e^x - \int e^x \cdot 2x dx \\ &= x^2 e^x - 2 \int x e^x dx \\ u = x &\quad v = e^x \\ du = dx &\quad dv = e^x dx \end{aligned}$$

IBP again!

$$\begin{aligned} &= x^2 e^x - 2 \left[x e^x - \int e^x dx \right] \\ &= \boxed{x^2 e^x - 2x e^x + 2e^x + C} \end{aligned}$$

Sometimes we may need to combine multiple methods.

Ex: Evaluate $\int \sin x \cos x e^{\sin x} dx$

Solution: Let's first clean up the integral using a

u -substitution.

$$\begin{aligned} \int \sin x \cos x e^{\sin x} dx &= \int \underbrace{\sin x \cdot e^{\sin x}}_{=ue^u} \cdot \underbrace{\cos x dx}_{=du} \\ \text{Let } u = \sin x, \text{ so } du &= \cos x dx \\ &= \int ue^u du &= wv - \int v dw \quad \text{where} \\ &\qquad\qquad\qquad \text{IBP} \\ &= ue^u - \int e^u du &w=u & v=e^u \\ &&dw=du & dv=e^u du \\ &= ue^u - e^u + C \\ &= \sin x e^{\sin x} - e^{\sin x} + C \end{aligned}$$

Sometimes IBP can also help when the integrand

doesn't look like a product!

Ex: $\int \ln x dx$

Solution: Let $u = \ln x$ $v = x$

$$du = \frac{1}{x} dx \quad dv = 1 dx$$

Then $\int \ln x dx = (\ln x) \cdot x - \int x \cdot \frac{1}{x} dx$

$$= x \ln x - \int 1 dx$$
$$= \boxed{x \ln x - x + C}$$

Ex: Evaluate $\int \arctan x dx$

Solution: Let $u = \arctan x$ $v = x$

$$du = \frac{1}{1+x^2} dx \quad dv = 1 dx$$

Then

$$\int \arctan x dx = x \cdot \arctan x - \int \frac{x}{1+x^2} dx$$

u-substitution!
Let $u = 1+x^2$, so
 $du = 2x dx \Rightarrow dx = \frac{du}{2x}$

$$= x \cdot \arctan x - \int \frac{x}{u} \left(\frac{du}{2x} \right)$$
$$= x \cdot \arctan x - \frac{1}{2} \int \frac{1}{u} du$$
$$= x \cdot \arctan x - \frac{1}{2} \ln|u| + C$$

$$= x \cdot \arctan x - \frac{1}{2} \ln |1+x^2| + C$$

Here's one more interesting type of IBP problem...

Ex: Evaluate $\int e^x \sin x \, dx$

Solution: Use IBP with $u = \sin x$ $v = e^x$
 $du = \cos x \, dx$ $dv = e^x \, dx$

$$\begin{aligned} \int e^x \sin x \, dx &= e^x \sin x - \int e^x \cos x \, dx \quad \text{IBP again!} \\ &\qquad\qquad\qquad u = \cos x \qquad v = e^x \\ &\qquad\qquad\qquad du = -\sin x \, dx \qquad dv = e^x \, dx \\ &= e^x \sin x - \left[e^x \cos x - \int e^x (-\sin x) \, dx \right] \\ &= e^x \sin x - e^x \cos x - \underbrace{\int e^x \sin x \, dx}_{\text{Aha! This is exactly the integral we started with!}} \end{aligned}$$

If $I = \int e^x \sin x \, dx$, then we've just shown that

$$I = e^x \sin x - e^x \cos x - I,$$

hence $2I = e^x (\sin x - \cos x)$, and therefore

$$I = \int e^x \sin x \, dx = \frac{e^x(\sin x - \cos x)}{2} + C$$

Using integration by parts with definite integrals is

no different — just don't forget the bounds!

$$\int_a^b u \, dv = [uv]_a^b - \int_a^b v \, du$$

Ex: Evaluate $\int_1^e x \ln x \, dx$.

Solution: Let $u = \ln x \quad v = \frac{x^2}{2}$

$$du = \frac{1}{x} \, dx \quad dv = x \, dx$$

$$\begin{aligned} \int_1^e x \ln x \, dx &= \left[\frac{x^2}{2} \ln x \right]_1^e - \int_1^e \frac{x^2}{2} \cdot \frac{1}{x} \, dx \\ &= \left(\frac{e^2}{2} \underbrace{\ln e}_{=1} - \frac{1^2}{2} \underbrace{\ln 1}_{=0} \right) - \int_1^e \frac{x^2}{2} \, dx \\ &= \frac{e^2}{2} - \left[\frac{x^3}{6} \right]_1^e \\ &= \frac{e^2}{2} - \left(\frac{e^3}{6} - \frac{1^3}{6} \right) = \boxed{\frac{e^2}{4} + \frac{1}{4}} \end{aligned}$$