

⑤ The Ratio and Root Tests

Our final two convergence tests can, in some cases, tell us that a series converges absolutely.

The Ratio Test

Consider a series $\sum a_n$ and suppose $L = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$ exists or is ∞ .

- (i) If $L < 1$, $\sum a_n$ converges absolutely
- (ii) If $L > 1$, $\sum a_n$ diverges
- (iii) If $L = 1$, the test is inconclusive. The series could converge absolutely, conditionally, or diverge.

Remark: The ratio test is often effective when dealing

with factorials: We define $0! = 1$ and

$$n! = n(n-1)(n-2) \cdots 3 \cdot 2 \cdot 1 \quad (n \in \mathbb{N})$$

$$\text{e.g., } 5! = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 120$$

Ex: Determine whether each series below converges

absolutely, converges conditionally, or diverges.

(a) $\sum_{n=0}^{\infty} \frac{(-1)^n \cdot 3^n}{n!}$

Solution: We'll use the ratio test!

$$\begin{aligned} L &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+1} \cdot 3^{n+1}}{(n+1)!}}{\frac{(-1)^n \cdot 3^n}{n!}} \right| \\ &= \lim_{n \rightarrow \infty} \frac{3^{n+1}}{3^n} \cdot \frac{n!}{(n+1)!} = \frac{n!}{(n+1) \cdot n!} = \frac{1}{n+1} \\ &= \lim_{n \rightarrow \infty} 3 \cdot \frac{1}{n+1} = 0 \end{aligned}$$

Since $L < 1$, $\sum_{n=0}^{\infty} \frac{(-1)^n 3^n}{n!}$ converges absolutely by

the ratio test.

(b) $\sum_{n=1}^{\infty} \frac{9^n}{n \cdot 2^{n+1}}$

Solution: Again, let's try the ratio test!

$$\begin{aligned}
 L &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{\frac{q^{n+1}}{(n+1) \cdot 2^{n+2}}}{\frac{q^n}{n \cdot 2^{n+1}}} \right| \\
 &= \lim_{n \rightarrow \infty} \frac{q^{n+1}}{q^n} \cdot \frac{2^{n+1}}{2^{n+2}} \cdot \frac{n}{n+1} \\
 &= \lim_{n \rightarrow \infty} \frac{q}{2} \left(\frac{n}{n+1} \right) \stackrel{LH}{=} \frac{q}{2}
 \end{aligned}$$

Since $L > 1$, $\sum_{n=1}^{\infty} \frac{q^n}{n \cdot 2^{n+1}}$ diverges by the ratio test.

$$(c) \sum_{n=2}^{\infty} \frac{(-1)^n}{n^2 \cdot \ln(n)}$$

Solution: Maybe the ratio test will work again?

$$\begin{aligned}
 L &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+1}}{(n+1)^2 \ln(n+1)}}{\frac{(-1)^n}{n^2 \cdot \ln(n)}} \right| \\
 &= \lim_{n \rightarrow \infty} \frac{n^2}{(n+1)^2} \cdot \frac{\ln(n)}{\ln(n+1)}
 \end{aligned}$$

$$\begin{aligned}
 &= \underbrace{\lim_{n \rightarrow \infty} \left(\frac{n}{n+1} \right)^2}_{\stackrel{\text{L'H}}{=} \left(\lim_{n \rightarrow \infty} \frac{1}{1} \right)^2 = 1} \cdot \underbrace{\lim_{n \rightarrow \infty} \frac{\ln(n)}{\ln(n+1)}}_{\stackrel{\text{L'H}}{=} \lim_{n \rightarrow \infty} \frac{\frac{1}{n}}{\frac{1}{n+1}}} = \underline{\underline{1}} \\
 &= \lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right) = 1
 \end{aligned}$$

Uh oh... if $L = 1$, the ratio test is inconclusive!

To determine if $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2 \cdot \ln(n)}$ converges absolutely,

we'll need to examine

$$\sum_{n=2}^{\infty} \left| \frac{(-1)^n}{n^2 \cdot \ln(n)} \right| = \sum_{n=2}^{\infty} \frac{1}{n^2 \cdot \ln(n)} .$$

Note that $\frac{1}{n^2 \cdot \ln(n)} \leq \frac{1}{n^2}$ and $\sum_{n=2}^{\infty} \frac{1}{n^2}$ is a

convergent p-series (as $p=2 > 1$). Thus, $\sum_{n=2}^{\infty} \frac{1}{n^2 \cdot \ln(n)}$

Converges, so $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2 \cdot \ln(n)}$ Converges absolutely.

The Root Test

Consider a series $\sum a_n$ and suppose $L = \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}$ exists or is ∞ .

(i) If $L < 1$, $\sum a_n$ converges absolutely

(ii) If $L > 1$, $\sum a_n$ diverges

(iii) If $L = 1$, the test is inconclusive. The series

could converge absolutely, conditionally, or diverge.

Remark: The root test is often effective when dealing with series involving expressions like $(f(n))^\frac{1}{n}$.

Ex: Apply the root test to $\sum_{n=1}^{\infty} \left(\frac{n+1}{3n+1} \right)^n$.

Solution: $L = \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}$

$$= \lim_{n \rightarrow \infty} \sqrt[n]{\left| \left(\frac{n+1}{3n+1} \right)^n \right|}$$

$$= \lim_{n \rightarrow \infty} \frac{n+1}{3n+1}$$

$$= \frac{1}{3}$$

Since $L < 1$, the series converges absolutely by the root test.

Ex: Apply the root test to $\sum_{n=1}^{\infty} \frac{(-1)^n e^{n^2}}{n^n}$.

$$\begin{aligned}\text{Solution: } L &= \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \lim_{n \rightarrow \infty} \sqrt[n]{\left| \frac{(-1)^n e^{n^2}}{n^n} \right|} \\ &= \lim_{n \rightarrow \infty} \left(\frac{e^{n^2}}{n^n} \right)^{1/n} \\ &= \lim_{n \rightarrow \infty} \frac{e^n}{n} \\ &\stackrel{LH}{=} \lim_{n \rightarrow \infty} \frac{e^n}{1} = \infty.\end{aligned}$$

Thus, since $L > 1$, the series diverges by the root test.