

§5.2 - Geometric Series

A geometric series is a series of the form

$$\sum_{n=0}^{\infty} a \cdot r^n = a + ar + ar^2 + ar^3 + \dots$$

where a, r are constants. We refer to r as the common ratio.

Ex: $\sum_{n=0}^{\infty} 2 \cdot \left(\frac{3}{2}\right)^n = 2 + 2\left(\frac{3}{2}\right) + 2\left(\frac{3}{2}\right)^2 + \dots$

This is a geometric series with $a=2$, $r = \frac{3}{2}$.

Ex: $\sum_{n=0}^{\infty} 3\left(\frac{-1}{5}\right)^n = 3 - \frac{3}{5} + \frac{3}{25} - \frac{3}{125} + \dots$

This is a geometric series with $a=3$, $r = -\frac{1}{5}$.

Question: When does a geometric series $\sum_{n=0}^{\infty} ar^n$ converge? When does it diverge?

Answer: Let's consider a few cases...

Case I: $r = 1$

We have $\sum_{n=0}^{\infty} ar^n = \underbrace{a + a + a + \dots}_{\text{sum blows up!}} \Rightarrow \text{divergent!}$

Case II: $r = -1$

We have $\sum_{n=0}^{\infty} ar^n = \underbrace{a - a + a - a + \dots}_{\text{sum never stabilizes!}} \Rightarrow \text{divergent}$

Case III: $r \neq \pm 1$

Let's examine the partial sum S_N . We have

$$\textcircled{1} \quad S_N = a + \cancel{ar} + \cancel{ar^2} + \dots + \cancel{ar^N}$$

$$\textcircled{2} \quad \underline{r \cdot S_N = \cancel{ar} + \cancel{ar^2} + \dots + \cancel{ar^N} + ar^{N+1}}$$

$$\textcircled{1} - \textcircled{2} \quad S_N - r \cdot S_N = a - ar^{N+1}$$

$$\Rightarrow (1-r) S_N = a(1-r^{N+1})$$

$$\Rightarrow \boxed{S_N = \frac{a(1-r^{N+1})}{1-r}}$$

To check convergence, we consider $\lim_{N \rightarrow \infty} S_N$:

If $r > 1$ or $r < -1$, then r^{N+1} blows up as $N \rightarrow \infty$;

but if $-1 < r < 1$, then $r^{N+1} \rightarrow 0$ as $N \rightarrow \infty$. Hence,

$$\lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \frac{a(1 - r^{N+1})}{1-r} = \begin{cases} \frac{a}{1-r} & \text{if } |r| < 1 \\ \text{DNE} & \text{if } |r| > 1. \end{cases}$$

The Geometric Series Test

Consider the geometric series $\sum_{n=0}^{\infty} ar^n$.

(i) If $|r| < 1$, then the series converges. In

particular, $\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$

(ii) If $|r| \geq 1$, then the series diverges.

Ex: $\sum_{n=0}^{\infty} 2 \cdot \left(\frac{3}{2}\right)^n$ is geometric with $a=2$, $r=3/2$.

Since $|r| \geq 1$, this series diverges!

Ex: $\sum_{n=0}^{\infty} 3 \left(\frac{-1}{5}\right)^n$ is geometric with $a=3$, $r=-1/5$.

Since $|r| < 1$, this series converges. Specifically,

$$\sum_{n=0}^{\infty} 3\left(\frac{-1}{5}\right)^n = \frac{a}{1-r} = \frac{3}{1-(-1/5)} = \frac{3}{6/5} = \boxed{\frac{5}{2}}$$

Example: Let's use geometric series to show that

$$0.99999\dots = 1!$$

Solution: $0.99999\dots = 0.9 + 0.09 + 0.009 + \dots$

$$= 9\left(\frac{1}{10}\right) + 9\left(\frac{1}{10}\right)^2 + 9\left(\frac{1}{10}\right)^3 + \dots$$

$$= \sum_{n=1}^{\infty} 9\left(\frac{1}{10}\right)^n$$

This is a geometric series with $r = \frac{1}{10}$. Since $|r| < 1$,

the series converges. However, the sum is NOT

$$\frac{a}{1-r} = \frac{9}{1-\frac{1}{10}} = \frac{9}{9/10} = 10$$

The formula $\sum_{n=n_0}^{\infty} a \cdot r^n = \frac{a}{1-r}$ only works if $n_0 = 0$!

We have a couple options to get around this...

Option 1: Add and subtract terms to create a sum starting at $n_0 = 0$.

$$\begin{aligned}\sum_{n=1}^{\infty} 9 \left(\frac{1}{10}\right)^n &= \underbrace{\sum_{n=1}^{\infty} 9 \left(\frac{1}{10}\right)^n + 9 \left(\frac{1}{10}\right)^0}_{\text{Geometric, } a=9, r=\frac{1}{10}} - 9 \left(\frac{1}{10}\right)^0 \\ &= \sum_{n=0}^{\infty} 9 \left(\frac{1}{10}\right)^n - 9 \left(\frac{1}{10}\right)^0 \\ &= \frac{9}{1 - \frac{1}{10}} - 9 \\ &= 10 - 9 \\ &= \boxed{1}\end{aligned}$$

Option 2: Reindex the sum to start at $n = 0$.

$$\sum_{n=1}^{\infty} 9 \left(\frac{1}{10}\right)^n = \sum_{n=0}^{\infty} 9 \left(\frac{1}{10}\right)^{n+1}$$

← Increased by 1 to make up for it!

↑ Reduced by 1

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \left(\frac{9}{10}\right) \left(\frac{1}{10}\right)^n \quad \leftarrow \begin{array}{l} \text{Geometric,} \\ a = \frac{9}{10}, r = \frac{1}{10} \end{array} \\
&= \frac{9/10}{1 - \frac{1}{10}} \\
&= \frac{9/10}{9/10} \\
&= \boxed{1}
\end{aligned}$$

Ex: $\sum_{n=2}^{\infty} \frac{3^{n+1}}{2^{2n}}$ doesn't yet look like a geometric

series ... but let's rewrite it as

$$\sum_{n=2}^{\infty} \frac{3^{n+1}}{2^{2n}} = \sum_{n=2}^{\infty} \frac{3 \cdot 3^n}{(2^2)^n} = \underbrace{\sum_{n=2}^{\infty} 3 \left(\frac{3}{4}\right)^n}_{\text{Geometric with } r = 3/4}$$

Since $|r| < 1$, the series converges. In particular,

$$\begin{aligned}
\sum_{n=2}^{\infty} \frac{3^{n+1}}{2^{2n}} &= \sum_{n=2}^{\infty} 3 \left(\frac{3}{4}\right)^n \\
&= \sum_{n=0}^{\infty} 3 \left(\frac{3}{4}\right)^{n+2}
\end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} 3 \cdot \left(\frac{3}{4}\right)^2 \cdot \left(\frac{3}{4}\right)^n \\
&= \sum_{n=0}^{\infty} \frac{27}{16} \cdot \left(\frac{3}{4}\right)^n \\
&= \frac{27/16}{1 - 3/4} = \boxed{\frac{27}{4}}
\end{aligned}$$

Ex: Use geometric series to write $2.81313\overline{13}$ as a fraction.

Solution:

$$2.81313\overline{13} = 2.8 + (0.013 + 0.00013 + 0.0000013 + \dots)$$

$$= 2.8 + 0.013 \underbrace{\left(1 + \frac{1}{100} + \frac{1}{100^2} + \dots\right)}_{\text{Geometric, } a=1, r = \frac{1}{100}}$$

$$= \frac{28}{10} + \frac{13}{1000} \left(\frac{1}{1 - \frac{1}{100}}\right)$$

$$= \frac{28}{10} + \frac{13}{1000} \left(\frac{100}{99}\right)$$

$$= \frac{28 \cdot 99}{10 \cdot 99} + \frac{13}{990} = \frac{2772 + 13}{990} = \boxed{\frac{2785}{990}}$$