

TEST	STATEMENT	NOTES	EXAMPLES
Geometric Series Test	If $a, r \in \mathbb{R}$ with $a \neq 0$ then the geometric series $\sum_{n=0}^{\infty} ar^n = \begin{cases} \frac{a}{1-r} & \text{if } r < 1, \\ \text{divergent} & \text{if } r \geq 1. \end{cases}$	- The N^{th} partial sum is given by $S_N = \frac{a(1-r^{N+1})}{1-r}.$	$\sum_{n=0}^{\infty} \frac{(-3)^n}{4^n}$ $\sum_{n=1}^{\infty} \frac{2^{2n}}{5^{n+1}}$
Divergence Test	If $\lim_{n \rightarrow \infty} a_n \neq 0$ or $\lim_{n \rightarrow \infty} a_n$ DNE, then $\sum_{n=1}^{\infty} a_n$ diverges.	- Often a good test to start with. - If $\lim_{n \rightarrow \infty} a_n = 0$, no conclusions can be made. (e.g., $\sum \frac{1}{n}$ diverges and $\sum \frac{1}{n^2}$ converges.)	$\sum_{n=1}^{\infty} \frac{n-1}{3n-1}$ $\sum_{n=1}^{\infty} \cos\left(\frac{1}{n}\right)$
Integral Test	Suppose $f(x)$ is continuous, positive, and decreasing on $[1, \infty)$. (i) If $\int_1^{\infty} f(x) dx$ converges, then $\sum_{n=1}^{\infty} f(n)$ converges. (ii) If $\int_1^{\infty} f(x) dx$ diverges, then $\sum_{n=1}^{\infty} f(n)$ diverges.	- Useful when $\int_1^{\infty} f(x) dx$ is easy to calculate. - When convergent, we have the remainder estimate $\int_{N+1}^{\infty} f(x) dx \leq R_N \leq \int_N^{\infty} f(x) dx$ where $R_N = S - S_N$ and S is the sum of the series.	$\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$ $\sum_{n=1}^{\infty} \frac{e^{1/n}}{n^2}$
p-Series Test	The series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges when $p > 1$ and diverges when $p \leq 1$.	- Often used with comparison tests.	$\sum_{n=1}^{\infty} \frac{1}{(2n)^2}$ $\sum_{n=1}^{\infty} \frac{1}{n}$
Comparison Test	Suppose that $0 \leq a_n \leq b_n$ for all n sufficiently large. (i) If $\sum b_n$ converges, then $\sum a_n$ converges. (ii) If $\sum a_n$ diverges, then $\sum b_n$ diverges.	- No conclusions if $\sum b_n$ diverges or $\sum a_n$ converges.	$\sum_{n=1}^{\infty} \frac{n+2}{(n+1)^3}$ $\sum_{n=1}^{\infty} \frac{\sqrt[3]{n}}{\sqrt{n^3+n+3}}$

TEST	STATEMENT	NOTES	EXAMPLES
<p>Limit Comparison Test</p>	<p>Suppose that $\sum a_n$ and $\sum b_n$ are series of positive terms, and let</p> $L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n}.$ <p>(i) If L exists and $0 < L < \infty$, then $\sum a_n$ and $\sum b_n$ either both converge or both diverge.</p> <p>(ii) If $L = 0$ and $\sum b_n$ converges, then $\sum a_n$ converges.</p> <p>(iii) If $L = \infty$ and $\sum b_n$ diverges, then $\sum a_n$ diverges.</p>	<p>- Usually works well with fractions involving polynomials, roots, or exponentials.</p> <p>- When applying this test to $\sum a_n$, we usually define b_n using only the most dominant parts of a_n.</p>	$\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt[3]{5 + n^7}}$ $\sum_{n=1}^{\infty} \frac{2^n + 3^n}{4^n + 5^n}$
<p>Alternating Series Test</p>	<p>Consider the series</p> $\sum_{n=1}^{\infty} (-1)^{n+1} b_n = b_1 - b_2 + b_3 - b_4 + \dots,$ <p>where $b_n > 0$ for all n. If</p> <p>(i) $\{b_n\}$ is a decreasing sequence, and</p> <p>(ii) $\lim_{n \rightarrow \infty} b_n = 0$,</p> <p>then $\sum_{n=1}^{\infty} (-1)^{n+1} b_n$ converges.</p>	<p>- When convergent, we have the remainder estimate</p> $ S - S_N \leq b_{N+1},$ <p>where S is the sum of the series.</p>	$\sum_{n=0}^{\infty} (-1)^n \frac{\sqrt{n}}{2n + 3}$ $\sum_{n=1}^{\infty} (-1)^n \sin\left(\frac{\pi}{n}\right)$ $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^{n!}}$
<p>Ratio Test</p>	<p>Suppose that $L = \lim_{n \rightarrow \infty} \left \frac{a_{n+1}}{a_n} \right$ exists or is equal to ∞.</p> <p>(i) If $L < 1$, then $\sum a_n$ converges absolutely.</p> <p>(ii) If $L > 1$, then $\sum a_n$ diverges.</p> <p>(iii) If $L = 1$, the test is inconclusive.</p>	<p>- Useful when the terms of the series involve factorials.</p>	$\sum_{n=1}^{\infty} \frac{10^n}{n \cdot 4^{2n+1}}$ $\sum_{n=1}^{\infty} \frac{(2n)!}{n! 2^n}$ $\sum_{n=1}^{\infty} \frac{(-1)^n \ln(n)}{3^n}$
<p>Root Test</p>	<p>Suppose that $L = \lim_{n \rightarrow \infty} \sqrt[n]{ a_n }$ exists or is equal to ∞.</p> <p>(i) If $L < 1$, then $\sum a_n$ converges absolutely.</p> <p>(ii) If $L > 1$, then $\sum a_n$ diverges.</p> <p>(iii) If $L = 1$, the test is inconclusive.</p>	<p>- Useful when terms of the series involve n^{th} powers.</p>	$\sum_{n=1}^{\infty} (\tan^{-1} n)^n$ $\sum_{n=1}^{\infty} \frac{n^n}{n^3 e^n}$ $\sum_{n=1}^{\infty} \left(\frac{n+1}{2n+1} \right)^{2n}$